

LIST OF TABLES

Table	Page
2-1 Plantation areas of Boakaew Watershed Management Station	14
2-2 Growths of <i>Pinus kesiya</i> in a series of pine plantations of 21 age-class stands	18
2-3 Wood productions of <i>Pinus kesiya</i> in a series of plantations of 21 age-class stands	21
2-4 Timber volumes of <i>Pinus kesiya</i> with different timber quality and stem-girth classes in 21 age-class plantations	22
2-5 Growths of succession trees in a series of plantations of 21 age-class stands	23
2-6 Wood productions of succession trees in a series of pine plantations of 21 age-class stands	26
2-7 Timber volumes of succession trees with different timber quality and stem-girth classes in 21 age-class plantations	27
2-8 Quantitative characteristics of tree species in a series of pine plantations	47
2-9 Species diversity index (SWI) of tree species in a series of pine plantations	68
2-10 The number of tree individuals of different height classes in a series of pine plantations	89
2-11 The number of tree individuals of different stem-girth classes in a series of pine plantations	110
2-12 Total number of seedling and ground-covered species in a series of pine plantations	132
2-13 Overall quantitative data of plant communities in a series of <i>Pinus kesiya</i> plantations	137
3-1 Species list, growth forms and seed/fruit characteristics of tree species in fragmented forests	144
3-2 Quantitative characteristics of tree species in fifteen fragmented forests	149
3-3 Species diversity index (SWI) of tree species in fifteen fragmented forests	151
3-4 Indexes of forest conditions (FCI) in fifteen fragmented forests	154
3-5 Number of tree individuals with different height classes in fifteen fragmented forests	155
3-6 Number of tree individuals with different height classes in each fragmented forest	157
3-7 Number of tree individuals with different stem-girth classes in fifteen fragmented forests	169
3-8 Number of tree individuals with different stem-girth classes in each fragmented forest	171
3-9 Quantitative characteristics of tree species in fifteen fragmented forests	185
3-10 Species diversity index (SWI) of tree species in each fragmented forest	196

LIST OF TABLES (CONTINUED)

Table	Page
3-11 Quantitative characteristics of tree species in clusters	212
3-12 Overall data of plant communities in fifteen fragmented forests	213
3-13 Total numbers of seedling and ground-covered species in fifteen fragmented forests	215
3-14 Species richness among fragmented forest and pine plantations	221
4-1 Some physical properties in soil profiles under pine plantations	228
4-2 Some chemical properties in soil profiles under pine plantations	235
4-3 Extractable bases and acidity, cation exchange capacity and base saturation in soil profiles under pine plantations	238
4-4 Assessment of fertility levels in soil profiles under pine plantations	239
4-5 Overall data of plant communities in pine plantations	241
4-6 Topography and soil profile development in five fragmented forests	242
4-7 Some physical properties in soil profiles under fragmented forests	260
4-8 Some chemical properties in soil profiles under fragmented forests	267
4-9 Extractable bases and acidity, cation exchange capacity and base saturation in soil profiles under fragmented forests	270
4-10 Assessment of fertility levels in soil profiles under fragmented forests	271
4-11 Overall data of plant communities in five fragmented forests	273
4-12 Some physical properties in soil profile under opened area	274
4-13 Some chemical properties in soil profile under opened area	280
4-14 Extractable bases and acidity, cation exchange capacity and base saturation in soil profile under opened area	281
4-15 Assessment of fertility levels in a soil profile under opened area	281

LIST OF FIGURES

Figure		Page
2-1	The location of study area	15
2-2	The relation between tree height and dbh of <i>Pinus kesiya</i> in pine plantations	19
2-3	The relation between tree volume and the product of the square of dbh (D^2) and tree height (H) of <i>Pinus kesiya</i> by using power equation	19
2-4	Growth models of tree height, dbh and stem volume of <i>Pinus kesiya</i> of 21 age-class stands	20
2-5	Mean volumes and volume increments of <i>Pinus kesiya</i> of 21 age-class stands	22
2-6	Mean height and dbh of <i>Pinus kesiya</i> and other broad-leaved trees in 21 age-class plantations	24
2-7	Mean height and dbh increments of <i>Pinus kesiya</i> and other broad-leaved trees in 21 age-class plantations	24
2-8	Over view of pine plantations at Boakaew Watershed Management Station	25
2-9	The natural succession in pine plantations	25
2-10	Tree densities of <i>Pinus kesiya</i> and succession broad-leaved trees with different stem-girth classes in 21 age-class plantations	46
2-11	Natural regeneration in <i>Pinus kesiya</i> plantations	131
3-1	The relation between dbh and tree height in fragmented forests	153
3-2	The size class distribution of gbh in fragmented forests	153
3-3	Over views pictures of fragmented forests	184
3-4	Similarity percentages of plant communities among fragmented forests	208
3-5	Spatial distribution of dominant tree species in fragmented forests located on different altitudes	209
3-6	Human disturbances through tree cutting in fragmented forests	211
3-7	Dendrogram for the cluster analysis of fragmented lower montane forests of Boakaew highland watershed. The horizontal dash line represents reference points for delimiting cluster 1, 2, 3 and 4.	212
3-8	Some seedlings and ground-covered species in fragmented forests	214
3-9	Modes of fruits/seeds dispersal	220
4-1	Variations of bulk densities (left) and gravel amounts (right) along soil profiles under pine plantations	229
4-2	Distribution of soil particles along soil profiles under pine plantations	229
4-3	Comparison of bulk densities and gravel amounts in surface soils (0-10 cm) of five age class pine plantations	230
4-4	The pH values and contents of soil organic matter, carbon, nitrogen and other extractable nutrients in soil profiles under pine plantations	236
4-5	Comparison of soil reaction in surface soils (0-10 cm) among different age-class pine plantations	237
4-6	C/N ratios, cation exchange capacity (CEC) and base saturation percentages (BS) in soil profiles under pine plantations	237

LIST OF FIGURES (CONTINUED)

Figure	Page
4-7 Study site and soil profile of pedon 1 (The 1 st fragmented forest)	253
4-8 Study site and soil profile of pedon 2 (The 2 nd fragmented forest)	254
4-9 Study site and soil profile of pedon 3 (The 3 rd fragmented forest)	255
4-10 Study site and soil profile of pedon 4 (The 4 th fragmented forest)	256
4-11 Study site and soil profile of pedon 5 (The 5 th fragmented forest)	257
4-12 Variations of bulk densities (left) and gravel amounts (right) in soil profiles under fragmented forests	261
4-13 Distribution of soil particles in soil profiles under fragmented forests	261
4-14 The pH values and contents of organic matter, carbon, nitrogen, and other extractable nutrients in soil profiles under fragmented forests	268
4-15 C/N ratios, cation exchange capacity (CEC) and base saturation percentages (BS) in soil profiles under fragmented forests	269
4-16 Variations of bulk densities (left) and gravel amounts (right) in soil profiles of pine plantations, opened area and the 5 th fragmented forest	274
4-17 Comparison of bulk densities and gravel amounts in surface soils (0-5 and 5-10 cm) among pine plantations, opened area and the 5 th fragmented forest	275
4-18 Distribution of soil particles along profile in opened area	275
4-19 The pH values and contents of organic matter, carbon, nitrogen and extractable nutrients in soil profiles of pine plantations, opened area and the 5 th fragmented forest	278
4-20 C/N ratio, cation exchange capacity (CEC) and base saturation percentage (BS) in soil profiles of pine plantations, opened area and the 5 th fragmented forest	279