

เอกสารอ้างอิง

- เนาวรัตน์ ศิริศิลป์. 2527. การปฏิบัติการวิเคราะห์คุณภาพและพืช. ภาควิชาปฐมศาสตร์และอนุรักษศาสตร์, คณะเกษตรศาสตร์, มหาวิทยาลัยเชียงใหม่. เชียงใหม่. 108-112 หน้า.
- ปรีชา วงศ์ศิริศักดิ์, วิเชียร ภานุมาส, พรภพ เชาวสันต์, ชนากยุ คลสมบูรณ์, นันทกร บุญเกิด และ เย็นใจ วงศ์สัก. 2529. การคัดเลือกสายพันธุ์ราชบูรณะ ถัวเหลืองที่เหมาะสมสำหรับถัวเหลืองพันธุ์ จังหวัดเชียงใหม่ 81 ในสภาพท้องที่. เอกสารวิชาการค้านปฐวิทยา เล่มที่ 1, การประชุมวิชาการประจำปี 2529. กองปฐวิทยา, กรมวิชาการเกษตร.
- ศรีศุกร์ นิลารณ์. 2532. ราชบูรณะสายพันธุ์พื้นเมืองสำหรับถัวเหลืองที่ดีในภาคเหนือของประเทศไทย. วิทยานิพนธ์เพื่อปริญญาวิทยาศาสตรมหาบัณฑิต สาขาวิชาปฐมศาสตร์, มหาวิทยาลัยเชียงใหม่, เชียงใหม่. 176 หน้า.
- ศูนย์สหกิจการเกษตร. 2531. สหกิจการเกษตรของประเทศไทย ปีเพาะปลูก 2530/31. เอกสารสหกิจการเกษตร เลขที่ 408. สำนักงานเศรษฐกิจการเกษตร, กระทรวงเกษตรและสหกรณ์, กรุงเทพมหานคร. 276 หน้า.
- Abu-Shakra, S.S., D.A. Phillips and R.C. Huffaker. 1978. Nitrogen fixation and delayed leaf senescence in soybeans. Science. 199 : 973-975.
- Boonkerd. N, 1982. Evaluation of Rhizobium japonicum in field grown soybeans. Proceedings of National Workshop in Recent Advances in Nitrogen Fixation Research : Their Implication for Thailand. Chulalongkorn University, Bangkok. pp 133-148.

- Boonkerd, N., D.F. Weber and D.F. Bezwick. 1975. Influence of Rhizobium japonicum strains and inoculation methods on soybeans grown in rhizobia-populated soil. Agron. J. 70 : 547-549.
- Boonkerd, N., P. Wadisirisak, W. Thananusorn, and T. Arayangul. 1986. Competition for nodule site between inoculated and indigenous Bradyrhizobium japonicum strains. In W. Wallace, and S.E. Smith (eds.), The Eight Australian Nitrogen Fixation Conference. The Australian Institute of Agricultural Science, Adelaide. pp 173-174.
- Caldwell, B.E. 1966. Inheritance of a strain specific ineffective nodulation in soybeans. Crop Sci. 6 : 427-428.
- Caldwell, B.E., K. Hinson and H.W. Johnson. 1966. A strain-specific ineffective nodulation reaction in the soybean, Glycine max L. Merrell. Crop Sci. 6 : 495-496.
- Caldwell, B.E. and G. Vest. 1968. Nodulation interactions between soybean genotype and serogroups of Rhizobium japonicum. Crop Sci. 8 : 680-682.
- Caldwell, B.E. 1969. Initial competition of root-nodule bacteria on soybeans in a field environment. Agron. J. 61:813-815.
- Caldwell, B.E. and E.E. Hartwig. 1970. Serological distribution of soybean root nodule bacteria. in soil of southeastern USA. Agron. J. 62 :621-622.

- Clark, F.E. 1957. Nodulation responses of two near-isogenic lines of the soybean. Can. J. Microbiol. 3 : 113-123.
- Darmirgi, S.M., L.R. Frederick and I.C. Anderson. 1967. Serogroup of Rhizobium japonicum in soybean nodules as affected by soil types. Agron. J. 59 : 10-12.
- Date, R.A. 1986. Problems in application of BNF in field crops. In Z.H. Shamsudden *et al* (eds). Proceedings of a UNESCO Regional Symposium and Workshop on Biotechnology of Nitrogen Fixation in the Tropics. Universiti Pertanian Malaysia, Serdang, Selangor. pp. 283-300.
- Devine, T.E. and D.F. Weber. 1977. Genetic specificity of nodulation. Euphytica. 26 : 527-535.
- Devine, T.E. and B.H. Breithaupt. 1980. Phenotypic thermal stability of rhizobiotoxine-induced chlorosis and nodulation controlling gene, *rj*. Crop Sci. 20 : 394-396.
- Devine, T.E. 1985. Host range and compatibility of soybean with rhizobial microsymbionts. In R. Shibles (ed.). World Soybean Research Conference III. West View Press, Colorado. pp 484-492.
- Devine, T.E. and B.H. Breithaupt. 1981. Frequencies of Nodulation Response Alleles, *Rj2*, and *Rj4*, in Soybean Plant Introduction and Breeding Lines. Technical Bulletin NO. 1628. United State Department of Agriculture. pp 1-42.

- Duteau, N.M. 1985. Potential of the Fast-Growing Rhizobium fredii for Use in Soybean Inoculum : Physical and Genetic Characterization of the Nitrogen - Fixing Symbiosis Ph.D. Thesis, Iowa State Univ, Ames, Iowa.
- Eaglesham, A.R.J. and S. Hassouna. 1982. Foliar chlorosis in legumes induced by cowpea rhizobia. Plant and Soil. 65 : 425-428.
- Eaglesham, A.R.J. 1985. Comparison of nodulation promiscuity of US-and Asian-type soya beans. Trop. Agri. 60 : 105-109.
- Ellis, W.R. Ham G.E. and Schmidt E.L. 1984. Persistance and Recovery of Rhizobium japonicum inoculum in a field soil. Agron. J. 76 : 573-576.
- Erdman, L.W. 1944. New developments in legumes inoculation. Soil Sci. Soc. Am. Proc. 8 : 213-214.
- Erdman, L.W., H.W. Johnson and F. Clark. 1957. Varietal responses of soybeans to a bacterial-induced chlorosis. Agron. J. 49 : 267-271.
- Graham, P.H. 1981. Plant factors affecting symbiotic nitrogen fixation in legumes. In P.H. Graham, and S.C. Harris (eds). Biological Nitrogen fixation Technology for tropical Agriculture. Centro Internacional de Agricultura Tropical. Colombia. pp 22-37.
- Hardy, R.W.F., R.C. Burns and R.D. Holsten. 1973. Applications of the acetylene-ethylene assay for measurement of nitrogen fixation. Soil Biol. Biochem. 5 : 47-81.

- Harold, H.K., B.B. Bohlool, T.S. Hu and D.F. Weber. 1982. Fast-growing rhizobia isolated from root nodules of soybean. J. Sci. 51 : 1631-1632.
- Johnson, H.W. and F.E. Clark. 1958. Role of the root nodule in bacterial induced chlorosis of soybeans. Soil Sci. Soc. Am. Proc. 22 : 527-528.
- Johnson, H.W. and U.M. Means. 1962. Serological groups of Rhizobium japonicum recovered from nodules of soybeans (Glycine max) in field soils. Agron. J. 55 : 269-271.
- Jordan, P.C. 1982. Transfer of Rhizobium japonicum Buchanan 1980 to Bradyrhizobium gen. nov., a genus of slow-growing, root nodule bacteria from leguminous plants. Int. J. Syst Bacteriol. 32 : 136-139.
- Kapusta, G. and D.L Rouwenhorst. 1973. Influence of inoculum size on Rhizobium japonicum serogroup distribution frequency in soybean nodules. Agron. J. 69 : 916-919.
- Leonard, L.T. 1943. A simple assembly for use in the testing of cultures of rhizobia. J. Bacteriol. 45 : 523-527.
- Moaward, H. S.M.S Badrel-din and M.A Khalafallah. 1988. Persistance and competitiveness of three Bradyrhizobium japonicum inoculant strains in clay loam Nile Valley soil. Plant and Soil. 108 : 137-141.

- Na Lampang, A. 1976. Interrelationship between soybean varieties and indigenous Rhizobium strains in northeast Thailand. In R.M. Goodman (ed), Expanding the Use of Soybeans. INTSOY Series No. 10, University of Illinois, Urbans Champaign. pp 198-199.
- Nangju, D. 1980. Soybean response to indigenous Rhizobia as influenced by cultivar origin. Agron. J. 72: 403-406.
- Obaton, M. 1977. Effectiveness, saprophytic and competitive ability : three properties of Rhizobium essential for increasing the yield of inoculated legumes. In A. Ayanaba, and P.J. Dart (eds). Biological Nitrogen Fixation in Farming Systems of the tropics. A Wiley-Interscience Publication, Chichester. pp 127-134.
- Owens, L.D., J.F.T. Thompson, R.G. Pitcher and T. Williams. 1972. Structure of rhizobiotoxine, an antimetabolic enol-ether amino acid from Rhizobium japonicum. J. Chem. Soc. Chem. Commun. p. 714.
- Pulver, E.L., F. Brockman and H.C. Wien. 1982. Nodulation of soybean cultivars with Rhizobium spp. and their response to inoculation with R. japonicum. Crop Sci. 22 : 1065-1070.
- Scholla, M.H. and Elkan, G.H. 1984. Rhizobium fredii sp. nov., a fast-growing species that effectively nodulates soybean. Int. J. Syst. Bacteriol. 34 : 484-486.

- Sloger, C. 1969. Symbiotic effectiveness and N₂ fixation in nodulated soybean. Plant Physiol. 44 : 1666-1668.
- Somasegaran, P., H. Hoben and J. Halliday. 1982. The Niftal Manual for Method in Legume-Rhizobium Technology. College of Tropical Agriculture and Human Resources, University of Hawaii, Honolulu. 303 p.
- Williams, L.F. and D.L. Lynch. 1954. Inheritance of a nonnodulation character in the soybean. Agron. J. 46 : 28-29.
- Vest, G. 1970. R_{j3-a} gene conditioning ineffective nodulation in soybean. Crop Sci. 10 : 34-35.
- Vest, G. and B.E. Caldwell. 1972. R_{j4-a} gene conditioning ineffective nodulation in soybean. Crop Sci. 12 : 692-693.