

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
TABLE OF CONTENTS	x
LIST OF TABLES	xiv
LIST OF FIGURES	xvii
LIST OF APPENDIX TABLE	xxi
INTRODUCTION	1
LITERATURE REVIEW	4
Adaptive Response of Plant to Water Stress	4
Morphological Responses	4
Physiological Response	5
Growth Analysis as Related to Water Stress	6
Effect of Water Stress on Yield and Yield Component of Maize	7
Nitrogen Partitioning in Maize	8
CERES-Maize Model	9
EXPERIMENT 1 : Screening of Maize Genotypes for Drought Tolerance	10
Objective	10
Materials and Methods	10
Results and Discussion	16
Phenotypic correlation	16
Screening for drought tolerance	16

	Page
EXPERIMENT 2 : Responses of Morphological and Physiological to Drought Stress on Maize Genotypes under Different Soil Moisture Regimes	21
Objectives	21
Materials and Methods	21
Results and Discussion	26
Growth and development	26
Total dry weight	27
Growth rate	28
Dry matter partitioning	33
Partitioning coefficient	36
Relationship between LAI, LAD, SLW and yield	37
Relationship between LAI, LAD, SLW and irrigation water application	39
Root growth	41
Root distribution	43
Root length density and irrigation water application	45
Relationship between root length density and soil water extraction	47
Canopy temperature	49
Leaf senescence	49
Relationship between canopy temperature and leaf senescence	52
Leaf water potential	52
Relationship between canopy temperature and leaf water potential	52
Stomatal resistance	56

	Page
Water use and water use efficiency	58
Yield response to irrigation plus rainfall	58
Effect of water deficit on kernel yield	61
Yield components response to applied irrigation water	63
EXPERIMENT 3 : Nitrogen Partitioning in Maize Genotypes under Different Nitrogen Fertilization Levels and Soil Moisture Regimes	67
Objectives	67
Materials and Methods	67
Results and Discussion	70
Growth and development	70
Dry matter and nitrogen content	71
Dry matter and nitrogen partitioning	77
Nitrogen use efficiency	83
Effect of water and nitrogen deficits on yield of maize genotypes	85
Effect of water and nitrogen stresses on yield components of maize genotypes	87
EXPERIMENT 4 : Estimation of Genetic Coefficients of Some Thai Maize Varieties of CERES-Maize Model	91
Objectives	91
Materials and Methods	91
Results and Discussion	102

I. Genetic Coefficients Estimation	102
Observed phenological events	102
Estimation of genetic coefficients	104
Model testing	104
II. Model validation	113
Phenological Simulation	113
Growth and Yield Simulation	116

7 SUMMARY AND CONCLUSIONS	123
REFERENCES	127
APPENDIX	142
CURRICULUM VITAE	160

LIST OF TABLES

Table		Page
1	Thirty-one maize genotypes used in Experimental 1.	11
2	Phenotypic correlation coefficients among traits for wet regime (above diagonal, N = 31) and dry regime (below diagonal, N = 31) of thirty-one maize genotypes in 1998.	18
3	Average kernel yield (kg/ha) of thirty-one maize genotypes under different water regimes in 1998.	19
4	Linear regression parameters for responses of total dry matter to mm of irrigation plus rainfall received by three maize genotypes in 1998 and 1999.	27
5	Crop growth rate (CGR), Stalk growth rate (SGR), Leaf growth rate (LGR) and Kernel growth rate (KGR) of three maize genotypes in wet and dry regimes in 1998 and 1999.	32
6	Partitioning coefficient (%) of three maize genotypes in wet and dry regimes in 1998 and 1999.	36
7	Regression equation showing the relationship between growth attributes (X) and kernel yield (Y) of three maize genotypes in 1998 and 1999.	38
8	Linear regression parameters for responses of growth attributes to mm of irrigation plus rainfall received by three maize genotypes in 1998 and 1999.	40
9	Stomata number at lower and upper leaf surface of three maize genotypes under different moisture regimes in 1998.	53

Table	Page
10 Water use efficiency of 3 maize genotypes in wet and dry regimes in 1998 and 1999.	59
11 Effect of moisture regimes on kernel yield (kg/ha) of three maize genotypes for Experiment 2 in 1998 and 1999.	62
12 Linear regression parameters for responses of yield components to mm of irrigation plus rainfall received by three maize genotypes in 1998 and 1999.	64
13 Effect of moisture regimes on yield components of three maize genotypes for Experiment 2 in 1998 and 1999.	65
14 Relationship between kernel yield and yield components of three maize genotypes under different moisture regimes in 1998 and 1999.	66
15 Nitrogen use efficiency of maize genotypes under different nitrogen levels for Experiment 3 in 1998 and 1999.	84
16 Effect of moisture regimes on kernel yield (kg/ha) of three maize genotypes under different nitrogen levels for Experiment 3 in 1998 and 1999.	86
17 Effect of moisture regimes on yield components of three maize genotypes under different nitrogen levels for Experiment 3 in 1998.	89
18 Effect of moisture regimes on yield components of three maize genotypes under different nitrogen levels for Experiment 3 in 1999.	90
19 Phenology and growth coefficients for maize used in the DSSAT version 3.0.	95
20 Genetic coefficients of SW 1 obtained from DSSAT version 3.0.	96
21 Planting, silking and maturity dates from the experiments of NS 1, NSX 9210 and SW 3601 at Phitsanulok Field Crops Experiment Station in 1998-1999.	99

Table		Page
22	Observed phenological events of maize planted at Phitsanulok Field Crops Experiment Station in 1997-1998.	103
23	Genetic coefficients of NS 1, NSX 9210 and SW 3601 obtained after estimation by running GenCalc from DSSAT version 3.0.	104
24	Simulated and observed silking and maturity dates for 3 maize varieties.	106
25	Simulated and observed kernel yield and yield components of three maize varieties at five different planting dates.	112
26	Analysis of variance for observed kernel yield.	113
27	Simulated and observed silking and maturity dates (days) for 3 maize genotypes.	115
28	Simulated and observed kernel yield and yield components of three maize genotypes at three different planting dates.	121

LIST OF FIGURES

Figure		Page
1	Relationship of water applied and distance from line source for thirty-one maize genotypes in 1998.	14
2	Weekly values of rainfall, pan evaporation, solar radiation and maximum and minimum temperature during the crop growth period in 1997/98 and 1998/99.	15
3	Relationship between kernel yield potential (intercept, a) and drought susceptibility (slope, b) of thirty-one maize genotypes in 1998.	20
4	Relationship of water applied and distance from line source for maize experiment 2 in 1998 and 1999.	25
5	Dry matter accumulation in different plant parts of maize genotypes at wet and dry regimes in 1998.	30
6	Dry matter accumulation in different plant parts of maize genotypes at wet and dry regimes in 1999.	31
7	Percentage of total dry weight at various stages of growth and distribution of weight among plant parts of three maize genotypes at wet and dry regimes in 1998.	34
8	Percentage of total dry weight at various stages of growth and distribution of weight among plant parts of three maize genotypes at wet and dry regimes in 1999.	35
9	Average root length density at different growth stages of maize genotypes grown under wet (a), moderate (b) and dry (c) regimes in 1998.	42

Figure		Page
10	Average root length density at R ₄ various positions and different soil depths of 3 maize genotypes under wet (a) and dry (b) regimes in 1998.	44
11	Average root length density at various growth stages and different soil depths of 3 maize genotypes under wet (a) and dry (b) regimes in 1998.	46
12	The water extraction pattern of NS 1 (a), NSX 9210 (b) and SW 3601 (c) in dry regime of different depth of soil profile in 1998.	48
13	Changes in canopy temperature of three maize genotypes in wet and dry regimes in 1998 and 1999.	50
14	Percentage of leaf senescence in three maize genotypes at six growth stages in 1998 and 1999.	51
15	Changes in leaf water potential of three maize genotypes in wet and dry regimes in 1998 and 1999.	54
16	Relationship between leaf water potential and canopy temperature of three maize genotypes at five moistures during the six periods (71-81 DAE) in 1998.	55
17	Stomatal resistance in three maize genotypes in wet and dry regimes in 1998.	57
18	Relationship between kernel yield and water applied plus rainfall for three maize genotypes in 1998 (a) and 1999 (b).	60
19	Relationship of water applied and distance from line source for maize experiment 3 in 1998 and 1999.	69
20	Biomass in each plant parts of 3 maize genotypes under low and high nitrogen in wet regime in 1998.	73

Figure		Page
21	Biomass in each plant parts of 3 maize genotypes under low and high nitrogen in dry regime in 1998.	74
22	Total nitrogen content in each plant parts of 3 maize genotypes under low and high nitrogen in wet regime in 1998.	75
23	Total nitrogen content in each plant parts of 3 maize genotypes under low and high nitrogen in dry regime in 1998.	76
24	Percent of total dry weight in different plant parts of three maize genotypes : (a) NS 1 (b) NSX 9210 (c) SW 3601, at low and high nitrogen application under wet regime in 1998.	79
25	Percent of total dry weight in different plant parts of three maize genotypes : (a) NS 1 (b) NSX 9210 (c) SW 3601, at low and high nitrogen application under dry regime in 1998.	80
26	Percent of total nitrogen in different plant parts of three maize genotypes : (a) NS 1 (b) NSX 9210 (c) SW 3601, at low and high nitrogen application under wet regime in 1998.	81
27	Percent of total nitrogen in different plant parts of three maize genotypes : (a) NS 1 (b) NSX 9210 (c) SW 3601, at low and high nitrogen application under dry regime in 1998.	82
28	File components assigned to estimate genetic coefficients of maize in DSSAT v. 3.0.	97
29	Comparison between simulated and observed leaf, stalk and kernel dry weights of NS 1 at five planting dates; (a) November, (b) January, (c) March, (d) May and (e) July.	109
30	Comparison between simulated and observed leaf, stalk and kernel dry weights of NSX 9210 at five planting dates; (a) November, (b) January, (c) March, (d) May and (e) July.	110

Figure		Page
31	Comparison between simulated and observed leaf, stalk and kernel dry weights of SW 3601 at five planting dates; (a) November, (b) January, (c) March, (d) May and (e) July.	111
32	Comparison between simulated and observed days to silking and maturity of three maize genotypes in 1998-1999.	114
33	Comparison between simulated and observed leaf, stalk and kernel dry weights of NS 1 at three planting dates; (a) November, (b) January and (c) February.	118
34	Comparison between simulated and observed leaf, stalk and kernel dry weights of NSX 9210 at three planting dates; (a) November, (b) January and (c) February.	119
35	Comparison between simulated and observed leaf, stalk and kernel dry weights of SW 3601 at three planting dates; (a) November, (b) January and (c) February.	120
36	Comparison between simulated and kernel yield of NS 1, NSX 9210 and SW 3601 at three planting dates.	122

LIST OF APPENDIX TABLES

Appendix Table		Page
1	Chemical and physical soil properties at the experiment site.	142
2	Schedule of irrigation treatment applied during the 1998 and 1999 growing season.	143
3	Simple linear regression (N = 12) between grain yield and reduction in water applied of thirty-one maize genotypes in 1998.	144
4	Soil properties for Phen series from ThaiSIS.	145
5	FILEA for estimation of genetic coefficients of three maize varieties.	146
6	FILEX for estimation of genetic coefficients of NS 1.	147
7	FILEX for estimation of genetic coefficients of NSX 6210.	150
8	FILEX for estimation of genetic coefficients of SW 3601.	153
9	Genetic coefficients and standard deviation for three maize varieties by using GenCalc.	156
10	FILEA for validation of CERES-Maize model.	157
11	FILEX for validation of CERES-Maize model.	158