

## References

- Abawi, G.S. and M.A. Pastor-Corrales. 1990. Seed transmission and effect of fungicide seed treatments against *Macrophomina phaseolina* in dry edible beans. *Turrialba* 40(3):334-339.
- Agarwal, V.K. and J.B. Sinclair. 1987. Principles of Seed Pathology. CRC press, Inc. Boca Raton, Florida. Vol. I &II.
- Agarwal, V.K.; S.B. Mathur, and P. Neergaard. 1972. Some aspects of seed health testing with respect to seed-borne fungi of rice, wheat, blackgram, greengram and soybean grown in India. *Indian Phytopathology* XXV:91-100.
- Alagarsamy, G. and Sivaprakasam, K. 1988. Effect of antagonists in combination with carbendazim against *Macrophomina phaseolina* infection in cowpea. *J. Biol. Control.* 2(2): 123-125.
- Baker, K. F. 1972. Thermotherapy of planting material. *Phytopathology* 52:1244-1255.
- Basher, M. A. and R. Bharat 1994. Antagonistic potential of root region microflora of chickpea against *Fusarium oxysporum* f. sp. *ciceri*. *Bangladesh J. Botany* 23(1):13-19.
- Basir, M.; M. Zubair and B. A. Malik 1987. Disease resistance sources and utilization in breeding improved mungbean in Pakistan p623-630. In Mungbean - Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.
- Begum, M. M.; I. Hossain and M. S. Haque. 1998. Biocontrol of seed borne *Fusarium oxysporum* with *Trichoderma harzинium*. *Bangladesh J. Environ. Sci.* 4:128-133.

- Bilgrami, K.S.; R.K. Sinha, and T. Prasad. 1978. Effect of fungal flora on seed contents of moong. Indian Phytopathology 31:476-479.
- Bilgrami, K.S.; T. Prasad,; Jamaluddin, and A.K.Roy. 1976. Studies on the deterioration of some pulses by fungi. Indian Phytopathology 29: 274-277.
- Burman, U. and S. Lodha. 1996. *Macrophomina phaseolina* induced changes in plant water relations of resistant and susceptible cowpea genotypes. Indian Phytopathology 49(3):254-259.
- Chainuvati, C.; N. Potan, and T. Worasan. 1987. Mungbean and black gram production and development in Thailand p657-666. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service, November 16-20, 1987.
- Chaudhary, K. C. B. and A. K. Pal. 1982. Infection of sannhemp (*Crotalaria juncea*) seeds by *Macrophomina phaseolina*. Seed Sci. & Technol. 10:151-153.
- Chen, C.Y.; S.C.S. Tsou, and H.H.Wang. 1987. Utilization patterns of mungbean in the Chienese diet. p498-507. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.
- Chinsawangwatanakul, A. and P. Surin. 1980. Mungbean diseases in Thailand. In Diseases and insect pests of mungbean and blackgram- a bibliography, AVRDC. February 1988. Series no-5.
- Chohan, J.S. and J. Kaue. 1975. Seed mycoflora of sunflower and control of seed-borne pathogens. Indian J. of Mycology and Plant Pathology 5:210-211.

- Chou, J. and W. Wu., 1995. Seed-borne fungal pathogens of ornamental flowering plants. *Seed Sci. & Technol.* 23:201-209.
- Clavet, C.; J. Para, and J. M. Barea. 1990. Interactions of *Trichoderma* spp. with *Glomus mosseae* and two wilt pathogenic fungi. *Agriculture, Ecosystems and Environment* 29(1-4):59-65.
- Davet, P. 1987. Criteria of selecting *Trichoderma* clones antagonistic to sclerotial fungi in soil. *Bulletin OEPP*. 17(4): 535-540.
- De, R.K.; R.K. Chaudhary, and Naimuddin. 1996. Comparative efficacy of bio-control agents and fungicides for controlling chickpea wilt caused by *Fusarium oxysporum* f. sp *ciceri*. *Indian J. of Agril. Sci.* 66(6):370-373.
- Delouche, J. C. and C. C. Baskin. 1973. Accelerated aging technique for predicting the relative storability of seed lots. *Seed Sci. & Technol.* 1:427-452. In Gupta, I.J.; A. F. Schmitthenner and M. B. McDonald. 1993. Effect of storage fungi on seed vigour of soybean. *Seed Sci. & Technol.* 21:581-591.
- Deng, T.C. 1984. The major diseases of mungbean and their control. *Pesticide World*. 12:47-49 In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Deshmukh, P. P. and J. G. Raut. 1992. Antagonism by *Trichoderma* spp. on five plant pathogenic fungi. *New Agriculturist*. 3(2):127-130.
- Dhar, V. and A.K. Sarbhoy. 1989. Studies on the germination and longevity of pycnidiospores of *Macrophomina phaseolina*. *Indian Phytopathology* 42 (1) :123-127.
- Dhedhi, B. M.; O. M. Gupta, and V. A. Patel. 1990. Antagonistic effect of micro-organisms to *Fusarium oxysporum* f. sp. *ciceri*. *Indian J. Mycology and Plant Pathology* 20(1):70-71.

- Ehteshamul-Haque, S. and A. Ghaffar. 1995. Role of *Bradyrhizobium japonicum* and *Trichoderma* spp. in the control of root rot disease of soybean. *Acta Mycologica* 30(1):35-40.
- Ehteshamul-Haque, S.; R. Y. Hashmi and A. Ghaffar. 1992. Biological control of root rot disease of lentil. *Lens Newsletter* 19(2): 43-45.
- Elad, Y.; R. Barak.; I. Chet, and Y. Henis. 1983. Ultrastructural studies of the interaction between *Trichoderma* spp. and plant pathogenic fungi. *Phytopathologische Zeitschrift*. 107(2):168-175.
- Ellis, M. A.; G. E. Galvez, and J. B. Sinclair. 1976. Effect of pod contact with soil on fungal infection of dry bean seeds. *Plant Disease Reporter* 60 (11):974-976.
- Ellis, M.A.; E.H. Paschal; P.E. Powell and F.D. Tenne. 1979. Internally seedborne fungi of soya bean in Puerto Rico and their effect on seed germination and field emergence. *Trop. Agric (Trinidad)*. 56(2):171-174.
- Fakir, G.A.; M.H. Rao, and M.J. Thirumalachar. 1976. Seed transmission of *Macrophomina phaseolina* in sunflower. *Plant Disease Reporter* 60(9): 736-737.
- Farzana, A.; Ghaffar, A. and Ali, F. (1991). Effect of seed treatment with biological antagonists on rhizosphere mycoflora and root infecting fungi of soybean. *Pakistan J. Botany* 23(2):183-188.
- Fernandez, G.C.J. and S. Shanmugasundaram. 1987. The AVRDC mungbean improvement program: the past, present and future p58-70. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.

- Fernandez, M. R. 1992. The effect of *Trichoderma harzianum* on fungal pathogens infesting soybean residues. *Soil Biology and Biochemistry* 24 (10):1027-1029.
- Ferrata, M. and V. D'Ambra. 1985. Morphological aspects of the parasitism of *Trichoderma harzianum* Rifai on hyphae and sclerotia of *Sclerotium rolfsii* Sacc. *Rivista di Pathologia Vegetale* 21(2):53-60.
- Fuhlbohm, M. J.; E. A. B. Aitken, and M. J. Ryley. 1997. Seed infection of mungbean by *Macrophomina phaseolina*. *Australian Phytopathological Society Conference '97*. 39p.
- Gangopadhyay, S.; T.D. Wyllie, and V.D. Luedders. 1970. Charcoal rot disease of soybean transmitted by seeds. *Plant Disease Reporter* 54 (12):1088-1091.
- Ghosh, J. and B. Nandi. 1986. Deteriorative abilities of some common storage fungi of wheat. *Seed Sci. & Technol.* 14:141-149.
- Gowda, C.L.L. and A.K Kaul. 1982. Pulses in Bangladesh, BARI & FAO of the United Nations 138-166.
- Grewal, J.S. and M. Pal. 1971. Report of the work done on disease of arhar and dry root rot of mung during 1970. Hissar: Haryana Agril. Univ. 109-111. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Grewal, J.S. 1988. Diseases of pulse crops - an overview. *Indian Phytopathology* 41(1): 1-14.
- Grezes-Besset, B.; N. Lucante,; V. Kelechian,; R. Dargent, and H. Muller. 1996. Evaluation of castor bean resistance to sclerotial wilt disease caused by *Macrophomina phaseolina*. *Plant Disease* 80: 842-846.

- Grondeau, C. and R. Samson. 1994. A review of thermotherapy to free plant materials from pathogens, especially seeds from bacteria. Critical Reviews in Plant Sciences 13(1):57-75.
- Gupta, I. J. and H. S. Cheema. 1990. Effect of microsclerotia of *Macrophomina phaseolina* and seed dressers on germination and vigour of sesame seed. Seed Research 18(2):169-172.
- Haran, S.; H. Schickler,; A. Oppenheim, and I. Chet. 1996. Differential expression of *Trichoderma harzianum* chitinase during mycoparasitism. Phytopathology 86(9):980-985.
- Harman, G.E.; Chet, I. and Baker, R. 1980. *Trichoderma hamatum* effects on seed and seedling disease induced in radish and pea by *Pythium* spp. or *Rhizoctonia solani*. Phytopathology 70:1167-1172.
- Himani, B. 1996. Influence of environmental conditions on antagonistic activity of *Trichoderma* spp. against *Fusarium udum*. Indian J. Mycology and Plant Pathology 26(1):58-63.
- Holliday, P. and E. Punithalingam. 1970. *Macrophomina phaseolina*. Descriptions of pathogenic fungi and bacteria. No. 275. Commonwealth Mycological Institute, Kew, Surrey, England.
- Holliday, P. 1980. Fungal diseases of tropical crops. Cambridge University Press. 254-255.
- Hooda, I. And R.K. Grover.1988. Effect of age, quantity of inoculum and isolates of *Macrophomina phaseolina* on the pathogenesis of mungbean and its control by chemicals. Indian Phytopathology 41 (1): 107-117.
- Hussain, S.; Ghaffar, A. and Aslam, M. 1990. Biological control of *Macrophomina phaseolina* charcoal rot of sunflower and mungbean. J. Phytopathology 130: 157-160.

- Iqbal, S. M.; A. Bakhsh,; S. Hussain, and A. B. Malik. 1995. Microbial antagonism against *Sclerotium rolfsii* Sacc. The cause of collar rot of lentil. *Lens Newsletter* 22(1-2): 48-49.
- ISTA. 1976. International Rules for Seed Testing. International Seed Testing Association. *Seed Sci. & Technol.* 4: 3-177.
- ISTA. 1993. International Rules for Seed Testing. *Seed Sci. & Technol.* 21. Supplement. 1-288.
- Jacobs, D. and O. Kamoen. 1987. Roll of cell wall lysing enzymes of *Trichoderma harzинium*. *Med. Fac. Landbouw. Rijks. Gent.* 56(26):751-758.
- Jadeja, R. G. and V. A. Patel. 1989. Biochemical changes due to infection of *Macrophomina phaseolina* in lima bean plants. *Indian J. Mycology and Plant Pathology* 19(1):135-136.
- Jain, N.K. and M.N. Khare. 1972. Chemical control of *Rhizoctonia bataticola* causing diseases of urid. 6(4):461-465. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Kabeere, F.; J. G. Hampton, and M. J. Hill. 1997. Transmission of *Fusarium graminearum* (Schwabe) from maize seeds to seedlings. *Seed Sci. & Technol.* 25:245-252.
- Khan, A.L.; K.B. Alam, and M.J. Talukder. 1977. Seed-borne fungi of pulses in Bangladesh. *Bangladesh J. Agriculture* 2(1): 163-166.
- Lifshitz, R.; M. T. Windham and R. Baker. 1986. Mechanism of biological control of pre-emergence damping-off of pea by seed treatment with *Trichoderma* spp. *Phytopathology* 76(7):720-725.

- Lim, T.K. and B. K. Teh. 1990. Antagonistic *in vitro* of *Trichoderma* spp. against several basidiomycetous soil-borne pathogens and *Sclerotium rolfsii*. Z. Pflkrank. Und Pflschutz. 97(1):33-41.
- Maden, S.; D. Singh,; S. B. Mathur. and P. Neergaard. 1975. Detection and location of seed-borne inoculum of *Ascochyta rabiei* and its transmission in chick pea. Seed Science and Technology 3:667-681.
- Mahanty, N.N. and V.B. Rao. 1968. Fungal diseases of pulses. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Maheshwari, R. K.; S. K. Mathur and A. Mathur. 1984. Effect of seed-borne fungi on seed viability and seedling vigour. Indian Phytopathology. 37:713-714.
- Maite, D. and C. Sen. 1985. Integrated biocontrol of *Sclerotium rolfsii* with nitrogenous fertilizers and *Trichoderma harzianum*. Indian J. Agril. Sci. 55(7):464-468.
- Maude, R.B. 1983. Eradicative seed treatments. Seed Sci. and Technol. 11:907-920.
- Maude, R.B. 1996. Seedborne diseases and their control. Principal and practice. CAB International. Walligford Oxon OX10 8DE UK.
- McGee, D. C. 1988. Maize Diseases. A reference for seed technologists. APS Press. The American Phytopathological Society, St Paul, Minnesota.
- Mihail, J. D. and S. M. Alcorn. 1982. Quantative recovery of *Macrophomina phaseolina* sclerotia from soil. Plant Disease 66: 662-663.
- Miller, J.C.J. and G.C.J. Fernandez. 1987. Selecting and breeding for enhanced N<sub>2</sub> fixation in mungbean p111-123. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.

- Morris, D.L. 1948. Quantitive Determination of carbohydrate with Dreywood's Anthrone reagent. *Science* 107:254-255. In Yoshida, S.; D. A. Forno; J.H. Cock and K.A. Gomez. 1976 Laboratory manual for physiological studies of rice. 3<sup>rd</sup> edition. The International Rice Resesarch Institute. 46-49.
- Mukhopadhyay, A.N. 1989. National seminar and VII workshop of AICRP on biological control Lucknow, October 23-25, 1989.
- Mukhopadhyay, A.N. 1994. Biocontrol of soil borne fungal plant pathogens - current status, future prospect and potential limitations. *Indian Phytopathology* 47(2):119-126.
- Mukhopadhyay, D. and B. Nandi. 1976. Carbohydate content of jute plants infected with *Macrophomina phaseolina*. *Indian Phytopathology* 29:208-209.
- Mutto, S.; V. D'Ambra and M. Ferrata. 1986. Ultrastructural aspects of the parasitism of *Trichoderma harzianum* on sclerotia of *Sclerotium rolfsii*. *Phytopathologia Mediterranea* 25(1-3):10-18.
- Nath, R.; S.B. Mathur, and P. Neergaard.1970. Seed-borne fungi of mungbean (*Phaseolus aureus* Roxb.) from India and their significance. *Proc. Seed Test. Ass.* 35:225-241.
- Nayak, B. K. and N. Behera. 1994. Effect of *Macrophomina phaseolina* on seed germination and seedling rot in *Phaseolus mungo*. *J. Mycopathol. Res.* 32(1): 71-74.
- Neergaard, P. 1977. Seed Pathology Vol. I. Storage fungi p282-297. In *Seed Pathology*, Vol II The Macmillan Press, London and Barings.
- Neergaard, P. 1979. *Seed Pathology Vol. II*. The Macmillan Press, London and Barings.

- Nik, W.Z. 1983. Seed borne fungi of soybean (*Glycine max* (L.) Merrill) and mungbean (*Vigna radiata* (L.) Wilczek) and their pathogenic potential. Malaysian Applied Biology 12(1):21-28.
- Nine, Y.L. 1986. Opportunities for research on diseases of pulse crops. Indian Phytopathology 39(3):333-342.
- Ossoble, A.A. 1987. Status of mungbean in Somalia p637-640. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.
- Pal, M. 1996. Pulses disease scenario. Indian Phytopathology 49(2): 129-131.
- Pannu, R.K., and D. P. Singh. 1987. Influence of water deficits on morphophysiological and yield behavior of mungbean (*Vigna radiata* (L.) Wilczek) p252-259. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.
- Papavizas, G. C. and N. G. Klag. 1975. Isolation and quantitative determination of *Macrophomina phaseolina* from soil. Phytopathology. 65:182-187.
- Pichitporn, S. and C. Thavarasook. 1990. Blackgram cultivars p17-22. In Proceedings of the mungbean meeting '90, Chiang Mai, Thailand, February 23-24, 1990.
- Pratt, R. G.; M. R. McLaughlin,; G. A. Pederson, and D. E. Rowe. 1998. Pathogenicity of *Macrophomina phaseolina* to mature plant tissues of alfalfa and white clover. Plant Disease 82:1033-1038.
- Prabhavat, S. 1990. Mungbean utilization in Thailand p9-16. In Proceedings of the mungbean meeting '90, Chiang Mai, Thailand. February 23-24, 1990. 9-15.

- Purushotham, S.P.; K. L. Patkar; H.S. Prakash and H.S. Shetty. 1996. Storage fungi and their influence on rice seed quality. Indian Phytopathology 49(2) : 152-156.
- Putasamai, K. and P. Surin. 1988. Charcoal rot of blackgram p242-257. In 3<sup>rd</sup> report of soybean research. 22-23 Nov. 1988. (in Thai).
- Quyen, M.V. 1987. Mungbean for cropping systems in Vietnam p669-678. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.
- Rath, G.C. and G.N. Routray. 1978. Internally seed transmitted diseases of moong and urid. Science and Culture. 44(1):40-41. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Raut, J.G. 1983. Transmission of seed-borne *Macrophomina phaseolina* in sunflower. Seed Sci. & Technol. 11:807-814.
- Raut, J.G. and B.B. Bhombe. 1984. Longevity of *Macrophomina phaseolina* in sunflower seeds. Indian Phytopathology 37:333-334.
- Reddy, M.R.S. and J. Subbayya. 1981. *Macrophomina phaseolina* on seed health of blackgram (*Phaseolus mungo* L.). Current Research. 10 (4):58. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Reuveni, R.; A. Nachmias and J. Krikun. 1983. The role of seed-borne inoculum on the development of *Macrophomina phaseolina* on melon. Plant Disease 67(3):280-281.
- Richardson, M. J. 1979. Annotated list of seed-borne diseases. 3<sup>rd</sup> edition, Proc. Int. Seed Test. Assoc. 23:1-320.

- Sadashivaiah, A.S.; K.G. Ranganathaiah, and D.N. Gowda. 1986. Seed health testing of *Helianthus annus* with special reference to *Macrophomina phaseolina*. Indian Phytopathology 39:445-447.
- Sandhu, A.; R. D. Singh and U. Kant. 1998. Changes in carbohydrate contents and hydrolyzing enzymes in cowpea infected by *Macrophomina phaseolina* (Tassi) Goid. Annals of Arid Zone 37 (4):423-425.
- Saxena, N. and D. Karan. 1991. Effect of seed-borne fungi on protein and carbohydrate contents of sesame and sunflower seeds. Indian Phytopathology 44:134-136.
- Saxena, R.M. and S. Sinha. 1979. Field mycoflora of *Vigna radiata* (L.) Wilczek and *Vigna mungo* (L.) Hepper in relation to pre-emergence and post-emergence mortalities. Seed Research 7(2):159-164.
- Scholefield, S. and M. J. Griffin. 1979. New or unusual records of plant diseases and pests. Plant Pathology 28:155-156.
- Sharada, R. and H. S. Shetty. 1987. Location and transmission of *Macrophomina phaseolina* in blackgram seeds. Indian Phytopathology 40(2): 194-196.
- Sharma, B. K. and B. M. Singh. 1990. Biological control of white rot of pea caused by *Sclerotium sclerotiorum* (Lib.) de Bary. J. Biological control 4 (2): 132-134.
- Sharma, S. R. 1988. Evaluation of different fungicides against ashy stem blight (*Macrophomina phaseolina*) of french bean. Indian J. Plant Protection 16:285-287.

- Shukla, D.N. and S.N. Bhargava. 1976. Some pathogenic fungi from pulses and oil seed crops. National academy of sciences, India, proceedings (Sec. B: biological science). 46(B): 531-532. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Singh, A. and T.P. Bhowmik. 1991. Prevalence and severity of root rot of sesamum caused by *Macrophomina phaseolina*. Indian Phytopathology 44: 235-238.
- Singh, D. 1991. Biocontrol of *Sclerotinia sclerotiorum* (Lib.) de Bary by *Trichoderma harzianum*. Tropical Pest Management 37(4):374-378.
- Singh, D. V. 1984. Effects of the metabolites of seed-borne fungi on wheat seed germination and seedling vigour. Indian Phytopathology 37: 343-346.
- Singh, I. and J.S. Chohan. 1977. Seed borne fungi in black gram (*Phaseolus mungo*) in Punjab. Indian J. Mycology & Plant Pathology 6:80.
- Singh, K.; S.S. Virmani, and J.S. Chohan. 1974. Field screening of urid (*Phaseolus mungo*) varieties against maturity wilt phase of the dry root rot disease caused by *Macrophomina phaseolina* in the Punjab. Indian J. Mycology & Plant Pathology 3:194-195.
- Singh, R. R. and P. Shukhla. 1987. Amino acid changes in black gram leaves infected with *Colletotrichum truncatum*. Indian Phytopathology 40:241-242.
- Singh, T. and D. Singh. 1982. Transmission of seed-borne inoculum of *Macrophomina phaseolina* from seed to plant. Proc. Indian Acad. Sci. (Plant Sci.). 91(4):357-370.

- Singh, V.P.; A. Chhabra, and R.P.S. Kharb.1987. Production and utilization of mungbean in India p486-497. In Mungbean- Proceedings of the Second International Symposium, Bangkok, Thailand. AVRDC tropical vegetable information service.
- Sinha, M.K. and T. Prasad. 1977. Deterioration of 'Arhar' seeds by *Aspergillus flavus*. Indian Phytopathology 30: 70-72.
- Sinha, O.K. and Khare, M.N. (1977). Control of seed-borne *Macrophomina phaseolina* and *Fusarium equiseti* by hot water treatment of cowpea seeds. Seed Research 5(1):20-22.
- Sinha, O.K. and Khare, M.N. 1977. Site of infection and further development of *Macrophomina phaseolina* and *Fusarium equiseti* in naturally infected cowpea seeds. Seed Sci. & Technol. 5:721-725.
- Sinha, O.K. and M.N. Khare.1977. Chemical control of *Macrophomina phaseolina* and *Fusarium equiseti* associated with cowpea seeds. Indian Phytopathology 30: 337-340.
- Sivan, A. and I. Chet. 1989. The possible role of competition between *Trichoderma harzинium* and *Fusarium oxysporum* on rhizosphere colonization. Phytopathology 79: 198-203.
- Srichuwong, S. 1992. Seed-borne infection of *Colletotrichum truncatum* (Schw.) Andrus and Moore in soybean seeds and its control. - Ph. D. Thesis, Pertanian University Malaysia, Malaysia.
- Sugha, S. K.; B. K. Sharma and P. D. Tyagi. 1993. Factors affecting development of collar rot of gram (*Cicer arietinum*) caused by *Sclerotium rolfsii*. Indian J. Agril. Sci. 63(6):382-385.
- Suhag, L.S. and D. Suryanarayana. 1977. Some aspects of seed health testing with respect to seed borne fungi of pulse crops grown in Haryana. Indian J. Mycology & Plant Pathology 6:32-36.

- Sultana, N.; A. K. Khanzada, and A. Gaffar. 1994. Location of *Macrophomina phaseolina* in seeds of pumpkin and development of charcoal rot disease. *Pakistan J. Botany* 26(1):17-180.
- Sumbali, G. and R. S. Mehrotra. 1982. Post infection chemical changes in round gourds infected with *Geotrichum candidum*. *Indian J. Mycology and Plant Pathology*. 12(1): 48-49.
- Sumitha, R. and S. J. Gaikwad. 1995. Checking Fusarium wilt of pigeon pea by biological means. *J. Soils and Crops* 5(2):163-165.
- Tarr, S.A.J. 1972. *The Principles of Plant Pathology*. Macmillan, London, UK, 632pp.
- Thirumalachar, M. J.; Paul Neergaard, and G. A. Fakir. 1977. Methods for pathogenicity tests of seed-borne *Macrophomina phaseolina* isolated from different hosts. *Phytopath. Z.* 88:234-237.
- Tripathi, H.S.; Singh, R.S. and Chaube, H.S. 1987. Effect of dry heat treatment on the survival of *Ascochyta rabiei*. *International chickpea newsletter*. 16:13
- Upadhyay, J.P. and A. N. Mukhopadhyay. 1986. Biological control of *Sclerotium rolfsii* by *Trichoderma harzianum* in sugarbeet. *Tropical Pest Management* 32(3):215-220.
- Vidhyasekaran, P.; G. Arjunan, and K. Ranganathan. 1976. Field tolerance of some blackgram varieties to root rot disease caused by *Macrophomina phaseolina*. *Madras Agril. J.* 63(3): 176-178. In Diseases and insect pests of mungbean and blackgram - a bibliography, AVRDC. February 1988. Series no-5.
- Vidhyasekaran, P.; N. Ramadoss,; K. Ranganathan and V. Krishnasamy. 1983. Increase in protein content of rice due to helminthosporiose infection. *Indian Phytopathology* XXVI: 736-738.

- Vidhyasekaren, P. and G. Arjunan. 1978. Studies on leaf blight of urd bean (*Vigna mungo*) caused by *Macrophomina phaseolina*. Indian Phytopathology 31:361-362.
- Watanabe, T. 1972. *Macrophomina phaseoli* found in commercial kidney bean seed and in soil, and pathogenicity to kidney bean seedlings. Ann. Phytopath. Soc. Japan 38: 100-105.
- Watanasit, A. and W. Thanomsub. 1995. Mungbean and blackgrams. In A guidebook for field crops production in Thailand. Field Crops Research Institute, Department of Agriculture. 45-55.
- Weerated, C. (1995). Detection of seed-borne fungi in 14 lines of mungbean seeds (*Phaseolus aureus*) by blotter method. In the second National Plant Protection Conference. 2:397-400.
- Yang, C.Y. 1987. Mungbean diseases and their control - a review. 1-35.
- Yoshida, S.; D. A. Forno; J.H. Cock and K.A. Gomez. 1976. Laboratory manual for physiological studies of rice. 3<sup>rd</sup> edition. The International Rice Resesarch Institute. 14-16.