

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
TABLE OF CONTENTS	x
LIST OF TABLES	xiv
LIST OF FIGURES	xvii
LIST OF PLANT SPECIES	xix
ABBREVIATIONS	xxi
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	3
2.1 Boron in plants	3
2.1.1 Boron Uptake and translocation	3
2.1.2 Boron function in plants	4
a) Cell wall	4
b) Membrane	5
c) Phenol metabolism	6
2.1.3 Boron in reproductive growth	8
2.2 Boron deficiency symptoms	8
2.2.1 Visible symptoms	8
2.2.2 Pollen sterility	10

2.2.3 Boron concentration in plant tissues	13
2.2.4 Assessment of grain set failure	13
2.3 B efficiency	15
2.3.1 Definition	15
2.3.2 Genotypic variation	16
2.3.3 Mechanisms	18
CHAPTER 3 Variation in boron responses in wheat (<i>Triticum aestivum</i> L.) genotypes during vegetative and reproductive growth	
3.1 Introduction	26
3.2 Materials and Methods	27
3.2.1 Sand culture experiment	27
3.2.2 Solution culture experiment	29
3.3 Results	32
3.3.1 Sand culture experiment	32
3.3.1.1 The duration of the apical development	32
3.3.1.2 The number of leaves	32
3.3.1.3 The flag leaf B concentration	36
3.3.1.4 The number of spikelet spike ⁻¹	38
3.3.1.5 Grain set	39
3.3.1.6 Relative response of grain set to B	40

3.3.2	Solution culture	41
3.3.2.1	Plant deficiency symptom	41
3.3.2.2	Plant growth	41
3.4	Discussion	42
3.4.1	Classifying wheat genotypes for boron efficiency	42
3.4.2	Fang 60 vs SW 41	45
CHAPTER 4 The effect of boron supply on boron uptake and distribution in wheat genotypes		
4.1	Introduction	47
4.2	Materials and methods	48
4.3	Results	49
4.3.1	Yield and yield component and GSI	49
4.3.2	Boron concentration in plant parts	55
4.3.3	Boron content in plant parts	60
4.3.4	Relative B distribution with in the whole plant	63
4.3.5	Relative B distribution in the upper shoot of the main stem	65
4.4	Discussion	
CHAPTER 5 The effect of B on pollen development in B-efficient (cv. Fang 60) and B-inefficient (cv. SW 41) wheat (<i>Triticum aestivum</i> L.) genotypes.		

5.1 Introduction	71
5.2 Materials and methods	72
5.3 Results and discussion	73
CHAPTER 6 Enhanced boron transport into the ear of wheat as a mechanism for boron efficiency	
6.1 Introduction	78
6.2 Materials and methods	80
6.3 Results and discussion	83
6.3.1 Pollen viability	83
6.3.2 Ear boron concentration	85
6.3.4 Leaf boron concentration	85
6.3.5 Shoot B and pattern of B distribution	86
6.3.6 Long distance transport of boron	88
6.3.7 Conclusions	91
CHAPTER 7 General Discussion	
7.1 Classifying wheat genotypes for boron efficiency	92
7.2 The role of B in anther development	93
7.3 Mechanisms of B efficiency	94
7.4 Suggestion for future research	96
REFERENCES	98
CURRICULUM VITAE	112

LIST OF TABLES

Table		Page
3.1	The effect of B (0 μM B (B0), 10 μM B (B10)) on the time taken (the number of days) for the spike to develop from floral initiation to terminal spikelet stage.	33
3.2	The effect of B (0 μM B (B0), 10 μM B (B10)) on the time taken (the number of days) for the spike to develop from double ridge to terminal spikelet stage.	34
3.3	The effect of B (0 μM B (B0), 10 μM B (B10)) on the number of leaves per plant after 14 (D14), 25 (D25) and 39 (D39) days growth.	35
3.4	The effect of B (0 μM B (B0), 10 μM B (B10)) on the number of leaves per plant after 31 (D31) days growth.	36
3.5	The effect of B (0 μM B (B0), 10 μM B (B10)) on B concentration (mg kg^{-1}) in the flag leaves at mid boot and full boot stages.	37
3.6	The effect of B (0 μM B (B0), 10 μM B (B10)) on the number of spikelet spike ⁻¹ .	38
3.7	The effect of B (0 μM B (B0), 10 μM B (B10)) on grain set.	39
3.8	Relative responses of grain set to low B	41
4.1	The effect of B on grain set index (%) in five wheat genotypes.	51

Table	Page
4.2 Day of full ear emergence, tiller plant ⁻¹ , spikelet plant ⁻¹ , top dry weight and straw yield of wheat genotypes as influenced by B	54
4.3 Tissue B concentration ($\mu\text{g g}^{-1}$) of the flag leaf sheath (a), the Penultimate leaf sheath (b), stem below penultimate leaf node (c) and stem above flag leaf node (d) at full boot as influenced by B.	59
4.4 Boron content ($\mu\text{g pot}^{-1}$) of flag leaf sheath (FLS), penultimate leaf sheath (PLS), spike, stem above flag leaf node (SAFL) and stem segment between flag leaf and penultimate leaf node (SSFP) at full boot as influenced by B.	62
4.5 The effect of B on the relative distribution within the whole plant in wheat genotypes. (SAPL=stem above penultimate leaf node, SBPL=stem below penultimate leaf node)	64
5.1 Reaction to KI/I ₂ staining for starch in the pollen of two wheat cultivars at anthesis after 5 days of B treatment earlier at tetrad-young microspore stages.	77
6.1 Total B concentration in plant parts (mg kg^{-1} dry weight) and relative B distribution (% of total shoot B) after 5 days of varied B supply.	87
6.2 Dry weight of ear on the main stem (g plant^{-1}) at day 0 (D0), day 1 (D1) and day 5 (D5) under different boron supply.	88

Table	Page
6.3 The content of ^{11}B in the ear on the main stem ($\mu\text{g ear}^{-1}$) at day 0 (D0), day 1 (D1) and day 5 (D5) under different boron supply.	89
6.4 The content of ^{10}B in the ear on the main stem ($\mu\text{g ear}^{-1}$) at day 1 (D1) and day 5 (D5) under different boron supply.	90
6.5 The ratio of $^{10}\text{B} : ^{11}\text{B}$ in the ear on the main stem at day 1 (D1) and day 5 (D5) under different boron supply.	90

LIST OF FIGURES

Figure	Page
2.1 Spike at anthesis. (a) Normal spike from +B plant. (b) Sterile spike from -B plant.	9
2.2 Scanning electron micrographs (A,b) and light micrographs (c,d) of wheat pollen towards anthesis.	10
2.3 Stages in development of spring wheat ears, anthers and microspores.	12
2.4 Possible mechanisms of genotypical differences in B efficiency in wheat.	24
3.1 Spike initiation and development.	28
3.2 The effect of B supply on total shoot and root dry weight in two genotypes of wheat grown for 16 days in solution culture. Bars represent means of 3 replications \pm se.	42
4.1 The effect of B on total grain yield, grain yield from the main stem and from tillers in five wheat genotypes.	50
4.2 The effect of B on grain spike ⁻¹ , grain spikelet ⁻¹ and spike plant ⁻¹ in five wheat genotypes.	53
4.3 The effect of B on B concentration in the whole shoot and the root in five wheat genotypes.	57
4.4 The effect of B on B concentration in the spike in five wheat genotypes	57


Figure	Page
4.5 The effect of B on B concentration in the flag leaf, the penultimate leaf, the remainder of the whole shoot and the stem segment between flag leaf and penultimate leaf node (SSFP) in five wheat genotypes.	58
4.6 The effect of B on B content in the whole shoot and the remainder of the whole shoot in five wheat genotypes.	61
4.7 The effect of B on B content in the flag leaf and the penultimate leaf in five wheat genotypes.	61
4.8 The effect of B on the relative distribution in the upper part of the shoot of the main stem in wheat genotypes.	66
5.1 The effect of short term B deficiency (5 days corresponding to the tetrad–young microspore) on pollen viability (%) in two wheat cultivar by fluorochromatic (FCR) test.	75
5.2 The pollen at anthesis by FCR test. (a) viable pollen (b) nonviable	76
5.3 Pollen at anthesis showing the nuclei by DAPI.	76
(a) +B, Fang 60.	
(b) +B, SW 41.	
(c) –B, Fang 60.	
(d) –B, SW 41	

LIST OF PLANT SPECIES

Botanical names of plant species cited in this book.

Apple	<i>Malus sylvestris</i>
Barley	<i>Hordeum vulgare</i>
Beet	<i>Beta vulgaris</i>
Black gram	<i>Vigna mungo</i>
Broccoli	<i>Brassica oleracea</i> L. convar. <i>botrytis</i> var. <i>botrytis</i>
Cauliflower	<i>Brassica oleracea</i> L. convar. <i>botrytis</i> var. <i>italica</i> Plenck
Celery	<i>Apium graveolens</i>
Cotton	<i>Gossypium hirsutum</i>
Grape	<i>Vitis vinifera</i>
Green gram	<i>Vigna radiata</i>
Lentil	<i>Lens culinaris</i>
Lucerne	<i>Medicago sativa</i>
Maize	<i>Zea mays</i>
Mung bean	<i>Phaseolus mungo</i>
Mustard	<i>Brassica juncea</i>
Oil seed rape	<i>Brassica napus</i>
Peanut	<i>Arachis hypogaea</i>
Pear	<i>Pyrus communis</i>
Red clover	<i>Trifolium pratense</i>

Sesame	<i>Sesame indicum</i>
Squash	<i>Cucurbita maxima</i> Duch.
Sumflower	<i>Helianthus annuus</i>
Sunflower	<i>Helianthus annuus</i>
Tobacco	<i>Nicotiana tabaccum</i>
Tomato	<i>Lycopersicon esculentum</i>
Wheat	<i>Triticum aestivum</i>
White clover	<i>Trifolium repens</i>



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
 Copyright© by Chiang Mai University
 All rights reserved

ABBREVIATIONS

ATPase	adenosine triphosphatase
B	boron
D	day
DAPI	4'-6-Diamidino-2-phenylindole 2HCL, Sigma Lot 104F-0542)
FCR	fluorochromatic
G	genotype
GA	gibberellin
GSI	grain set index (%)
IAA	indole acetic acid
KI/I ₂	iodine test
MES	2-[N-Morpholino]ethanesulfonic acid
PPO	polyphenoloxidases
RG-II	rhamnogalacturonan II
SAPL	stem above penultimate leaf node
SAPL	stem below penultimate leaf node
SSFP	stem segment between flag leaf and penultimate leaf node
UV	ultraviolet