#### TABLE OF CONTENTS

	Page	
Acknowledgements	iii	
Abstract in Thai	iv	
Abstract in English	vi	
Table of contents	viii	
List of tables	xi	
List of figures	xvii	
List of appendix tables	xviii	
Chapter 1 Introduction	1	
Objectives of the experiments	2	
Chapter 2 Review of Literature	3	
1. Origin and distribution	3	
2. Botany of longan	3	
3. Flowering process	4	
4. The control of flowering	5	
5. Factors affecting flowering	6	
5.1 External or environmental factors	6	
5.2 Internal or endogeneous factors	8	
6. Potassium chlorate and flowering of longan	15	
6.1 Potassium chlorate property	15	
6.2 Effect of potassium chlorate on plant		
physiology and biochemistry	15	

# TABLE OF CONTENTS (CONTINUED)

		Page
	6.3 Effect of potassium chlorate on flowering of longan	17
	6.4 Effect of potassium chlorate on hormonal	
	changes in longan	19
Chapter 3	Materials and Methods	20
	1. Materials	20
	1.1 Experimental plants	20
	1.2 Others apparatus	21
	2. Methods	21
	2.1 Experiment 1 The concentration effect of potassium chlorate on flower induction of derooted air-layered longan cv. Daw	21
	2.2 Experiment 2 Effect of potassium chlorate on	
	flowering and morphological change of terminal	
	buds in rooted and derooted air-layered longan cv.  Daw	22
	2.3 Experiment 3 Effect of potassium chlorate on physiological and biochemical changes in leaves	
	and shoots of derooted air-layered longan cv.Daw	23
Chapter 4	Results	32
Chapter 5	Discussion	90
Chapter 6	Conclusion	96

#### TABLE OF CONTENTS (CONTINUED)

	Page
References	97
Appendix	108
Curriculum vitae	126

# ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright<sup>©</sup> by Chiang Mai University All rights reserved

#### LIST OF TABLES

Table		Page
1	The effect of potassium concentrations on flower induction of derooted air-layered longan cv. Daw	33
2	The chlorophyll a contents (mg·gFW <sup>-1</sup> ) of leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	41
3	The changing percentage of chlorophyll a (%) of leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	42
4	The chlorophyll b contents (mg·gFW <sup>-1</sup> ) of leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	43
5	The changing percentage of chlorophyll b (%) of leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	44
6	Total chlorophyll contents (mg·gFW <sup>-1</sup> ) of leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of	sity e (
	500 ppm	44

Гable		Page
7	The changing percentage of total chlorophyll (%) of leaves of air-	
	layered (R) and derooted air-layered (DR) longan cv. Daw, non-	
	treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	45
8	Total nonstructural carbohydrate (TNC) (mg D-glucose equivalent·gDW-1)	
	of leaves and shoots of air-layered (R) and derooted air-layered (DR)	
	longan cv. Daw, non-treated and treated with potassium chlorate	
	(+KClO <sub>3</sub> ) at the concentration of 500 ppm	47
9	The changing percentage of total nonstructural carbohydrate	
	(TNC) (%) of leaves and shoots of air-layered (R) and derooted	
	air-layered (DR) longan cv. Daw, non-treated and treated with	
	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	48
10	Total sugar (TS) (mg D-glucose equivalent gDW-1) of leaves and	
	shoots of air-layered (R) and derooted air-layered (DR) longan cv.	
	Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at	
	the concentration of 500 ppm	50
11	The changing percentage of total sugar (TS) (%) of leaves and	
	shoots of air-layered (R) and derooted air-layered (DR) longan cv.	
	Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at	
	the concentration of 500 ppm	51
12	Reducing sugar (RS) contents (mg D-glucose equivalent·gDW-1) of	
12	leaves and shoots of air-layered (R) and derooted air-layered (DR)	
	longan cv. Daw, non-treated and treated with potassium chlorate	52
	(+KClO <sub>3</sub> ) at the concentration of 500 ppm	53

Table		Page
13	The changing percentage of reducing sugar (RS) contents (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR)	
	longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	54
14	The total nitrogen contents (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	57
15	The changing percentage of nitrogen (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
16	concentration of 500 ppm  The carbohydrate per nitrogen ratios (C:N) of air-layered (R) and	58
	derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	59
17 <b>17</b>	The changing percentage of carbohydrate per nitrogen ratios (C:N) (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with	Kl sido
18	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm  The percentage of nitrate (NO <sub>3</sub> <sup>-</sup> ) (%) in leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and	e (
	treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	62

Table		Page
19	The changing percentage of nitrate (NO <sub>3</sub> ) (%) in leaves of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	63
20	The percentage of phosphorus (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	65
21	The changing percentage of phosphorus (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	66
22	The percentage of potassium (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	68
23	The changing percentage of potassium (%) of leaves and shoots of air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	69
24	The percentage of calcium (%) of leaves and shoots of air-layered	
	(R) and derooted air-layered (DR) longan cv. Daw, non-treated	
	and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration	
	of 500 ppm	71

Table		Page
25	The changing percentage of calcium (%) of leaves and shoots of	
	air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	72
26	The nitrate reductase activity (NRA) (µmoleNO <sub>2</sub> -h <sup>-1</sup> ·gFW <sup>-1</sup> ) of	
	leaves of air-layered (R) and derooted air-layered (DR) longan cv.	
	Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at	
	the concentration of 500 ppm	74
27	The changing percentage of nitrate reductase activity (NRA) (%)	
	of leaves of air-layered (R) and derooted air-layered (DR) longan	
	cv. Daw, non-treated and treated with potassium chlorate	
	(+KClO <sub>3</sub> ) at the concentration of 500 ppm	75
28	IAA contents (μM·gFW <sup>-1</sup> ) of air-layered (R) and derooted air-	
	layered (DR) longan cv. Daw, non-treated and treated with	
	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	77
29	The changing percentage of IAA (%) of air-layered (R) and	
	derooted air-layered (DR) longan cv. Daw, non-treated and treated	
		78
	with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	10
30	Gibberellin-like substances contents µg GA3 (Kyowa) equivalent gFW <sup>-1</sup>	
	of air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	80

Table		Page
31	The changing percentage of gibberellin-like substances (GAs) (%)	
	of air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	81
	Carlinia libraria de CVA (a Libraia animia de CVA)	
32	Cytokinin-like substances contents (CKs) (µg kinetin equivalent·gFW <sup>-1</sup> )	
	of air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	84
33	The changing percentage of cytokinin-like substances (CKs) (%)	
	of air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	85
34	The gibberellin-like substance per cytokinin-like substance ratios	
	(GAs:CKs) of air-layered (R) and derooted air-layered (DR)	
	longan cv. Daw, non-treated and treated with potassium chlorate	
	(+KClO <sub>3</sub> ) at the concentration of 500 ppm	86
35	Ethylene contents (ppm) of air-layered (R) and derooted air-	
	layered (DR) longan cv. Daw, non-treated and treated with	
	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	88
36	The changing percentage of ethylene (%) of air-layered (R) and	
	derooted air-layered (DR) longan cv. Daw, non-treated and treated	
	with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	89

#### xvii

#### LIST OF FIGURES

Figure		Page
1	One year-old air-layered longan cv. Daw which used as experimental	
	plants	20
2	Non-shading greenhouse with automatic water sprayer	22
3	The toxic symptoms of potassium chlorate at high concentration	
	on derooted air-layered longan cv. Daw (A:normal; B:leaf drop;	
	C, D and E:chlorosis; F:dehydration)	34
4%	Floral buds of derooted air-layered longan cv. Daw treated with	
	KClO <sub>3</sub> 500 ppm	35
5	The terminal bud of air-layered longan cv. Daw at 10 (A), 15 (B),	
	20 (C) and 25 (D) days, non-treated with KClO <sub>3</sub> (ab = axillary	
	bud, am = apical meristem, yl = young leaf)	36
6	The terminal bud of derooted air-layered longan cv. Daw at 10	
	(A), 15 (B), 20 (C) and 25 (D) days, non-treated with $KClO_3$ (ab	
	= axillary bud, am = apical meristem, yl = young leaf)	37
7	The terminal bud of air-layered longan cv. Daw at 10 (A), 15	
	(B), 20 (C) and 25 (D) days, treated with KClO <sub>3</sub> (ab = axillary	
	bud, am = apical meristem, yl = young leaf, fb = floral bud)	38
8	The terminal bud of derooted air-layered longan cv. Daw at 10	
	(A), 15 (B), 20 (C) and 25 (D) days, treated with KClO <sub>3</sub> (ab =	
	axillary bud, am = apical meristem, yl = young leaf, fb = floral	
	bud) ights reserve	39
9	Floral buds induced by potassium chlorate of derooted air-layered	
	longan at 35-40 DAT	40

#### LIST OF APPENDIX TABLES

Appendix table		Page
1	The chlorophyll a changes (µg·gFW <sup>-1</sup> ) of leaves of air-layered	
	(R) and derooted air-layered (DR) longan cv. Daw, non-treated	
	and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	109
2	The chlorophyll b changes (µg·gFW <sup>-1</sup> ) of leaves of air-layered	
	(R) and derooted air-layered (DR) longan cv. Daw, non-treated	
	and treated with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	110
3	Total chlorophyll changes (µg·gFW <sup>-1</sup> ) of leaves of air-layered	
	(R) and derooted air-layered (DR) longan cv. Daw, non-treated	
	and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	111
4	Total nonstructural carbohydrate (TNC) (mg D-glucose equivalent	
	·gDW <sup>-1</sup> ) changes of leaves and shoots of air-layered (R) and	
	derooted air-layered (DR) longan cv. Daw, non-treated and treated	
	with potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	112
5	Total sugar (TS) (mg D-glucose equivalent gDW <sup>-1</sup> ) changes of	
นสิทธิ	leaves and shoots of air-layered (R) and derooted air-layered	
	(DR) longan cv. Daw, non-treated and treated with potassium	
	bt(C) by Chiang Mai I hivor	113
		113
6	Reducing sugar (RS) contents (mg D-glucose equivalent·gDW <sup>-1</sup> )	
	changes of leaves and shoots of air-layered (R) and derooted air-	
	layered (DR) longan cv. Daw, non-treated and treated with	
	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	114

# LIST OF APPENDIX TABLES (CONTINUED)

Appendix table		Page
7	The total nitrogen content (%) changes of leaves and shoots of	
	air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	115
8	The carbohydrate per nitrogen ratio (C:N) (%) changes of leaves	
	and shoots of air-layered (R) and derooted air-layered (DR)	
	longan cv. Daw, non-treated and treated with potassium chlorate	
	(+KClO <sub>3</sub> ) at the concentration of 500 ppm	116
9	The percentage of nitrate (NO <sub>3</sub> ) changes in leaves of air-layered	
	(R) and derooted air-layered (DR) longan cv. Daw, non-treated	
	and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	117
10	The percentage of phosphorus (%) changes of leaves and shoots	
	of air-layered (R) and derooted air-layered (DR) longan cv.	
	Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> )	
	at the concentration of 500 ppm	118
$G_{11}$	The percentage of potassium (%) changes of leaves and shoots	
	of air-layered (R) and derooted air-layered (DR) longan cv.	
	Daw, non-treated and treated with potassium chlorate (+KClO <sub>3</sub> )	
	at the concentration of 500 ppm	119
12	The percentage of calcium (%) changes of leaves and shoots of	
	air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	120

#### LIST OF APPENDIX TABLES (CONTINUED)

Appendix table		Page
13	The nitrate reductase activity (NRA) ( $\mu$ moleNO <sub>2</sub> -h <sup>-1</sup> ·gFW <sup>-1</sup> )	
	changes of leaves of air-layered (R) and derooted air-layered	
	(DR) longan cv. Daw, non-treated and treated with potassium	
	chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	121
14	IAA changes (μM·gFW <sup>-1</sup> ) of air-layered (R) and derooted air-	
	layered (DR) longan cv. Daw, non-treated and treated with	
	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	122
15	Gibberellin-like substances changes (µg GA <sub>3</sub> (kyowa)·gFW <sup>-1</sup> ) of	
	air-layered (R) and derooted air-layered (DR) longan cv. Daw,	
	non-treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	123
16	Cytokinin-like substances changes (µg kinetin-gFW-1) of air-	
	layered (R) and derooted air-layered (DR) longan cv. Daw, non-	
	treated and treated with potassium chlorate (+KClO <sub>3</sub> ) at the	
	concentration of 500 ppm	124
S 21 317	Ethylene changes (ppm) of air-layered (R) and derooted air-	
	layered (DR) longan cv. Daw, non-treated and treated with	
	potassium chlorate (+KClO <sub>3</sub> ) at the concentration of 500 ppm	125