
TABLE OF CONTENTS

0101013	Page
Acknowledgements	iii
Abstract (English)	v
Abstract (Thai)	ix
List of Tables	xvi
List of Figures	xix
Chapter 1 Introduction	1
Chapter 2 Literature review	4
2.1 Definition of genetic diversity	4
2.2 Definition of local varieties	4
2.3 Structure of genetic diversity in local varieties	5
2.3.1 Allelic richness and genotype diversity	5
2.3.2 Adaptation to the local environment	6
2.3.3 Diversity to meet temporal environmental variation	7
2.3.4 Continuing crop evolutionary process	7
2.4 Threat to local varieties	8
2.5 Farmers management of diversity	149
2.6 Factors affecting genetic diversity of local populations2.7. Rice diversity base on ecosystem in Laos	10
2.7.1 Rice diversity in the rainfed lowland environment	11
2.7.2 Rice diversity in the rainfed upland environment	12

2.8 (Conservation of traditional rice varieties in Laos	13
2.9 I	Measurement of genetic diversity of local rice varieties	13
2	2.9.1 Measurement of morphological and physiological	
	characteristics	14
	2.9.2 Measurement of genetic diversity with DNA level	15
Chapter 3 Ma	terials and Methods	17
3.1	Household survey	17
	3.1.1 Site selection	17
	3.1.2 Methodology	17
	3.1.3 Data analysis	18
3.2	Characterization of local rice varieties	21
	3.2.1 Farmers' seed characterization	21
	3.2.2 Progeny tested	21
	3.2.3 DNA analysis	22
	3.2.4 Data analysis	24
Chapter 4 Res	sults	25
4.1	Field survey	25
ลิขสิทชิ	Characterization of local rice varieties	31
Copyright	.2.1 Farmers' seed characterization .2.2 Progeny test	31
All r	Morphological characteristics S E Morphological characteristics	41
	Physiological characteristics	62
	DNA analysis of local rice varieties	77



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

LIST OF TABLES

Table	0101013	Page
3.1	Description of local rice in the study areas	20
3.2	Primer sequences, repeat motif, expected PCR product size, annealing	
	temperature and chromosomal locations of six microsatellite primers.	23
4.1.1	Number of household, populations, ethnic group and main staple in	
	four villages, Ban Houyman (HM) and Ban Thapho (TP) of Phonxay	
	(PX) district and Ban Ladthahae (LTH) and Ban Houyleung (HL) of	
	Pak Ou (PO) district, Luang Prabang province	26
4.1.2	Distribution and percentage (in parenthesis) of number rice varieties,	
	ecosystem, endosperm type and maturity of rice varieties in four	
	villages	27
4.1.3	Rank and proportion of rice varieties (no of household) in four villages	28
4.1.4	Number of rice varieties per household, rice seed exchange and	
	storage in four villages	30
4.2.1	Husk color of rice varieties in farmers' seed lots in four villages	33
4.2.2	Pericarp color of rice varieties in farmers' seed lots in four villages	34
4.2.3	Percentage of non-glutinous, glutinous and Shannon-Weaver Index	itv
	(H') of rice varieties in farmers' seed lots in four villages	35
4.2.4	Seed length of rice varieties in farmers' seed lots in four villages	36
4.2.5	Seed width of rice varieties in farmers' seed lots in four villages	37
426	100 seeds weight of rice varieties in farmers' seed lots in four villages	38

	4.2.7	Leaf blade color of rice varieties in progeny test in four villages	45
	4.2.8	Basal leaf sheath color of rice varieties in progeny test in four villages	46
	4.2.9	Leaf blade pubescence of rice varieties in progeny test in four villages	47
	4.2.10	Auricle color of rice varieties in progeny test in four villages	48
	4.2.11	Collar color of rice varieties in progeny test in four villages	49
	4.2.12	Ligule color of rice varieties in progeny test in four villages	50
	4.2.13	Ligule shape of rice varieties in progeny test in four villages	51
	4.2.14	Stigma color of rice varieties in progeny test in four villages	52
	4.2.15	Apicullus color of rice varieties in progeny test in four villages	53
	4.2.16	Pericarp color of rice varieties in progeny test in four villages	54
	4.2.17	Husk color of rice varieties in progeny test in four villages	55
	4.2.18	Grain awning of rice varieties in progeny test in four villages	57
	4.2.19	Husk pubescence of rice varieties in progeny test in four villages	58
	4.2.20	Days to flowering of rice varieties in progeny test in four villages	63
	4.2.21	Number of tillers per plant of rice varieties in progeny test in four	
		villages	65
	4.2.22	Culm length (cm) of rice varieties in progeny test in four villages	67
a	4.2.23	Number of panicles per plant of rice varieties in progeny test in four	11
	NOV/K	villages Chiang Mai I Iniversi	69
^	4.2.24	Panicle length (cm) of rice varieties in progeny test in four villages	71
A	4.2.25	Number of seeds per panicle of rice varieties in progeny test in four	C
		villages	73
	1226	Percent of filled seeds of rice varieties in progeny test in four villages	75

4.2.27	Number of alleles of six microsatellite loci, of twelve samples of five	
	local rice varieties collected from different villages	77
4.2.28	Gene diversity per locus for six markers loci, of twelve samples of five	
	local rice varieties collected from different villages	78
4.2.29	Genetic parameter for samples of twelve samples of five local rice	
	varieties collected different villages	80
4.2.30	Genetic parameter for varieties of twelve samples of five local rice	
	varieties collected different villages	81
4.2.31	Genetic parameter for villages of twelve samples of five local rice	
	varieties collected different villages	82
	The state of the s	

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

Figure	0101013	Page
3.1	Location of Luang Prabang provinces in Lao PDR	19
4.1.1	Percent of household planted popular varieties in Ban Houyman (HM) and	
	Ban Thapho (TP) of Phonxay (PX) district, Luang Prabang province	29
4.1.2	Percent of household planted popular varieties in Ban Ladthahae (LTH) and	
	Ban Houyleung (HL) of Pak Ou (PO) district, Luang Prabang province	29
4.2.1	Average of seed length and coefficient of variance of rice varieties in	
	farmers' seed lots in four villages, Ban Houyman (HM) and Ban Thapho	
	(TP) of Phonxay (PX) district and Ban Ladthahae (LTH) and Ban	
	Houyleung (HL) of Pak Ou (PO) district, Luang Prabang province	39
4.2.2	Average of seed width and coefficient of variance of rice varieties in	
	farmers' seed lots	39
4.2.3	Grain shapes of rice varieties in four villages in farmers' seed lots	40
4.2.4	Variant distance clustering by UPGMA methods showing morphological	
	relationship among samples of local rice varieties	59
4.2.5	Morphological characteristics of local rice varieties. Local rice varieties	
Copy	exhibited color of leaf blade, Basal leaf sheath, collar and auricle	60
4.2.6	Morphological characteristics of local rice varieties. Local rice varieties	
N II II	exhibited color of husk, apiculus, awning and pericarp	61
4.2.7	Distribution of days to flowering of rice varieties in progeny test	64
4.2.8	Distribution of number to tillering of rice varieties in progeny test	66

4.2.9	Distribution of culm length (cm) of rice varieties in progeny test	68
4.2.10	Distribution of number of panicle per plant of rice varieties in progeny test	70
4.2.11	Distribution of panicle length (cm) of rice varieties in progeny test	72
4.2.12	Distribution of number of seed per panicle of rice varieties in progeny test	74
4.2.13	Distribution of percent of filled seeds of rice varieties in progeny test	76
4.2.14	Microsatellite amplification product from of twelve samples of five local	
	rice varieties genomic DNA with primer RM316.	78
4.2.15	Genetic distance clustering by UPGMA methods showing genetic	
	relationship among samples of local rice varieties	83

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved