


TABLE OF CONTENTS

Acknowledgements	iii
Abstract in English	v
Abstract in Thai	ix
Table of Contents	xiii
List of Tables	xvi
List of Figures	xxii
List of Appendices	xxv
CHAPTER 1 INTRODUCTION	1
CHAPTER 2 LITERATURE REVIEW	4
2.1 Classification of <i>Oryza</i> species	4
2.2 Identification of wild species	6
2.3 Asian common wild rice (<i>O. rufipogon</i>)	6
2.3.1 Habitat of common wild rice	6
2.3.2 Genetic diversity of common wild rice	7
2.3.3 Gene flow	9
2.3.4 The importance of wild rice growth stage	9
2.4 Germplasm and conservation of common wild rice	12
2.5 Genetic diversity used for microsatellite analysis	13
2.6 Usefulness of wild rice in plant breeding program	14

CHAPTER 3 MATERIALS AND METHODS	15
3.1 Genetic diversity of common wild rice in Cambodia	15
3.1.1 Survey and collection of common wild rice populations in Cambodia	15
3.1.2 Genetic structure of common wild rice using microsatellite markers	19
3.2 Characterization of F ₂ populations between cultivated rice and common wild rice	24
3.2.1 Genetic materials	24
3.2.2 Experimental procedure	26
3.2.3 Data analysis	26
CHAPTER 4 RESULTS	27
4.1 Genetic diversity of common wild rice populations in Cambodia	27
4.1.1 Survey of common wild rice.	27
4.1.2 Genetic structure of common wild rice population	33
4.2 Morphological and physiological characteristic of parent and F ₂ segregation analysis.	49
4.2.1 Morphological characterization of plants and seed of F ₂ segregation and parents.	49
4.2.2 Morphological segregation of F ₂ s	53
4.2.3 Physiological characteristics of F ₂ segregation population analysis	62

CHAPTER 5 DISCUSSION	77
CHAPTER 6 CONCLUSION	82
REFERENCES	84
APPENDICES	93
CURRICULUM VITAE	105



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF TABLES

Table	Page
2.1 <i>Oryza</i> species with their chromosome number, genome group and geographical distribution	5
3.1 Description of common wild rice collected from four regions in Cambodia	18
3.2 Description of cultivated rice collected from three regions in Cambodia used in this study	18
3.3 Primer sequences, repeat motif, expected PCR product size, annealing temperature and chromosomal locations of five microsatellite primers	20
3.4 Description of cultivated rice varieties and common wild rice populations were used in the experiment	24
4.1 Ecotype, habitat, and morphological character at anthesis of 10 common wild rice populations survey in Cambodia	29
4.2 Ecotype, habitat, and morphological character of 7 cultivated rice survey in Cambodia	31
4.3 Number of alleles samples of 10 common wild rice populations and 7 local rice varieties collected from different province, based on five microsatellite loci.	34

4.4	Allele frequencies of 7 local rice varieties and 10 wild rice populations collected from four locations in Cambodia based on microsatellite marker (RM 20)	37
4.5	Allele frequencies of 7 local rice varieties and 10 wild rice populations collected from four locations in Cambodia based on microsatellite marker (RM 164)	38
4.6	Allele frequencies of 7 local rice varieties and 10 wild rice populations collected from four locations in Cambodia based on microsatellite marker (RM 225)	39
4.7	Allele frequencies of 7 local rice varieties and 10 wild rice populations collected from four locations in Cambodia based on microsatellite marker (RM 341)	40
4.8	Allele frequencies of 7 local rice varieties and 10 wild rice populations collected from four locations in Cambodia based on microsatellite marker (RM 588)	41
4.9	Effective numbers of alleles (A_e), Average number of alleles per locus (N_a), Expected heterozygosity (H_e), Observed heterozygosity (H_o), Inbreeding coefficient (F_{is}), Out-crossing rate (t) of populations revealed by microsatellite.	43
4.10	Genetic diversity estimated for 10 common wild rice population in	

Three locations in Cambodia	45
4.11 Genetic differentiation among (F_{ST}) matrix values determined across five microsatellites loci in ten common wild rice populations in Cambodia.	46
4.12 Proportion of inferred cluster (Q) of 7 cultivated rice and 10 common wild rice populations to each inferred population ($K=2$)	48
4.13 Morphological characterization of plants and seeds of parents and F_2 segregation population	50
4.14 Plant type of F_2 populations derived from crosses between cultivated and wild rice	57
4.15 Leaf-blade color of F_2 population derived from crosses between cultivated and wild rice	57
4.16 Leaf-sheath color of F_2 population derived from crosses between cultivated and wild rice	58
4.17 Auricle color of F_2 population derived from crosses between cultivated and wild rice	58
4.18 Ligule color of F_2 population derived from crosses between cultivated and wild rice	58
4.19 Internodes color of F_2 population derived from crosses between cultivated and wild rice	59

4.20	Stigma color of F ₂ population derived from crosses between cultivated and wild rice	59
4.21	Apiculus color of F ₂ population derived from crosses between cultivated and wild rice	59
4.22	Presentation on awn of F ₂ population derived from crosses between cultivated and wild rice	60
4.23	Awn color of F ₂ population derived from crosses between cultivated and wild rice	60
4.24	Hull color of F ₂ population derived from crosses between cultivated and wild rice	60
4.25	Panicle type of F ₂ population derived from crosses between cultivated and wild rice	61
4.26	Pericarp color of F ₂ population derived from crosses between cultivated and wild rice	61
4.27	Range, mean and standard deviation (Sd) of tillers per plant of Parents and F ₂ populations from crosses between cultivated rice (IR66) x wild rice from Kampong Thom province (KTWR), and Sen pidao (SPD) x wild rice from Takeo province (TKWR)	62
4.28	Range, mean and standard deviation (Sd) of plant height (cm) of Parents and F ₂ populations from crosses between cultivated rice	

- (IR66) x wild rice from Kampong Thom province (KTWR), and Sen
pidao (SPD) x wild rice from Takeo province (TKWR) 64
- 4.29 Range, mean and standard deviation (Sd) of days to flowering of
Parents and F₂ populations from crosses between cultivated rice
(IR66) x wild rice from Kampong Thom province (KTWR), and Sen
pidao (SPD) x wild rice from Takeo province (TKWR) 65
- 4.30 Range, mean and standard deviation (Sd) of panicle length (cm) of
Parents and F₂ populations from crosses between cultivated rice
(IR66) x wild rice from Kampong Thom province (KTWR), and Sen
pidao (SPD) x wild rice from Takeo province (TKWR) 67
- 4.31 Range, mean and standard deviation (Sd) of branch panicle number
of parents and F₂ populations from crosses between cultivated rice
(IR66) x wild rice from Kampong Thom province (KTWR), and Sen
pidao (SPD) x wild rice from Takeo province (TKWR) 68
- 4.32 Range, mean and standard deviation (Sd) of spikelet per panicle of
parents and F₂ populations from crosses between cultivated rice
(IR66) x wild rice from Kampong Thom province (KTWR), and Sen
pidao (SPD) x wild rice from Takeo province (TKWR) 70
- 4.33 Range, mean and standard deviation (Sd) of seed fertility (%) of
parents and F₂ populations from crosses between cultivated rice


	(IR66) x wild rice from Kampong Thom province (KTWR), and Sen pidao (SPD) x wild rice from Takeo province (TKWR)	71
4.34	Range, mean and standard deviation (Sd) of seed shattering (%) of parents and F ₂ populations from crosses between cultivated rice (IR66) x wild rice from Kampong Thom province (KTWR), and Sen pidao (SPD) x wild rice from Takeo province (TKWR)	73
4.35	Range, mean and standard deviation (Sd) of seed width (mm) of parents and F ₂ populations from crosses between cultivated rice (IR66) x wild rice from Kampong Thom province (KTWR), and Sen pidao (SPD) x wild rice from Takeo province (TKWR)	75
4.36	Range, mean and standard deviation (Sd) of seed length (mm) of parents and F ₂ populations from crosses between cultivated rice (IR66) x wild rice from Kampong Thom province (KTWR), and Sen pidao (SPD) x wild rice from Takeo province (TKWR)	75

LIST OF FIGURES

Figure	Page
3.1 Collection sites of common wild rice in Cambodia	17
3.2 Collection sites of two common wild rice biotypes crossed with two improved rice varieties.	25
4.1 Views of habitat of common wild rice populations collected in October 16-25, 2007.	32
4.2 A UPGMA dendrogram based on Nei's distance showing genetic relationships among the 10 populations of common wild rice in Cambodia.	46
4.3 Population structures of common wild rice and cultivated rice collected from 4 locations (Phnom Penh, Kandal, Takeo and Prey Veng) of Cambodia was assumed $K=2$.	47
4.4 Morphological characteristics of plant type, internode color, leaf sheath color, anther length and pericarp color of parent and F_2 segregation population.	52
4.5 Distribution of number of tillers per plant of F_2 populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	63
4.6 Distribution of number of plant height (cm) of F_2 populations	

	between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	64
4.7	Distribution of number of days to flowering of F ₂ populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	66
4.8	Distribution of number of panicle length of F ₂ populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	67
4.9	Distribution of number of branch panicle of F ₂ populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	69
4.10	Distribution of number of spikelet panicle of F ₂ populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	70
4.11	Distribution of number of seed fertility of F ₂ populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	72
4.12	Distribution of number of seed shattering of F ₂ populations between Sen pidao (SPD) x wild rice from Takeo province (TKWR) and IR66 x wild rice from Kampong Thom province (KTWR).	73

- 4.13 Distribution of seed shape of F_2 populations between cultivated rice
(Sen pidao) x wild rice (Takeo) compared with their parents. 76
- 4.14 Distribution of seed shape of F_2 populations between cultivated rice
(IR66) x wild rice (Kompong Thom) compared with their parents. 76



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

LIST OF APPENDICES

App.		Page
1	Wild rice survey sheet	93
2	Descriptor state and Meaning of morphological and other characters of common wild rice	94
3	DNA extraction	95
4	PCR Processing	97
5	Polyacrylamine Gel Electrophoresis Preparing (PAGE)	98
6	Band patterns of 10 wild rice populations and 7 local rices at RM20 lucus	100
7	Band patterns of 10 wild rice populations and 7 local rices at RM164 lucus	101
8	Band patterns of 10 wild rice populations and 7 local rices at RM225 lucus	102
9	Band patterns of 10 wild rice populations and 7 local rices at RM341 lucus	103
10	Band patterns of 10 wild rice populations and 7 local rices at RM588 lucus	104