

REFERENCES

- Adoukonou-Sagbadja H., Wagner C., Dansi A., Ahlemeyer J., Dainou O., Akpagana K., Ordan F. And W. Friedt. 2007. Genetic diversity and population differentiation of traditional fonio millet (*Digitaria* spp.) landraces from different agro-ecological zones of West Africa. *Theor Appl Genet* 115:917-931.
- Akagi H., Yokozeki Y., Inagaki A., and T. Fujimura. 1996. Microsatellite DNA markers for rice chromosomes. *Theor Appl Genet* 93:1071-1077.
- Allinne C., Mariac C., Vigouroux Y., Bezancon G., Couturon E., Moussa D., Tidjani M., Pham J.L. and T. Robert. 2007. Role of seed flow on the pattern and dynamics of pearl millet (*Pennisetum glaucum* [L.] R. Br.) genetic diversity assessed by AFLP markers: a study in south-western Niger. *Genetica* DOI 10.1007/s10709-007-9197-7.
- Almekinders C.J.M., Louwaars N.P. and G.H. de Bruijn. 1994. Local seed systems and their importance for an improved seed supply in developing countries. *Euphytica* 78:207-216.
- Alvarez N., Garine E., Khasah C., Dounias E., Hossaert-McKey M. and D. McKey. 2005. Farmers' practices, metapopulation dynamics, and conservation of agricultural biodiversity on-farm: a case study of sorghum among the Duupa in sub-sahelian Cameroon. *Biol Conserv* 121:533-543.
- Bajracharya J., Steele K.A., Jarvis D.I., Sthapit B.R. and J.R. Witcombe. 2006. Rice landrace diversity in Nepal: variability of agro-morphological traits and SSR markers in landraces from a high-altitude site. *Field Crop Res* 95:327-335.

- Baker H.G. 1974. The evolution of weeds. *Ann Rev Ecol Syst* 5:1-24
- Balma D., Baniya B.K., Bramel-Cox P., De N.N., Djimadoum M., Ellis E., Ha N.P., Hodgkin T., Louette D., Mellas H., Nassif F., Ndung'u-Skilton J., Ortega-Paczka R., Rana R.B., Sawasogo M. Tripp R. and F.R. Valdivia. 2000. Chapter 6 Seed system. pp 95-106. In Jarvis D.I. Myer L. Klemick H. Guarino L. Smale, M. Brown A.H.D. Sadiki M. Sthapit B. and T. Hodgkin. (Eds.) *A training guide of in situ conservation on-farm. Version 1*. International Plant Genetic Resources Institute, Rome, Italy.
- Barbier P. 1989a. Genetic variation and ecotypic differentiation in the wild rice species *Oryza rufipogon*. I. Population differentiation in life-history traits and isozymic loci. *Jpn J Genet* 64: 259–271.
- Barbier P. 1989b. Genetic variation and ecotypic differentiation in the wild rice species *Oryza rufipogon*. II. Influence of the mating system and life-history traits on the genetic structure of populations. *Jpn J Genet* 64: 273–285.
- Barnaud A., Deu M., Garine E., McKey D and I. Joly. 2007. Local genetic diversity of sorghum in village in northern Cameroon: structure and dynamics of landraces. *Theor Appl Genet* 114:237-248.
- Beebe S., Toro Ch. O., Gonzalez A. V., Chacon M.I., and D.G. Debouck. 1997. Wild-weed-crop complexes of common bean (*Phaseolus vulgaris* L., Fabaceae) in the Andes of Peru and Columbia, and their implications for conservation and breeding. *Genet Resour Crop Evol* 44:73-91.
- Bell R. and V. Seng. 2004. Rainfed lowland rice-growing soils of Cambodia, Laos, and North-east Thailand. P. 161-173. In Seng V., Craswell E., Fukai S. and K.

Fischer. (Eds). *Water in agriculture*. ACIAR Proceedings No. 116e (printed version published in 2004).

Bellon M. and S.B. Brush. 1994. Keepers of maize in Chiapas, Mexico. *Econ Bot* 48:196–209.

Bellon M.R. 1997. Understanding farmer seed and variety systems On-farm conservation as a process: an analysis of its components. In *Using diversity enhancing and maintaining genetic resources on-farm*, eds. Sperling L, Loevinsohn M (ISBN 0-88936-833-3).

Bohonak J.A. 2002. IBD (isolation by distance): a program for analyses of isolation by distance. *J Hered* 93:153-154.

Bracco M., Lia V.V., Gottlieb A.M., Hernandez C.J. and Poggio L. 2009. Genetic diversity in Maize landraces from indigenous settlements of Northeastern Argentina. *Genetica* 135:39-49.

Brown A.H.D. 1978. Isozymes, plant population genetics structure and genetic conservation. *Theor Appl Genet* 52:145-157.

Brown A.H.D. 2000. Chapter 2 The genetic structure of crop landrace and the challenge to Conserve them *in situ* on farm. P.29-49. In Brush, S.B. (Ed), *Gene in the field: On-Farm Conservation of Crop Diversity*, Lewis Publishing Co., USA.

Brush S.B. 1995. *In situ* conservation of landraces in centers of crop diversity. *Crop Science* 35:346-354.

Buckler E. IV, Thornsberry J., and S. Kresovich. 2001. Molecular diversity, structure and domestication of grasses. *Genet Res Camb* 77:213-218.

Burger C.J., Chapman A.M., and J.M. Burke. 2008. Molecular insights into the evolution of crop plants. *Am J Bot* 95(2):113-122.

- Cai W.H., Wang K.X. and H. Morishima. 2004. Comparison of population genetic structures of common wild rice (*Oryza rufipogon* Griff.), as revealed by analyses of quantitative traits, allozymes and RFLPs. *Heredity* 92:409-417.
- Cai W. and H. Morishima. 2002. QTL cluster reflect character association in wild and cultivated rice. *Theor Appl Genet* 104:1217-1228.
- Caicedo A. and B. Schaal. 2004. Population structure and phylogeography of *Solanum pimpinellifolium* inferred from a nuclear gene. *Mol Ecol* 13: 1871-1882.
- Cao Q., Lu B.R., Xia H., Rong J., Sala F., Spada A., and F. Grassi. 2006. Genetic diversity and origin of weedy rice (*Oryza sativa* f. *spontanea*) population found in North-eastern China revealed by simple sequence repeat (SSR) markers. *Ann Bot* 98:1241-1252.
- Cavalli-Sforza L.L. and A.W.F. Edwards. 1967. Phylogenetic analysis: models and estimation procedures. *Evolution* 21(3):550-570.
- Ceccarelli, S. 1994. Specific Adaptation and Breeding for Marginal Conditions. *Euphytica* 77:205-219.
- Chang T.T. 1976. The origin, evolution, cultivation, dissemination, and diversification of Asian and African rices. *Euphytica* 25:425-441.
- Chase M. W. and H. G. Hills. 1991. Silica gel: An ideal material for field preservation of leaf samples for DNA studies. *Taxon*. 40:215-220.
- Chen X., Temnykh S., Xu Y., Cho G.Y. and S.R. McCouch. 1997. Development of a microsatellite framework map providing genome-wide coverage in rice (*Oryza sativa* L.). *Theor Appl Genet* 95:553-567.

- Cheng C.Y., Motohashi R., Tsuchimoto S., Fukuta Y., Ohtsubo H., and E. Ohtsubo. 2003. Polyphyletic origin of cultivated rice based on the interspersed pattern of SINEs. *Mol Biol Evol* 20:67-75.
- Chitrakon S. 1995. Characterization, evaluation and utilization of wild rice germplasm in Thailand. Pathum Thani Rice Research Center. Thonburi, Pathum Thani, Thailand. 143 p.
- Clark R.M., Linton E., Messing J., and F. Doebley. 2004. Pattern of diversity in the genomic region near the maize domestication gene *tb1*. *Proc Natl Acad Sci USA* 101(3):700-707.
- Cureton A., Newbury H., Raybould A., and B. Ford-Lloyd. 2006. Genetic structure and gene flow in wild beet populations: the potential influence of habitat on transgene spread and risk assessment. *J Appl Ecol* 43:1203-1212.
- De Wet J.M.J and J.R. Harlan. 1975. Weeds and domesticates: evolution in the man-made habitat. *Econ Bot* 29:99-107.
- Delouche J.C., Burgos N.R., Gealy D.R., de San Martin D.Z., Labrada R., Larinde M. and C. Rosell. 2007. Weedy Rices – Origin, Biology, Ecology, and Control, *FAO Plant Production and Protection Paper 188*, FAO, Rome (2007) 144 pp.
- Dennis J.V. 1987. Farmer Management of Rice Variety Diversity in Northern Thailand. Ph.D. dissertation, Cornell University.
- Doebley J.F., Gaut B.S., and B.D. Smith. 2006. The molecular genetics of crop domestication. *Cell* 127:1309-1321.
- Doyle J.J. and J.L. Doyle. 1987. A rapid DNA isolation procedure from small quantities of fresh leaf tissues. *Phytochem Bull.* 19:11-15.

- Elias M., Muhlen S.G., McKey D., Roa C. and J. Tohme. 2004. Genetic diversity of traditional south American landraces of cassava (*Manihot Esculenta* Crantz): an analysis using microsatellites. *Econ Bot* 58(2):242-256.
- Ellstrand N.C., Prentice H.C. and J.F. Hancock. 1999. Gene flow and introgression from domesticated plants into their wild relatives. *Annu Rev Ecol Syst* 30:539-563.
- Epperson K.B. and T. Li. 1996. Measurement of genetic structure within populations using Moran's spatial autocorrelation statistics. *Proc Natl Acad Sci USA* 93:10528-10532.
- Excoffier L., Smouse E.P., and M.J. Quattro. (1992) Analysis of molecular variance inferred from metric distance among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131:479-491
- Eyre-Walker A.R., Gaut R.L., Hilton H., Feldman D.L., and B.S. Gaut. 1998. Investigation of the bottleneck leading to the domestication of maize. *Proc Natl Acad Sci USA* 95:4441-4446.
- Fukunaga K., Hill J., Vigouroux Y., Matsuoka Y., Jesus S. G., Liu K., Buckler S.E. and J. Doebley. 2005. Genetic diversity and population structure of teosinte. *Genetics* 169:2241-2254.
- Fukuoka S., Suu D.T., Ebana K., Trinh N.L. Nagamine T. and K. Okuno. 2006. Diversity in phenotypic profiles in landrace populations of Vietnamese rice: A case study of agronomic characters for conserving crop genetic diversity on farm. *Genet Resour Crop Evol* 53:753-761.

- Gao L.Z., Ge S., and D.Y. Hong. 2000a. Allozymic diversity and genetic structure of common wild rice *Oryza rufipogon* Griff., China. *Theor Appl Genet* 101:494–502.
- Gao L.Z., Ge S., Hong D.Y., Chen W., Jiang W.Z. and X.K. Wang 2000b. Genetic erosion in northern marginal population of common wild rice *Oryza rufipogon* Griff. and its conservation, revealed by allozyme analysis. *Hereditas* 133:47–53.
- Gao L.Z., Schaal B.A., Zhang C.H., Jia J.Z., and Y.S. Dong. 2002. Assessment of population genetic structure of common wild rice *Oryza rufipogon* Griff. detected by microsatellite DNA and allozyme loci. *Theor Appl Genet* 106:173–180.
- Gao L.Z. 2004. Population structure and conservation genetics of wild rice *Oryza rufipogon* (Poaceae): a region-wide perspective from microsatellite variation. *Mol Ecol* 13:1009–1024.
- Garris A.J., Tai T.H., Coburn J., Kresovich S. and S. McCouch. 2005. Genetic structure and diversity in *Oryza sativa* L. *Genetics* 169:1631-1638.
- Ge S., Oliveira G.C.X., Schaal B.A., Gao L.Z. and D.Y. Hong. 1999. RAPD variation within and between natural populations of the wild rice *Oryza rufipogon* from China and Brazil. *Heredity* 82:638–644.
- Gealy D.R., Tai T.H. and C.H. Sneller. 2002. Identification of red rice, rice, and hybrid populations using microsatellite markers. *Weed Science* 50:333-339.
- Goldringer I., Enjalbert J., Raquin A.L., and P. Brabant. 2001. Strong selection in wheat populations during ten generations of dynamic management. *Genet Sel Evol* 33:441-463.

- Gomez J.O., Blair W.M., Frankow-Lindberg E.B., and U. Gullberg. 2004. Molecular and phenotypic diversity of common bean landraces from Nicaragua. *Crop Sci* 44:1412-1418
- Goudet J. 2001. FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). <http://www.unil.ch/izea/software/fstat.html>.
- Gravuer K., von Wettberg E., and J. Schmitt. 2005. Population differentiation and genetic variation inform translocation decisions for *Liatris scariosa* var. *novae-angliae*, a rare New England grassland perennial. *Biol Conserv* 124:155-167.
- Gray A. 1996. Genetic diversity and its conservation in natural populations of plants. *Biodiversity letters* 3:71-80.
- Gu X.Y., Kianian S.F., and M.E. Foley. 2006. Dormancy genes from weedy rice respond divergently to seed development environments. *Genetics* 172:1199-1211.
- Hamrick J.L., Linhart Y.B. and J.B. Mitton. 1979. Relationships between life history characteristics and electrophoretically detectable genetic variation in plants. *Ann Rev Ecol Syst* 10:173-200.
- Harlan J.R. and J.M.J. de Wet. 1971. Towards a rational classification of cultivated plants. *Taxon* 20:509-517.
- Harlan J.R. 1992. *Crop and Man*. 2nd edition. Madison. Wisconsin, USA.
- He X.Q., Liu Q.C., Ishiki K., Zhai H., and Y.P. Wong. 2006. Genetic diversity and genetic relationships among Chinese sweetpotato landraces revealed by RAPD and AFLP markers. *Breed Sci* 56:201-207

- Hyten D., Song Q., Zhu Y., Choi I.Y., Nelson R.L., Costa J.M., Specht J.E., Shoemaker R.C. and P.B. Cregan. 2006. Impact of genetic bottlenecks on soybean genome diversity. *Proc Natl Acad Sci USA* 103:16666-16671
- Jaradat A.A., Shahid M. and A.Y. Al Maskri. 2004. Genetic diversity in the batini barley landrace from Oman: I. spike and seed quantitative and qualitative traits. *Crop Sci.* 44:304-315.
- Jarvis D.I. and T. Hodgkin. 1999. Wild relatives and crop cultivars: detecting natural introgressions and farmer selection of new genetic combinations in agroecosystems. *Mol Ecol* 8:S159-S173.
- Javis D.I., Brown A.H.D., Cuong P.H., Collado-Pandura L., Latournerie-Moreno L., Gyawali S., Tanto T., Sawadogo M., Mar I., Sadiki M., Hue N.T.N., AriasReyes L., Balma D., Bajracharya J., Castillo F., Rijal D., Belqadi L., Rana R., Saidi S., Ouedraogo J., Zangre R., Phrib K., Chavez J.L., Schoen D., Sthapit B., Santis P.D., Fedda C. and T. Hodgkin. 2008. A global perspective of the richness and evenness of traditional crop-variety diversity maintained by farming communities. *Proc Natl Acad Sci USA* 105(14):5326-5331.
- Keller L.F. and D.M. Waller. 2002. Inbreeding effects in wild populations. *Trends Ecol Evol* 17(5): 230-241.
- Khush G.S. 1997. Origin, dispersal, cultivation and variation of rice. *Plant Mol. Biol.* 35:25-34.
- Kimura M. and G.H. Weiss. 1964. The stepping stone model of population structure and the decrease of genetic correlation with distance. *Genetics* 49:561-576.
- Kumar S., Tamura K., Jakobsen B.I., and M. Nei. 2001. MEGA2: molecular evolutionary genetics analysis software. *Bioinformatics* 17:1244-1245.

- Kuroda Y., Sato Y.I., Bounphanousay C., Kono Y. and K. Tanaka. 2007. Gene flow from cultivated rice (*Oryza sativa* L.) to wild *Oryza* species (*O. rufipogon* Griff. and *O. nivara* Sharma and Shastry) on the Vientiane plain of Laos. *Euphytica* 142:75-83 DOI:10.1007/s10681-005-0529-3.
- Langevin S.A., Clay K. and Grace J.B. 1990. The incidence and effects of hybridization between cultivated rice and its related weed red rice (*Oryza sativa* L.). *Evolution* 44(4):1000-1008.
- Li C., Zhou A. and T. Sang. 2006. Genetic analysis of rice domestication syndrome with the wild annual species, *Oryza nivara*. *New Phytologist* 170:185-194.
- Li Z., Zhang H., Zeng Y., Yang Z., Shen S., Sun C. and X Wang. 2002. Studies on sampling schemes for the establishment of core collection of rice landraces in Yunnan, China. *Genet Resour Crop Evol* 49:67-74.
- Linhart Y.B. and M.C. Grant. 1996. Evolutionary significance of local genetic differentiation in plants. *Ann Rev Ecol Syst* 27:237-77
- Liu A. and J.M. Burke. 2006. Patterns of nucleotide diversity in wild and cultivated sunflower. *Genetics* 173:321-330.
- Liu K. and S.V. Muse. 2005. POWERMARKER: integrated analysis environment for genetic marker data. *Bioinformatics* 21:2128-2129.
- Londo J.P. and B.A. Schaal. 2007. Origin and population genetics of weedy red rice in the USA. *Mol Ecol* 16:4523-4535.
- Londo J.P., Chiang Y., Hung K., Chiang T. and B.A. Schaal. 2006. Phylogeography of Asian wild rice, *Oryza rufipogon*, reveals multiple independent domestications of cultivated rice, *Oryza sativa*. *Proc Natl Acad Sci USA* 103:9578 – 9583.

- Louette D. and M. Smale. 2000. Farmers' seed selection practices and traditional maize varieties in Cuzalapa, Mexico. *Euphytica* 113:25-41.
- Louette D., Charrier A. and J. Berthaud. 1997. *In situ* conservation of maize in Mexico: genetic diversity and maize seed management in a traditional community. *Econ Bot* 51:20-38.
- Loveless M.D. and J.L. Hamrick. 1984. Ecological determinants of genetic structure in plant populations. *Ann Rev Ecol and Syst* 15:65-95.
- Maneechote C., Jamjod S. and B. Rerkasem. 2004. Invasion of weedy rice in rice fields in Thailand: problems and management. *IRRN* 29(2):20-22.
- Martinez-Castillo J., Zizumbo-Villarreal D., Gepts P., Delgado-Valerio P., and P. Colunga-GarciaMarin. 2006. Structure and genetic diversity of wild population of lima bean (*Phaseolus lunatus* L.) from the Yucatan Peninsula, Mexico. *Crop Sci* 46:1071-1080.
- McCouch S.R., Teytelman L., Xu Y., Lobos K.B., Clare K., Walton M., Fu B., Maghirang R., Li Z., Xing Y., Zhang Q., Kono I., Yano M., Fjellstrom R., DeClerck G., Schneider D., Cartinhour S., Ware D. and L. Stein. 2002. Development and mapping of 2240 new SSR markers of rice (*Oryza sativa* L.). *DNA Research* 9:199-207.
- McCouch, S.R. 2004. Diversifying selection in plant breeding. *PloS Biology* 2 (10):1507-1512.
- Meesin, S. 2003. The structure of genetic diversity in a local Thai rice germplasm. MSc (Agronomy), Graduate School, Chiang Mai University Thailand
- Mengoni, A., Gonnelli, C., Galardi, F., Gabbrielli, R. and M. Bazzicalupo. 2000. Genetic diversity and heavy metal tolerance in populations of *Silene paradoxa*

L.(Caryophyllaceae): a random amplified polymorphic DNA analysis. *Mol Ecol* 9:1319-1324.

Mohammadi S.A. and B.M. Prasanna. 2003. Analysis of genetic diversity in crop plants-salient statistical tools and considerations. *Crop Sci* 43:1235-1248.

Morishima, H. and H.I. Oka. 1970. A survey of genetic variations in the populations of wild *Oryza* species and their cultivated relatives. *Jpn J Genet* 45:371-385.

Morishima, H. and P. Barbier. 1990. Mating system and genetic structure of natural populations in wild rice *Oryza rufipogon*. *Plant Species Biol* 5:31-39.

Neeraja, N.C., Hariprasad, S.A., Malathi, S. and A.E. Siddiq. (2005) Characterization of tall landraces of rice (*Oryza sativa* L.) using gene-derived simple sequence repeats. *Curr Sci* 88:149-152.

Nei, M. 1973. Analysis of gene diversity in subdivided populations. *Proc Nat Acad Sci USA* 70 (12):3321-3323.

Niruntrayakul, S. 2008. Gene flow between cultivated and wild rice. Ph.D. Thesis, Graduate School, Chiang Mai University, Chiang Mai, 143 p.

Noldin, J.S. 2000. Red rice status and management in the Americas. In Wild and weedy rice in rice ecosystem in Asia-a review. IRRN. pp21-24.

OAE (1998) Report on the survey of main season rice, 1996/97 season, Agricultural Statistic Document No. 9/1998

Oka, H.I. and W.T. Chang. 1961. Hybrid swarms between wild and cultivated rice species, *Oryza perennis* and *O. sativa*. *Evolution* 15:418-430.

Oka, H. 1988. *Origin of cultivated rice*. Japan Scientific Societies Press, Tokyo.

- Olsen, K. and A. Schaal. 2001. Microsatellite variation in cassava (*Manihot esculenta*, Euphorbiaceae) and its wild relatives: further evidence for a southern Amazonian origin of domestication. *Am J Bot* 88:131-142.
- Orlove, S.B. and B.S. Brush, 1996. Anthropology and the conservation of biodiversity. *Annu Rev Anthropol* 25:329-352.
- Ouborg, N.J., Piquot, Y. and J.M. van Groenendael. 1999. Population genetics, molecular markers and the study of dispersal in plants. *J Ecol* 87:551-568.
- Oupkaew P. 2009. Physiological and molecular variation in a gall midge resistant local rice variety, Muey Nawng. Master Thesis, Graduate School, Chiang Mai University.
- Palsson, S. 2004. Isolation by distance, based on microsatellite data, tested with spatial autocorrelation (SPAIDA) and assignment test (SPASSIGN). *Mol Ecol Notes* 4:143-145.
- Panaud, O., Chen, X. and S. McCouch. 1996. Development of microsatellite markers and characterization of simple sequence length polymorphism (SSLP) in rice (*Oryza sativa* L.). *Mol Gen Genet* 252:597-607.
- Papa, R and P. Gepts. 2004. Gene flow between crops and their wild progenitors. *Encyclopedia of Plant and Crop Science*. DOI:10.1081/E-EPCS 120017095
- Parzies, H.K., Spoor, W. and R.A. Ennos. 2004. Inferring seed exchange between farmers from population genetic structure of barley landrace Arabi Aswad from Northern Syria. *Genet Resour Crop Evol* 51:471-478.
- Peakall, R. and P.E. Smouse. 2006. GenA1Ex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Mol Ecol Notes* 6(1):288-295.

- Perales, R.H., Brush S.B. and C.O. Qualset 2003. Landraces of Maize in Central Mexico: an altitudinal transect. *Econ Bot* 57(1):7-20.
- Perales, R.H., Benz, F.B. and S. B. Brush. 2005. Maize diversity and ethnolinguistic diversity in Chiapas, Mexico. *Proc Natl Acad Sci USA* 102:949-954.
- Pingali, L.P. and S. Rajaram. 1998. Technological opportunities for sustaining wheat productivity growth toward 2020, 2020 vision brief 51 Washington D.C., USA.
- Pintasen S., Prom-u-thai C., Jamjod S., Yimyam N. and B. Rerkasem. 2007. Variation of grain iron content in a local upland rice germplasm from the village of Huai Tee Cha in northern Thailand. *Euphytica* DOI 10.1007/s10681-007-9421-7.
- Pressoir, G. and J. Berthaud. 2004. Population structure and strong divergent selection shape phenotypic diversification in maize landraces. *Heredity* 92:95-101.
- Pritchard, J. K. and W. Wen. 2004 Documentation for Structure software: version 2. See <http://pritch.bsd.uchicago.edu/software.html>
- Pritchard J.K., Stephens M., and P. Donnelly. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155: 945–959.
- Punyalue, A. 2006. Characterization of common wild rice populations from main rice growing region of Thailand. Master Thesis, Graduate School, Chiang Mai University
- Sanni, K.A., Fawole, I., Guei, R.G., Ojo, D.K., Somado, E.A., Ogunbayo, S.A. and I. Sanchez. 2007. Geographical patterns of phenotypic diversity in *Oryza sativa* landraces of Cote d'Ivoire. *Euphytica* 160:389-400.
- Sano Y. and H. Morishima. 1982. Variation in resource allocation and adaptive strategy of wild rice, *Oryza perennis* Moench. *Bot Gaz* 143(4):518-523.

- Sano Y., Morishima H. and H.I. Oka. 1980. Intermediate perennial-annual populations of *Oryza perennis* found in Thailand and their evolutionary significance. *Bot Mag Tokyo* 93:291-305.
- Savolainen, O., Pyhajarvi, T and T. Knurr. 2007. Gene flow and local adaptation in trees. *Ann Rev Ecol Evol Syst* 38:595-619.
- Schaal, B.A., Hayworth, A.D., Olsen, M.K., Rauscher, T.J. and A.W. Smith. 1998. Phylogeographic studies in plants: problems and prospects. *Mol Ecol* 7:465-474.
- Schmidt-Vogt, D. 2001. Secondary forests in swidden agriculture in the highland of Thailand. *J Trop For Sci* 13(4):748-767
- Second, G. 1982. Origin of the genic diversity of cultivated rice (*Oryza* spp.): study of the polymorphism scored at 40 isozyme loci. *Jpn J Genet* 57:25-57.
- Semon, M., Nielsen, R., Jones, M.P. and S.R. McCouch. 2005. The population structure of African cultivated rice *Oryza glaberrima* (Steud.): evidence for elevated levels of linkage disequilibrium caused by admixture with *O. sativa* and ecological adaptation. *Genetics* 169:1639-1647.
- Shishido R., Kikuchi M., Nomura K. and H. Ikehashi. 2006. Evaluation of genetic diversity of wild rice (*Oryza rufipogon* Griff.) in Myanmar using simple sequence repeats (SSRs). *Genet Resour Crop Evol* 53:179-186.
- Sirabanchongkran, A., Yimyam N., Boonma W., Rerkasem K., Coffey K., Pinedo-Vasquez, M. and C. Padoch. 2004. Varietal turnover and seed exchange: implications for conservation of rice genetic diversity on-farm. *IRRN* 29(2):12-14.
- Slatkin, M. 1993. Isolation by distance in equilibrium and non-equilibrium populations. *Evolution* 47:264-279.

- Smouse, P.E. and R. Peakall. 1999. Spatial autocorrelation analysis of individual multiallele and multilocus genetic structure. *Heredity* 82:561-573.
- Sokal, R.R. and D.E. Wartenberg. 1983. A test of spatial autocorrelation analysis using an isolation-by-distance model. *Genetics* 105:219-237.
- Soleri, D. and D.A. Cleveland. 1993. Farmer's genetic perceptions regarding their crop populations: an example with maize in the central valleys of Oaxaca, Mexico. *Econ Bot* 55:106-128.
- Song Z.P., Lu B-R., Zhu Y.G., J.K. Chen. 2003a. Gene flow from cultivated rice to the wild species *Oryza rufipogon* under experimental field conditions. *New Phytologist* 157: 67–665.
- Song, Z.P., Xu, X., Wang, B., Chen, J.K. and B.R. Lu. 2003b. Genetic diversity in the northernmost *Oryza rufipogon* populations estimated by SSR markers. *Theor Appl Genet* 107: 1492–1499.
- Song, Z., Li, B., Chen, J. and B.R. Lu. 2005. Genetic diversity and conservation of common wild rice (*Oryza rufipogon*) in China. *Plant Species Biol* 20:83-92.
- Song, Z., Lu, B.R. and J. Chen. 2004. Pollen flow of cultivated rice measured under experimental conditions. *Biodiversity Conserv* 13:579-590.
- Song, Z., Zhu, W., Rong, J., Xu, X., Chen, J. and B.R. Lu. 2006. Evidences of introgression from cultivated rice to *Oryza rufipogon* (Poaceae) populations based on SSR fingerprint: implications for wild rice differentiation and conservation. *Evol Ecol* 20:501-522.
- Supamongkol, P. 2006. Genetic diversity of local rice cv. *Meuy Nawng*. MSc (Agronomy), Graduate School, Chiang Mai University Thailand

- Sweeney, M. and S. McCouch. 2007. The complex history of the domestication of rice. *Ann Bot.* doi:10.1093/aob/mcm128
- Tamura K., Dudley J., Nei M. and S. Kumar. 2007. MEGA4: molecular evolution genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24(8):1596-1599.
- Tanksley, S.D. and S.R. McCouch. 1997. Seed banks and molecular maps: unlocking genetic potential from the wild. *Science* 277:1063-1066.
- Tarnita C.E., Ohtsuki H., Antal T., Fu F. and M.A. Nowak. 2009. Strategy selection in structured populations. *J Theor Biol* 259(3):570-581.
- Tiranti, B., and V. Negri. 2007. Selective microenvironmental effects play a role in shaping genetic diversity and structure in a *Phaseolus vulgaris* L. landrace: implications for on-farm conservation. *Mol Ecol* 16:4942-4955.
- Tosti, N, and V. Negri. 2005. On-going on-farm microevolutionary processes in neighbouring cowpea landraces revealed by molecular markers. *Theor Appl Genet* 110:1275-1283.
- Ullstrup, A.J. 1972. The impact of the southern corn leaf blight epidemics of 1970-1971. *Annu Rev Phytopathol* 10:37-50.
- Vaughan, D.A., Lu, B.R. and N. Tomooka. 2008. The evolving story of rice evolution. *Plant Sci* 174:394-408.
- Vaughan, D.A., Morishima, H., and K. Kadowaki. 2003. Diversity in the *Oryza* genus. *Curr Opin Plant Biol* 6:139-146.
- Vaughan, D.A., Sanchez, P.L., Ushiki, J., Kaga, A. and N. Tomooka. 2005. Chapter 16 Asian rice and weedy rice evolutionary perspectives. pp 257-277. In Gressel J.

- (Ed.) *Crop ferality and volunteerism*. CRC press, Taylor&Francis group, Boca Raton, FL, USA.
- Vaughan, L.K., Ottis, B.V., Prazak-Havey, A.M., Sneller, C., Chandler, J.M., and W.D. Park, 2001. Is all red rice found in commercial rice really *Oryza sativa*? *Weed Sci* 49:468-476.
- vom Brocke K., Weltzien E., Christinck A., Presterl T. and H.H. Geiger. 2003. Effects of farmers' seed management on performance and adaptation of pearl millet in Rajasthan, India. *Euphytica* 130:267-280.
- Ward, S. 2006. Genetic analysis of invasive plant populations at different spatial scales. *Biol Invasions* 8:541-552.
- Weir BS, and C.C. Cockerham. 1984. Estimating F-Statistics for the Analysis of Population Structure. *Evolution* 38:1358-1370.
- Wiley R.H. 1981. Social structure and individual ontogenies: problems of description, mechanism, and evolution. P. 105-133. In Bateson P. P. G. and P. H. Klopfer (Eds), *Perspicities in ethology*, vol. 4. Plenum Press, New York.
- Wongtamee, A. 2008. Evaluation of exotic genes contamination from crop rice (*Oryza sativa* L.) in common wild rice (*Oryza rufipogon* Griff.) populations. Master Thesis, Graduate School, Chiang Mai University
- Wood, D., and M.J. Lenne. 1997. The conservation of agrobiodiversity on-farm: questioning the emerging paradigm. *Biodiversity Conserv* 6:109-129.
- Wright, S. 1990. Evolution in Mendelian populations. *Bull Math Biol* 52:241-295.
- Wright, S. 1943. Isolation by distance. *Genetics* 28:114-138.

- Xiao, J., Li, J., Grandilloa, S., Ahn, S.N., Yuan, L., Tanksley, S.D. and S.R. McCouch. 1998. Identification of trait-improving quantitative trait loci alleles from a wild rice relative, *Oryza rufipogon*. *Genetics* 150:899-909.
- Yeh, F.C., Yang, R.C., and T.B.J. Boyle. 1999. POPGENE, version 1.32 Microsoft window-based freeware for population genetic analysis. (University of Alberta, Edmonton, Alberta Canada)
- Zeder, M.A., Emshwiller, E., Smith, B.D. and D.G. Bradley. 2006. Documenting domestication: the intersection of genetics and archaeology. *Trends Genet* 22(3):139-155.
- Zeven, C.A. 1999. The traditional inexplicable replacement of seed and seed ware of landraces and cultivars: a review. *Euphytica* 110:181-191.
- Zhou Q., Zheng X., Luo J., Gaut B.S. and S.Ge. 2007. Multilocus analysis of nucleotide variation of *Oryza sativa* and its wild relatives: severe bottleneck during domestication of rice. *Mol Biol Evol* 24(3):875-888.
- Zhou H.F., Zheng X.M., Wei R.X., Second G., Vaughan D.A. and S. Ge. 2008. Contrasting population genetic structure and gene flow between *Oryza rufipogon* and *Oryza sativa*. *Theor Appl Genet* 117:1181-1189.
- Zhou, H.F., Xie, Z.W. and S. Ge. 2003. Microsatellite analysis of genetic diversity and population genetic structure of a wild rice (*Oryza rufipogon* Griff.) in China. *Theor Appl Genet* 107:332-339.
- Zizumbo-Villarreal, D., Colunga-GarciaMarin, P., de la Cruz, E., Delgado-Valerio, P. and P. Gepts. 2005. Population structure and evolutionary dynamics of wild-weedy-domesticated complexes of common bean in a Mesoamerican region. *Crop Sci* 45:1073-1083.