Table of Contents

	Page
Acknowledgement	iii
Abstract in Thai	v
Abstract in English	vii
Table of Contents	ix
List of Tables	xiii
List of Figures	XV
List of Abbreviations	xvi
Chapter I Introduction	1
1.1 Background	1
1.2 Problem of rice production in Bangladesh	2
1.3 Rice pest status in Bangladesh	3
1.4 Current pest management practices	3
Chapter II Literature Review	5
2.1 Pest surveillance in rice production	5
2.2 Current situation of rice pest surveillance in Bangladesh	6
2.3 Three major rice insect pests of Bangladesh	7
2.4 Integrated Pest Management (IPM)	9
2.4.1 Current situation of rice IPM in Bangladesh	-12
2.4.2 Impact on IPM training	13
2.4.3 Integrated pest management (IPM) policy developments	14
2.4.4 Developments of community IPM	14
2.4.5 Farmer-farmer training	<u>15</u>
2.4.6 IPM in schools	15
2.4.7 IPM clubs/IPM congress	16
2.5 Geographic information systems (GIS) and IPM	16
2.6 Pest population mapping	18

Chapter III Research Methodology	21
3.1 Research procedures	22
3.2 Field and household survey	22
3.2.1Sampling techniques	25
3.2.2Data collection	25
3.2.2.1 Primary data collection	25
3.2.2.2 Secondary data collection	26
3.3 Data Analysis	26
3.3.1 GIS analysis	26
3.3.1.1 Software	27
3.3.1.2 Data acquisition	27
3.3.1.3 Design requirements	27
3.3.1.4 How to input data	28
3.3.2 Multiple regression model for factors influence benefit-cost ratio	29
3.3.3 Multinomial logistic model for adoption of surveillance method	29
Chapter IV Description of the Study Site	32
4.1 Description of Bangladesh socio-economic characteristics	32
4.1.1 Location and topology	32
4.1.2 Climate of Bangladesh	33
4.1.2.1 Rainfall	33
4.1.2.2 Temperature	34
4.1.3 Land use pattern in Bangladesh	35
4.1.4 Role of agriculture in Bangladesh economy	36
4.1.5 Rice farming in Bangladesh	37
4.1.6 Status of rice cultivation in different ecosystem	38
4.2 Description of Gazipur district	39
4.2.1 Socio-economic condition of Gazipur district	40
4.2.2 Problems of rice cultivation in Gazipur district	41
4.2.3 Pest infestation in Gazinur district	42

Chapter V Spatially-explicit Pest Infestation in Bangladesh	43
5.1 Spatial analysis three major insects of rice in 64 districts of Bangladesh	43
5.1.1 Stem borer infestation during the Year 1999-2008	44
5.1.2 Brown planthopper infestation during the year1999-2008	5(
5.1.3 Rice hispa infestation during the year 1999-2008	56
5.2 Conclusion	62
Chapter VI Comparison of Surveillance Method and Adoption of Surveillance Method	f 64
6.1 Types of farmers and sample size	64
6.1.1 Farming system practices by different types of farmers	67
6.1.2 Historical profile of major pest infestation record (7 years record)	67
6.1.3 Frequency of crop field visit by farmers in a week	68
6.1.4 Method of pest control	69
6.1.5 Analysis of pest-defender ratio in different farmer's field	70
6.1.6 Comparison of percentage of pest infestation in rice field	71
6.1.7 Comparison of labour force engaged in different types of rice farming	72
6.1.8 Comparison of benefit-cost ratio in different types of farmer's practice	73
6.2 Factors affecting benefit-cost ratio in different types of farmers	74
6.2.1 Benefit-cost ratio in different parameters in IPM farmer	74
6.2.2 Benefit cost ratio in different parameters in Non-IPM farmer	76
6.2.3 Benefit-cost ratio in different parameters in contact farmers' group	78
6.2.4 Benefit-cost ratio in different parameters for all farmers (n=125)	79
6.3 Multinomial logit model for farmers' surveillance method adoption	81
6.3.1 Factors affecting adoption of surveillance method	84

Chapter VII Summary of the Finding and Recommendations		86
7.1 Summary of the finding		86
7.1.1 Spatially-explicit find	ling	87
7.1.2 Socioeconomic sustai	nability comparison	87
7.1.3 Ecological sustainabi	lity comparison	88
7.1.4 Agronomic sustainab	ility comparison	89
7.1.5 Overall sustainability	comparison	89
7.2 Recommendation		89
7.2.1 Training and extension	on	90
7.3 Future research		90
7.4 Limitation of the study		91
References		92
Appendices		95
Appendix A		96
Appendix B		98
Appendix C		99
Curriculum Vitae		104

ลิขสิทธิมหาวิทยาลัยเชียงใหม Copyright[©] by Chiang Mai University All rights reserved

List of Tables

Table		Page
3.1	Comparison of monitoring system	23
4.1	Average temparature and rainfall	34
4.2	Land cover types and extent of Bangladesh 1992-1993	36
5.1	Spatial distribution of stem borer infestation during ten years period (district number and percentage)	47
5.2	Spatial distribution of BPH infestation during ten years period (district number and percentage)	53
5.3	Spatial distribution of rice hispa infestation during ten years period (district number and percentage)	59
6.1	Rice cultivated land extent in the sample	66
6.2	Farming system practices by different types of farmers	67
6.3	Historical profile of major pest infestation record (7 years record)	68
6.4	Frequency of crop field visit by farmers in a week	69
6.5	Different method taken for pest control	70
6.6	Pest-defender ratio in three types of farmer's field	71
6.7	Percentage of pest infestation in three different types of field	71
6.8	Average labour forces engaged in different types of farming	72
6.9	Average benefit-cost ratio in different types of farming	74
6.10 (a	a) Determinants of benefit-cost ratio in different parameters in IPM farmer	e 76
6.10 (ł	o) Determinants of benefit cost ratio in different parameters in Non-IPM farmer	77
6.10(c) Determinants of benefit cost ratio in different parameters in contact farmers' group	79

6.11 Determinants of benefit cost ratio in different parameters for all farmer (n=125)	81
6.12 Definition of variables and variables values in multinomial regression	82
6.13 Factors affecting adoption of surveillance method (likelihood ratio test)	85

List of Figures

Figure	Figure	
4.1	Map showing location of Gazipur district	40
5.1	Map showing stem borer infestation during the year 1999- 2002	48
5.2	Map showing stem borer infestation during the year 2003- 2006	49
5.3	Map showing stem borer infestation during the year 2007- 2008	50
5.4	Map showing brown planthopper infestation during the year 1999-2002	54
5.5	Map showing brown planthopper infestation during the year 2003-2006	55
5.6	Map showing brown planthopper infestation during the year 2007-2008	56
5.7	Map showing rice hispa infestation during the year 1999- 2002	60
5.8	Map showing rice hispa infestation during the year 2003- 2006	61
5.9	Map showing rice hispa infestation during the year 2007- 2008	62
6.1	Graph shows the land size frequency for different number of farmers	66

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

Abbreviations and Symbols

AESA = Agro Eco-system Analysis

AEZ = Agro Ecological Zone

BARI = Bangladesh Agricultural Research Institute

BRRI = Bangladesh Rice Research Institute

CRSP = Collaborative Research Support Program

DAE = Department of Agricultural Extension

DANIDA = Danish International Development Agency

FAO = Food and Agriculture Organization

FFS = Farmer Field School

FT = Farmer Trainer

GIS = Geographic Information System

GMO = Genetically Modified Organism

HYV = High Yielding Variety

ICP = Inter Country Program

IPM = Integrated Pest Management

NAP = National Agricultural Policy

SFS = Student Field School

SWTRC = Southwest Technical Resource Center

TOT = Training of Trainers

UNDP = United Nations Development Programme