

REFERENCES

- Ada GL. An Immunologist's View of HIV Infection. In : Merigan TC., Bartlett JG., Bolognesi D. Textbook of AIDS medicine, 2nd ed. Williams & Wilkins;1999:87-98.
- Albert J, Fenyo EM. Simple, sensitive, and specific detection of human immunodeficiency virus type 1 in clinical specimens by polymerase chain reaction with nested primers. *J Clin Microbiol.* 1990;28(7):1560-4.
- Barcellini W, Rizzardi GP, Borghi MO, Fain C, Lazzarin A, Meroni PL. TH1 and TH2 cytokine production by peripheral blood mononuclear cells from HIV-infected patients. *AIDS* 1994;8:757-762.
- Barcellini W, Rizzardi GP, Velati C, et al. In vitro production of type 1 and type 2 cytokines by peripheral blood mononuclear cells from high-risk HIV-negative intravenous drug users. *AIDS.* 1995;9:691-694.
- Bernard NF, Yannakis CM, Lee JS, Tsoukas CM. Human immunodeficiency virus (HIV)-specific cytotoxic T lymphocyte activity in HIV-exposed seronegative persons. *J Infect Dis* 1999;179:538-547.
- Beyrer C, Artenstein AW, Rugsao S, et al. Epidemiologic and biologic characterization of a cohort of human immunodeficiency virus type 1 highly exposed, persistently seronegative female sex workers in northern Thailand. *J Infect Dis* 1999;179:59-67.
- Biasin M, Caputo SL, Speciale L, et al. Mucosal and systemic immune activation is present in human immunodeficiency virus-exposed seronegative women. *J Infect Dis* 2000;182:1365-1374.
- Bongertz V., Costa CI., Veloso VG., et al. Vertical HIV-1 transmission: importance of neutralizing antibody titer and specificity. *Scand J Immunol.* 2001;53:302-309.
- Borkowsky W, Krasinski K, Moore T, Papeavangelou V. Lymphocyte proliferative responses to HIV-1 envelope and core antigens by infected and uninfected adults and children. *AIDS Res Hum Retroviruses* 1990; 6: 673-78.
- Broliden K, Hinkula J, Devito C, et al. Functional HIV-1 specific IgA antibodies in HIV-1 exposed, persistently IgG seronegative female sex workers. *Immunology Letters* 2001;79:29-36.
- Buchbinder SP, Katz MH, Hessel NA, O'Malley PM, Holmberg SD. Long-term HIV-1 infection without immunologic progression. *AIDS* 1994;8:1123-1128.
- Burke DS, McCutchan FE. Global distribution of Human Immunodeficiency Virus-1 clades. The Molecular Biology of Human Immunodeficiency Virus Type 1. In : Devita VT, Hallman S, Rosemberg SA, et al. editors. *AIDS : Etiology, diagnosis, treatment and prevention*, 4th ed. Philadelphia: Lippincott-Raven Publishers 1997. p.119-26.
- Burnett PR, VanCott TC, Polonis VR, Redfield RR, Birx DL. Serum IgA-mediated

- neutralization of HIV type 1. *J Immunol* 1994;152:4642-4648.
- Busch MP, Lee LL, Satten GA, et al. Time course of detection of viral and serologic markers preceding human immunodeficiency virus type 1 seroconversion: Implication for screening of blood and tissue donors. *Transfusion*. 1995;35:91-97.
- Canaris AD, Caruso A, Licenziati S, et al. Lack of polarized type 1 or type 2 cytokine profile in asymptomatic HIV-1-infected patients during a two-year bimonthly follow-up. *Scand J Immunol*. 1998;47:146-51.
- Cao Y., Qin L., Zhang L., Safrit J, Ho D. Virologic and immunologic characterization of Long-Term Survivors of Human Immunodeficiency Virus Type 1 Infection. *The New England Journal of Medicine*. 1995;332:201-208.
- Centers for Disease Control and Prevention. Persistent lack of detectable HIV-1 antibody in a person with HIV infection. Utah, 1995. *MMWR Morb Mortal Wkly Rep* 1996; 45:181-5.
- Cheynier R, Langlade-Demoyen P, Marescot MR, et al. Cytotoxic T lymphocyte responses in the peripheral blood of children born to human immunodeficiency virus-1-infected mothers. *Eur J Immunol* 1992; 22: 2211-7.
- Clerici M, Balotta C, Meroni L, et al. Type 1 cytokine production and low prevalence of viral isolation correlate with long-term nonprogression in HIV infection. *AIDS Res Hum Retroviruses*. 1996;12:1053-1061.
- Clerici M, Barassi C, Devito C, et al. Serum IgA of HIV-exposed uninfected individuals inhibit HIV through recognition of a region within the alpha-helix of gp41. *AIDS*. 2002;16:1731-1741.
- Clerici M, Berzofsky JA, Shearer GM, Tacket CO. Exposure to human immunodeficiency virus (HIV) type I indicated by HIV-specific T helper cell responses before detection of infection by polymerase chain reaction and serum antibodies. *J Infect Dis*. 1991;164:178-182.
- Clerici M, Giorgi JV, Chou C-C, et al. Cell-mediated immune response to human immunodeficiency virus (HIV) type 1 in seronegative homosexual men with recent sexual exposure to HIV-1. *J Infect Dis* 1992; 165: 1012-1019.
- Clerici M(a), Hakim FT, Venzon DJ, et al. Changes in interleukin-2 and interleukin-4 production in asymptomatic, human immunodeficiency virus-seropositive individuals. *J Clin Invest*. 1993;91:759-765.
- Clerici M(a), Levin JM, Kessler HA, et al. HIV-specific T-helper activity in seronegative health care workers exposed to contaminated blood. *JAMA*. 1994;5;271:42-46.
- Clerici M, Salvi A, Trabattoni D, et al. A role for mucosal immunity in resistance to HIV infection. *Immunol Lett*. 1999 ;66:21-25.
- Clerici M.(b), Shearer GM. A TH1-->TH2 switch is a critical step in the etiology of HIV infection. *Immunol Today*. 1993;14:107-111.
- Clerici M(b), Shearer GM. The Th1-Th2 hypothesis of HIV infection: new insights. *Immunol Today*. 1994;15:575-581.
- Clerici M(c), Sison AV, Berzofsky JA, et al. Cellular immune factors associated with mother-to-infant transmission of HIV. *AIDS*. 1993;7:1427-1433.
- Cocchi F, DeVico AL, Garzino-Demo A, et al. Identification of RANTES, MIP-1

- alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. *Science*. 1995;270:1811-1815.
- Cohn J. Exploiting the HIV-chemokine nexus. *Science* 1997;275:1261–1262.
- Cornelissen M, van den Burg R, Zorgdrager F, Lukashov V, Goudsmit J. *Pol* gene diversity of five human immunodeficiency virus type 1 subtypes: evidence for naturally occurring mutations that contribute to drug resistance, limited recombination patterns, and common ancestry for subtypes B and D. *J Virol*. 1997;70:6348-58.
- De Maria A, Cirillo C, Moretta L. Occurrence of human immunodeficiency virus type 1 (HIV-1) specific cytolytic T cell activity in apparently uninfected children born to HIV-1 infected mothers. *J Infect Dis* 1994;170: 1296-9.
- Dean M, Carrington M., Winkler C., et al. Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CKR-5 structural gene. *Science* 1996;273:1856–1862.
- Devito C, Broliden K, Kaul R, et al. Mucosal and plasma IgA from HIV-1-exposed uninfected individuals inhibit HIV-1 transcytosis across human epithelial cells. *J Immunol* 2000;165:5170-5176.
- Devito C, Hinkula J, Kaul R, et al. Mucosal and plasma IgA from HEPS individuals can neutralize HIV-1 infection. *AIDS* 2000;14:1917-20.
- Dorrell L, Hessel AJ, Wang M, et al. Absence of specific mucosal antibody responses in HIV-exposed uninfected sex workers from the Gambia. *AIDS* 2000;14:1117-1122.
- Douglas SD, Durako S, Sullivan KE, Camarca M, Moscicki AB, Wilson CM. TH1 and TH2 cytokine mRNA and protein levels in human immunodeficiency virus (HIV)-seropositive and HIV-seronegative youths. *Clin Diagn Lab Immunol*. 2003;10:399-404.
- Dyer WB, Ogg GS, Demoitie MA, et al. Strong human immunodeficiency virus (HIV)-specific cytotoxic T-lymphocyte activity in Sydney Blood Bank Cohort patients infected with nef-defective HIV type 1. *J. Virol*. 1999;73:436–443.
- Fakoya A, Matear PM, Filley E, et al. HIV infection alters the production of both type 1 and 2 cytokines but does not induce a polarized type 1 or 2 state. *AIDS*. 1997;11:1445-1452.
- Fan J, Bass HZ, Fahey JL. Elevated IFN-gamma and decreased IL-2 gene expression are associated with HIV infection. *J Immunol* 1993;151:5031-5040.
- Fowke KR, Nagelkerke NJD, Kimani J, et al. Resistance to HIV-1 infection among persistently seronegative prostitutes in Nairobi, Kenya. *Lancet* 1996, 348:1347-1351.
- Fujiwara T, Oda K, Yokota S, Takatsuki A, Ikehara Y, Brefeldin A causes disassembly of the Golgi complex and accumulation of secretory proteins in the endoplasmic reticulum. *J Biol Chem* 1988;263:18545-18552.
- Harrer T., Harrer E., Kalams S.A., et al. Cytotoxic T lymphocytes in asymptomatic long-term nonprogressing HIV-1 infection. Breadth and specificity of the response and relation to in vivo viral quasispecies in a person with prolonged infection and low viral load. *J. Immunol*. 1996;156:2616–2623.
- Hu YW, Birch P, Balaskas E, et al. Flow cytometric immunofluorescence assay for detection of antibodies to human immunodeficiency virus type 1 using


- insoluble precursor forms of recombinant polyproteins as carriers and antigens. *J Clin Microbiol* 1996;34:1412-1419.
- Jason J, Larned J. Single-cell cytokine profiles in normal humans: comparison of flow cytometric reagents and stimulation protocols. *J Immunol Methods*.1997;22;207:13-22.
- Jung T, Schauer U, Heusser C, Neumann C, Rieger C. Detection of intracellular cytokines by flow cytometry. *J Immunol Methods* 1993;26;159:197-207.
- Kanki P. Epidemiology and Natural History of Human Immunodeficiency Virus Type 2. The Molecular Biology of Human Immunodeficiency Virus Type 1. In : Devita VT, Hallman S, Rosenberg SA, et al. *AIDS : Etiology, Diagnosis, Treatment and Prevention*, 4th ed. Lippincott-Raven Publishers;1997:127-135.
- Kaul R(a), Dong T, Plummer FA, et al. CD8(+) lymphocytes respond to different HIV epitopes in seronegative and infected subjects. *J Clin Invest*. 2001;107:1303-1310.
- Kaul R(b), Plummer FA, Clerici M, Bomsel M, Lopalco L, Broliden K. Mucosal IgA in exposed, uninfected subjects: evidence for a role in protection against HIV infection. *AIDS* 2001;15:431-432.
- Kaul R, Plummer FA, Kimani J, et al. HIV-1-specific mucosal CD8+ lymphocyte responses in the cervix of HIV-1-resistant prostitutes in Nairobi. *J Immunol*. 2000;164:1602-1611.
- Kaul R, Trabattoni D, Bwayo JJ, et al. HIV-1-specific mucosal IgA in a cohort of HIV-1-resistant Kenyan sex workers. *AIDS* 1999;14;13:23-29.
- Kelso A. Th1 and Th2 subsets: paradigms lost? *Immunol Today* 1995;16:374-379.
- Klausner RD, Donaldson JG, Lippincott-Schwartz J, Brefeldin A: insights into the control of membrane traffic and organelle structure. *J Cell Biol* 1992;116:1071-1080.
- Klein SA, Dobmeyer JM, Dobmeyer TS, et al. Demonstration of the Th1 to Th2 cytokine shift during the course of HIV-1 infection using cytoplasmic cytokine detection on single cell level by flow cytometry. *AIDS* 1997;11:1111-1118.
- Kundu SK, Merigan TC. Heterogeneity of cytokine functions in HIV infection. *Immunology* 1997;91:234-238.
- Levy JA. The importance of the innate immune system in controlling HIV infection and disease. *Trends Immunol* 2001;22:312-316.
- Levy JA.(a). A Discovery and Nature of the HIV Virion. In : Levy JA. *HIV and the Pathogenesis of AIDS*, 1st ed. Washington, D.C. : ASM Press;1994:1-10.
- Levy JA.(b). Human and Cell Mediated Immune Responses To HIV Infection. In : Levy JA. *HIV and the Pathogenesis of AIDS*, 1st ed. Washington, D.C. : ASM Press;1994:153-180.
- Liles WC, van Voorhis C. Nomenclature and biologic significance of cytokines involved in inflammation and the host immune re-sponse. *J. Infect. Dis*. 1995;172:1573-1580.
- Lippincott-Schwartz J, Donaldson JG, Schweizer A, et al. Microtubule-dependent retrograde transport of proteins into the ER in the presence of brefeldin A suggests an ER recycling pathway. *Cell* 1990;9;60:821-836.
- Liu R, Paxton W, Choe S. et al. Homozygous defect in HIV 1 coreceptor accounts for

- resistance of some multiple exposed individuals to HIV 1 infection. *Cell* 1996;86 :1-11.
- Louisirirotchanakul S, Liu H, Roongpisuthipong A, et al. Genetic analysis of HIV-1 discordant couples in Thailand: association of CCR2 64I homozygosity with HIV-1-negative status. *J Acquir Immune Defic Syndr*. 2002;29:314-315.
- Lucey DR, Clerici M, Shearer GM. Type 1 and type 2 cytokine dysregulation in human infectious, neoplastic, and inflammatory diseases. *Clin Microbiol Rev*. 1996;9:532-562.
- Maino VC, Picker LJ. Identification of functional subsets by flow cytometry: intracellular detection of cytokine expression. *Cytometry* 1998;34:207-215.
- Mazzoli S, Lopalco L, Salvi A, et al. Human immunodeficiency virus (HIV)-specific IgA and HIV neutralizing activity in the serum of exposed seronegative partners of HIV-seropositive persons. *J Infect Dis* 1999;180:871-875.
- Mazzoli S, Trabattoni D, Lo Caputo S, et al. HIV-specific mucosal and cellular immunity in HIV-seronegative partners of HIV-seropositive individuals. *Nature Med* 1997;3:1250-1257.
- McCutchan FE, Hegerich PA, Brennan TP, et al. Genetic variants of HIV-1 in Thailand. *AIDS Res Hum Retroviruses*. 1992;8:1887-1895.
- McMichael AJ., Rowland-Jones SL. Cellular immune responses to HIV. *Nature* 2001; 410, 980-987.
- Meroni L, Trabattoni D, Balotta C, et al. Evidence for type 2 cytokine production and lymphocyte activation in the early phases of HIV-1 infection. *AIDS* 1996;10:23-30.
- Meyaard L, Hovenkamp E, Keet IP, et al. Single cell analysis of IL-4 and IFN-gamma production by T cells from HIV-infected individuals: decreased IFN-gamma in the presence of preserved IL-4 production. *J Immunol* 1996;157:2712-2718.
- Mosmann TR, Li L, Hengartner H, Kagi D, Fu W, Sad S. Differentiation and functions of T cell subsets. *Ciba Found Symp*. 1997;204:148-54; discussion 154-158.
- Nicastri E, Sarmati L, Ercoli L, et al. Reduction of IFN-gamma and IL-2 production by peripheral lymphocytes of HIV-exposed seronegative subjects. *AIDS* 1999;30;13:1333-1336.
- Nishanian P, Huskins KR, Stehn S, Detels R, Fahey JL. A simple method for improved assay demonstrates that HIV p24 antigen is present as immune complexes in most sera from HIV-infected individuals. *J Infect Dis*. 1990;162:21-28.
- Novitsky V, Rybak N, McLane MF, et al. Identification of human immunodeficiency virus type 1 subtype C Gag-, Tat-, Rev-, and Nef-specific elispot-based cytotoxic T-lymphocyte responses for AIDS vaccine design. *J Virol*. 2001;75:9210-9228.
- Oppenheim JJ., Ruscetti FW.. Cytokines. In: Tristram G, Parslow. Editor. *Medical Immunology*., 10th ed. New York: McGraw-Hill Companies, Inc. 2001; 148-166.
- Pala P, Hussell T, Openshaw PJ. Flow cytometric measurement of intracellular cytokines. *J Immunol Methods* 2000; 21;243:107-124.
- Pantaleo G(a), Fauci AS. New concepts in the immunopathogenesis of HIV infection.

- Annu Rev Immunol. 1995;13:487-512.
- Pantaleo G, Graziosi C, Fauci AS. New concepts in the immunopathogenesis of human immunodeficiency virus infection. *N Engl J Med*. 1993;328:327-35.
- Pantaleo G(b), Menzo S, Vaccarezza M, et al. Studies in subjects with long-term nonprogressive human immunodeficiency virus infection. *N Engl J Med* 1995;26;332:209-216.
- Pavlikis GN. The Molecular Biology of Human Immunodeficiency Virus type 1. In : Devita VT, Hallman S, Rosemberg SA, et al. editors. *AIDS : Etiology, diagnosis, treatment and prevention*, 4th ed. Philadelphia: Lippincott-Raven Publishers 1997. p.45-74.
- Picker LJ, Singh MK, Zdraveski Z, et al. Direct demonstration of cytokine synthesis heterogeneity among human memory/effector T cells by flow cytometry. *Blood* 1995;15;86:1408-1419.
- Pinto LA, Sullivan J, Berzofsky JA, et al. ENV-specific cytotoxic T lymphocyte responses in HIV seronegative health care workers occupationally exposed to HIV-contaminated body fluids. *J Clin Invest* 1995; 96:867–876.
- Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods* 1995;15;188:117-128.
- Psallidopoulos MC, Schnittman SM, Thompson LM 3rd, et al. Integrated proviral human immunodeficiency virus type 1 is present in CD4+ peripheral blood lymphocytes in healthy seropositive individuals. *Viol.* 1989;63:4626-4631.
- Reimer L, Mottice S, Schable C, et al. Absence of detectable antibody in a patient infected with human immunodeficiency virus. *Clin Infect Dis*. 1997;25:98-100.
- Romagnani S. Biology of human TH1 and TH2 cells. *J Clin Immunol*. 1995;15:121-9.
- Romagnani S. T-cell subsets (Th1 versus Th2). *Ann Allergy Asthma Immunol* 2000;85:9-18.
- Romagnani S, Maggi E, Del Prete G. HIV can induce a TH1 to TH0 shift, and preferentially replicates in CD4+ T-cell clones producing TH2-type cytokines. *Res Immunol*. 1994;145:611-617; discussion 617-618.
- Rosenberg ES, Billingsley JM, Caliendo AM, et al. Vigorous HIV-1-specific CD41 T-cell responses associated with control of viremia. *Science* 1997;278:1447–1450.
- Rowland-Jones SL, McMichael A. Immune responses in HIV-exposed seronegatives: have they repelled the virus? *Curr Opin Immunol* 1995;7:448-455.
- Rowland-Jones SL, Sutton J, Ariyoshi K, et al. HIV-specific cytotoxic T-cells in HIV-exposed but uninfected Gambian women. *Nat. Med*. 1995;1:59–64.
- Rugeles MT, Solano F, Diaz FJ, Bedoya VI, Patin PJ. Molecular characterization of the *CCR 5* gene in seronegative individuals exposed to human immunodeficiency virus (HIV). *J Clin Virology*. 2002;23:161–169.
- Salk J, Bretscher PA, Salk PL, Clerici M, Shearer GM. A strategy for prophylactic vaccination against HIV. *Science* 1993; 260: 1270–1272.
- Sester M, Sester U, Kohler H, et al. Rapid whole blood analysis of virus-specific CD4 and CD8 T cell responses in persistent HIV infection. *AIDS* 2000;14:2653-2660.

- Shearer GM. AIDS 5:245-253.1991, Clerici M., 1993 J Clin Invest 91:759-765, Clerici M, Shearer GM. A TH1-->TH2 switch is a critical step in the etiology of HIV infection. Immunol Today 1993; 14: 107-111.
- Shearer GM, Clerici M. Early T-helper cell defects in HIV infection. AIDS. 1991;5:245-253.
- Shearer GM, Clerici M. Protective immunity against HIV infection: has nature done the experiment for us? Immunol Today. 1996;17:21-24.
- Sligh JM, Roodman ST, Tsai CC. Flow cytometric indirect immunofluorescence assay with high sensitivity and specificity for detection of antibodies to human immunodeficiency virus (HIV). Am J Clin Pathol 1989;91:210-214.
- Sousa AE, Chaves AF, Loureiro A, Victorino RM. Comparison of the frequency of interleukin (IL)-2-, interferon-gamma-, and IL-4-producing T cells in 2 diseases, human immunodeficiency virus types 1 and 2, with distinct clinical outcomes. J Infect Dis. 2001;184:552-559.
- Spits H, Yssel H, Paliard X, Kastelein R, Figdor C, de Vries JE. IL-4 inhibits IL-2-mediated induction of human lymphokine-activated killer cells, but not the generation of antigen-specific cytotoxic T lymphocytes in mixed leukocyte cultures. J Immunol 1988;1;141:29-36.
- Sriwanthana B, Hodge T, Mastro TD, et al. HIV-specific cytotoxic T lymphocytes, HLA-A11, and chemokine-related factors may act synergistically to determine HIV resistance in CCR5 delta32-negative female sex workers in Chiang Rai, northern Thailand. AIDS Res Hum Retroviruses 2001;17:719-734.
- Stoneburner RL. Joint United Nations Programme on HIV/AIDS. The HIV/AIDS situation in mid :global and regional highlights. UNAIDS Fact Sheet. AIDS 1996;4:99-106.
- Stranford SA., Skurnick J, Louria D, et al. Lack of infection in HIV-exposed individuals is associated with a strong CD8(+) cell noncytotoxic anti-HIV response. Proc Natl Acad Sci U S A. 1999;96:1030-1035.
- Sullivan PS, Schable C, Koch W, et al. Persistently negative HIV-1 antibody enzyme immunoassay screening results for patients with HIV-1 infection and AIDS: serologic, clinical, and virologic results. Seronegative AIDS Clinical Study Group. AIDS 1999;14;13:89-96.
- Sutthent R, Foongladda S, Likanonskul S, et al. Detection of HIV-1 proviral DNA by polymerase chain reaction: a preliminary study in Bangkok. J Med Assoc Thai 1996;79:142-148.
- UNAIDS (2002) Fact Sheet Sub-Saharan Africa. Available <http://www.unaids.org/> (2003, December 22).
- Vilcek J, Le J. Immunology of cytokines: an introduction. In : Thomson AW. editor. The Cytokine Handbook. , 2nd ed. London: Academic Press Limited 1994; p.1-19.
- von Sydow M, Gaines H, Sonnerborg A, et al. Antigen detection in primary infection. Br Med J1988;296:238 -240.
- Weiblen BJ, Schumacher RT, Hoff R. Detection of IgM and IgA HIV antibodies after removal of IgG with recombinant protein G. J Immunol Methods 1990;9;126:199-204.
- Weniger BG, Takebe Y, Ou CY, Yamazaki S. The molecular epidemiology of HIV in

- Asia. AIDS. 1994;8 Suppl 2:S13-28.
- Whitmire JK, Asano MS, Murali-Krishna K, Suresh M, Ahmed R. Long-term CD4 Th1 and Th2 memory following acute lymphocytic choriomeningitis virus infection. J Virol 1998; 72:8281–8288.
- Williams SB, Flanigan TP, Cu-Uvin S, et al. Human immunodeficiency virus (HIV)-specific antibody in cervicovaginal lavage specimens obtained from women infected with HIV type 1. Clin Infect Dis. 2002;35:611-7.
- Zajac AJ, Blattman JN, Murali-Krishna K, et al. Viral immune evasion due to persistence of activated T cells without effector function. J Exp Med 1998;188:2205–2213.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright © by Chiang Mai University

All rights reserved