

REFERENCES

1. Centers for Disease Control and Prevention (CDC). National Diabetes Fact Sheet United States, 2003 [Online]. Available: http://www.cdc.gov/diabetes/pubs/pdf/ndfs_2003.pdf (4 March 2005).
2. Aekplakorn W, Stolk RP, Neal B, Suriyawongpaisal P, Chongsuvivatwong V, Cheepudomwit S, Woodward M. The prevalence and management of diabetes in Thai adults: The International Collaborative Study of Cardiovascular Disease in Asia. *Diabetes Care* 2003; 26(10):2758-2763.
3. Nesto RW. The relation of insulin resistance syndromes to risk of cardiovascular disease. *Rev Cardiovas Med* 2003; 4(6):S11-S18.
4. Grant PJ, Davies JA. Cardiovascular disease and diabetes. In: Pickup JC, Williams G. editor. *Textbook of diabetes* 2. 3 ed. Oxford: Blackwell Publishing; 2003. p. 56.1-56.24.
5. Williams G, Pickup, JC. *Handbook of Diabetes*. 2 ed. Oxford: Blackwell Science; 1999.
6. Taegtmeyer H, McNulty P, Young ME. Adaptation and maladaptation of the heart in diabetes: Part I general concepts. *Circulation* 2002; 105:1727-1733.
7. Davis WA, Knuiman M, Kendall PK, Grange V, Davis TME. Glycemic exposure is associated with reduced pulmonary function in type 2 diabetes. *Diabetes Care* 2004; 27(3):752-757.

8. Benbassat CA, Stern E, Kramer M, Lebzelter J, Blum I, Fink G. Pulmonary function in patients with diabetes mellitus (abstract). Am J Med Sci 2001; 332(3):127-132.
9. Melo E, Vianna EO, Gallo L, Foss NC, Terra-Filho J. Pulmonary function, cholinergic bronchomotor tone, and cardiac autonomic abnormalities in type 2 diabetes patients. Braz J Med and Biol Res 2003; 36:291-299.
10. Hedman A, Berglund L, Gustavsson BE, Reneland R, Lithell H. Relationships between muscle morphology and insulin sensitivity are improved after adjustment for intra-individual variability in 70-year-old men. Acta Physiol Scand 2000; 169:125-132.
11. Lillioja S, Young AA, Culter CL, Ivy JL, Abbott WGH., Zawadzki, JK, Yki-Jarvinen H, Chris L, Secomb TW, Bogardus C. Skeletal muscle capillary density and fiber type are possible determinants of in vivo insulin resistance in man. J Clin Invest 1987; 80:415-424.
12. Andersen H, Nielsen S, Mogensen CE, Jakobsen J. Muscle strength in type 2 Diabetes. Diabetes 2004; 53:1543-1548.
13. Estacio RO, Jeffers B, Regensteiner JG, Dickenson M, Wolfel EE, Schrier RW. The association between diabetic complications and exercise capacity in NIDDM patients. Diabetes Care 1998; 21(2):291-295.
14. Gregg EW, Beckles GLA, Williamson DF, Leveille SG, Langlois JA, Engelgau MM, Narayan KMV. Diabetes and physical disability among older U.S. Adults. Diabetes Care 2000; 23:1272-1277.

15. Gregg EW, Mangione CM, Cauley JA, Thompson TJ, Schwartz AV, Ensrud KE, Nevitt MC. Diabetes and incidence of functional disability in older women. *Diabetes Care* 2002; 25:61-67.
16. Volpato S, Blaum C, Resnick H, Ferrucci L, Fried LP, Guralnik JM. Comorbidities and impairments explaining the association between diabetes and lower extremity disability. *Diabetes Care* 2002; 25:678-683.
17. Volpato S, Blaum C, Ferrucci L, Ostir G, Cappola A, Fried LP, Guralnik JM. Progression of lower-extremity disability in older women with diabetes. *Diabetes Care* 2002; 26:70-75.
18. Wei M, Gibbons LW, Kampert JB, Nichaman MZ, Blair SN. Low cardiorespiratory fitness and physical Inactivity as predictors of mortality in men with type 2 diabetes. *Ann Intern Med* 2000; 132(8):605-611.
19. Ozdirenc M, Biberoglu S, Ozcan A. Evaluation of physical fitness in patients with type 2 diabetes mellitus. *Diabetes Res Clin Pract* 2003; 60:171-176.
20. American college of sports. ACSM's guidelines for exercise testing and prescription. 7th ed. Philadelphia: Lippincott Williams & Wilkins; 2005.
21. Power SK, Howley ET. Exercise Physiology: Theory and application to fitness and Performance. 5 ed. New York: McGraw- Hill; pp 145, 191, 2004.
22. American Thoracic Society (ATS). ATS statement: Guidelines for the six-minute walk test. *Am J Respir Crit Care Med* 2002; 166:111-117

23. Solway S, Brooks D, Lacasse Y, Thomas SA. Qualitative systematic overview of the measurement properties of functional walk tests used in the cardiorespiratory domain. *Chest* 2001; 119:256-270.
24. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004; 27(5):1047-1053.
25. World Health Organization (WHO). Laboratory diagnosis and monitoring of diabetes mellitus. Geneva, 2002 [Online].
Available: <http://whqlibdoc.who.int/hq/2002/9241590483.pdf> (28 January 2005).
26. Bennett SE, Karnes JL. Neurological disabilities. Philadelphia: Lippincott; pp 3-4, 1998.
27. World Health Organization (WHO). Definition, diagnosis and classification of diabetes mellitus and its complication; Part 1: Diagnosis and Classification of Diabetes Mellitus. Geneva, 1999 [Online]. Available:
http://www.staff.ncl.ac.uk/philip.home/who_dmg.pdf. (28 January 2005)
28. American Diabetes Association (ADA). Physical activity/ Exercise and Diabetes. *Diabetes Care* 2004; 27(supplement 1):558-562.
29. The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 2003; 26(1):S1-S20.
30. Hawley JA, Houmard JA. Introduction-preventing insulin resistance through exercise: A cellular approach. *Med Sci Sports Exerc* 2004; 36(7):1187-1190.

31. Reusch JEB. Current concepts in insulin resistance, type 2 Diabetes Mellitus and the Metabolic Syndrome. *Am J Cardiol* 2002; 90(5A):19G-26G.
32. Spranger J, Pfeiffer AFH. Diabetic microvascular complications. In: Mogensen CE, editor. *Hypertension and diabetes*. London: Lippincott Williams & Wilkins; 2002. p. 57-67.
33. Nathan DM. Long term complications of diabetes mellitus. *N Engl J Med* 1993; 328(23):1676-1685.
34. American Thoracic Society (ATS). Standardization of spirometry 1994 update. *Am J Respir Crit Care Med* 1995; 152:1107-1136.
35. Hughes JMB, Pride NB. Lung function tests: physiological principles and clinical applications. London: W. B. Saunders; 1999.
36. Yki-Jarvinen H. Insulin resistance in type 2 diabetes. In: Pickup JC, Williams, G., editor. *Textbook of diabetes 1*. Oxford: Blackwell Publishing; 2003. p. 22.1-22.19.
37. Nyholm B, Qu Z, Kaal S, Pedersen SB, Gravholt CH, Andersen JL, Saltin B, Schmitz O. Evidence of an increased number of type IIb muscle fibers in insulin-resistant first-degree relatives of patients with NIDDM. *Diabetes* 1997; 46:1822-1828.
38. Marin P, Andersson B, Krotkiewski M, Björntorp P. Muscle fiber composition and capillary density in women and men with NIDDM. *Diabetes Care* 1994;17(5):382-386.

39. Holmback AM, Porter MM, Downham D, LExell J,. Reliability of isokinetic ankle dorsiflexion strength measurement in healthy young men and women. Scand J Rehab Med 1999; 31:229-239.
40. Kramer JF. Reliability of knee extensor and flexor torques during continuous concentric-eccentric cycles. Arch Phy Med Rehabil 1990; 7:460-464.
41. Wennerberg D. Reliability of an isokinetic dorsiflexion and plantar flexion apparatus. Am J Sports Med 1991; 19(5):519-522.
42. Frontera WR, Hughes VA, Lutz KJ, Evans WJ. A cross sectional study of muscle strength and mass in 45 to 78 yr old men and women. J Appl Physiol 1991; 71(2):644-650.
43. Frontera WR, Hughes VA, Fielding RA, Fiatarone MA, Evans WJ, Roubenoff, R. Aging of skeletal muscle: a 12 yr longitudinal study. J Appl Physiol 2000; 88:1321-1326.
44. Wei M, Gibbons LW, Mitchell TL, Kampert JB, Lee CD, Blair SN. The association between cardiorespiratory fitness and impaired fasting glucose and type 2 diabetes mellitus in men. Ann Intern Med 1999; 130(2):89-96.
45. McGavock J, Mandic S, Lewanczuk R, Koller M, Muhll IV, Quinney A, Taylor D, Welsh R, Haykowsky M. Cardiovascular adaptations to exercise training in postmenopausal women with type 2 diabetes mellitus. Cardiovasc Diabetol 2004; 3 (3).

46. Maiorana A, O' Driscoll G, Goodman C, Taylor R, Green D. Combined aerobic and resistance exercise improve glycemic control and fitness in type 2 diabetes. *Diabetes Res Clin Pract* 2002; 56:115-123.
47. Boule NG, Haddad E, Kenny GP, Wells GA, Sigal RJ. Effects of exercise on glycemic control and body mass in type 2 Diabetes mellitus. *JAMA* 2001; 286(10):1218-1227.
48. Peirce NS. Diabetes and exercise. *Br J Sports Med* 1999; 33:161-173.
49. Tokmakidis SP, Zois CE, Volaklis KA, Kotsa K, Touvra A-M. The effects of a combined strength and aerobic exercise program on glucose control and insulin action in women with type 2 diabetes. *Eur J Appl Physiol* 2004; 92:437-442.
50. Bautmans I, Lambert M, Mets T. The six-minute walk test in community dwelling elderly: influence of health status. *BMC Geriatrics* 2004; 4 (6).
51. Troosters T, Gosselink R, Decramer M. Six minute walking distance in healthy elderly subjects. *Eur Respir J* 1999; 14:270-274.
52. Carter R, Holiday DB, Stocks J, Grothues C, Tiep B. Predicting oxygen uptake for men and women with moderate to severe chronic obstructive pulmonary disease. *Arch Phy Med Rehabil* 2003; 84:1158-1164.
53. Cahalin LP, Mathier MA, Semigran MJ, Dec GW, DiSalvo TG. The six minute walk test predicts peak oxygen uptake and survival in patients with advanced heart failure. *Chest* 1996; 110:325-332.

54. Demers C, McKelvie RS, Negassa A, Yusuf S. Reliability, validity, and responsiveness of the six-minute walk test in patients with heart failure. *Am Heart J* 2001; 142:698-703.
55. Pankoff BA, Overend TJ, Lucy SD, White KP. Reliability of the six-minute walk test in people with fibromyalgia. *Arthritis Care Res* 2000; 13(5):291-295.
56. Kervio G, Ville NS, Leclercq C, Daubert J-C, Carre F. Intensity and daily reliability of the six-minute walk test in moderate chronic heart failure patients. *Arch Phy Med Rehabil* 2004; 85:1513-1518.
57. Eaton T, Young P, Milne D, Wells AU. Six-minute walk, maximal exercise tests: reproducibility in fibrotic interstitial pneumonia. *Am J Respir Crit Care Med* 2005; 171:1150-1157.
58. Enright PL, Sherrill DL. Reference equations for the six-minute walk in healthy adults. *Am J Respir Crit Care Med* 1998; 158:1384-1387.
59. Stevens D, Elpern E, Sharma K, Szidon P, Ankin M, Kesten S. Comparison of hallway and treadmill six-minute walk tests. *Am J Respir Crit Care Med* 1999; 160:1540-1543.
60. Brooks D, Solway S, Weinacht K, Wang D, Thomas S. Comparison between an indoor and an outdoor 6-minute walk test among individuals with chronic obstructive pulmonary disease. *Arch Phy Med Rehabil* 2003; 84:873-876.
61. Wu G, Sanderson B, Bittner V. The 6-minute walk test: How important is the learning effect? *Am Heart J* 2003; 146:129-133.

62. Lord SR, Menz HB. Physiologic, psychologic and health predictors of 6-minute walk performance in older people. *Arch Phy Med Rehabil* 2002; 83:907-911.
63. กองออกกำลังกายเพื่อสุขภาพ กรมอนามัย กระทรวงสาธารณสุขคู่มือ. 2547. "แบบสัมภาษณ์การเคลื่อนไหวออกแรง/ออกกำลังกาย." [ระบบออนไลน์]. แหล่งที่มา <http://www.anamai.moph.go.th> (27 มกราคม 2548).
64. Scanlon PD, Connett JE, Waller LA, Altose MD, Bailey WC, Buist AS, Tashkin DP. Smoking cessation and lung function in mild-to-moderate chronic obstructive pulmonary disease: the lung health study. *AM J Respir Crit Care Med* 2000; 161:381-390.
65. Pride NB. Smoking cessation: effects on symptoms, spirometry and future trends in COPD. *Thorax* 2006; 56(Suppl II):ii7-ii10.
66. Weisell RC. Body mass index as an indicator of obesity. *Asia Pacific J Clin Nutr* 2002; 11(suppl):s681-s684.
67. Poh H, Eastwood PR, Cecins NM, Ho KT, Jenkins SC. Six-minute walk distance in healthy Singaporean adults cannot be predicted using reference equations derived from Caucasian populations. *Respirology* 2006;11(2):211-216.
68. Haffner SJ, Cassells H. Hyperglycemia as a cardiovascular risk factor. *The American Journal of Medicine* 2003;115(8A):6s-11s.
69. Jesudason DR, Leong D, Dunstan K, Wittert GA. Macrovascular risk and diagnostic criteria for type 2 diabetes. *Diabetes Care* 2003; 26:485-490.

70. Isokinetic muscle strength test. (No date). Exercising and testing [Online]. Available: <http://www.isokinetics.net/joints/kneeadvanced.htm>. (31 May 2005)
71. Aagaard P, Simonsen, EB, Magnusson SP., Larssonand B, Dyhre-Poulsen P. A new concept for isokinetic hamstring:quadriceps muscle strength ratio. Am J Sports Med 1998; 26:231-237.
72. Calmels PM, Nellen M, van der Borne I, Jourdin P, Minaire P. Concentric and eccentric isokinetic assessment of flexor-extensor torque ratios at the hip, knee, and ankle in a sample population of healthy subjects. Arch Phys Med Rehabil 1997; 78:1224-1230.
73. Kannus P. Ratio of hamstring to quadriceps femoris muscles' strength in the anterior cruciate ligament. Phys Ther 1988; 68:961-965.
74. Camarri B, Eastwood PR, Cecins NM, Thompson PJ, Jenkins S. Six minute walk distance in healthy subjects aged 55-75 years. Respir Med 2006; 100(4):658-665.