

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENT	iii
ABSTRACT ENGLISH	iv
ABSTRACT THAI	viii
TABLE OF CONTENTS	xi
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
ABBREVIATIONS	xx
 CHAPTER I: INTRODUCTION	
A. INTRODUCTION	
I. Bilirubin Metabolism	1
II. Function of Bilirubin in Serum	3
III. Hyperbilirubinemia and Clinical Significance of Bilirubin in Human Serum	3
IV. Neonatal Hyperbilirubinemia	5
V. Toxic effects of bilirubin on cellular functions and fragmentation of DNA in brain tissue	8
 B. LITERATURE REVIEW	
I. Neonatal Jaundice and Bilirubin Encephalopathy	10
II. Unconjugated Bilirubin Toxicity in Neural Cells	11

	Page
III. Association of Albumin and Bilirubin in Serum Effect to Cytotoxicity	11
IV. Mechanism of Unconjugated Bilirubin Toxicity in Neural Cells	13
V. Mechanism of Unconjugated Bilirubin Toxicity in Neural Cells Neonates	14
VI. Effect of Concentrations and Times of Unconjugated Bilirubin Expose to the cells	15
VII. Role of Bilirubin Induced Cells Apoptosis	15
VIII. Evidence Involved in Initiation of DNA Degradation	16
IX. Alternative Study on DNA Fragmentation <i>in vitro</i>	20
C. OBJECTIVES	22
 CHAPTER II: MATERIALS AND METHODS	
A. MATERIALS	
I. Instruments	23
II. Chemicals and Reagents	24
B. METHODS	
I. Spectrophotometric Determination of Bilirubin Absorption Spectra	25
II. Investigation of the Absorption Spectra of Bilirubin in the Presence of Effectors by Spectrophotometric Method	25
1. The effect of transition metal ions on bilirubin absorption spectra	25
1.1 Types of transition metal ions	25
1.2 The effect of transition metal ion concentrations on the absorption spectra of bilirubin	26
2. The effect of albumin on the interaction of transition metal ion with bilirubin observed by Spectrophotometric method	26

	Page
3. Confirmation of bilirubin binding with interaction metal ion	27
III. Method for Investigation of the Effect of Bilirubin Interacted with Transition Metal Ions on DNA Degradation <i>in vitro</i>	28
1. Investigation of the effect of bilirubin with or without albumin and transition metal ions concentrations on DNA degradation <i>in vitro</i>	28
1.1 The effect of bilirubin concentrations on the degradation of DNA with fixed transition metal ion concentration.	28
1.2 The effect of transition metal ion concentrations on DNA degradation with fixed bilirubin concentrations	28
2. Time dependent kinetic of DNA degradation	29
3. Protocol of agarose gel electrophoresis	29
4. Quantitation of DNA degradation	30
4.1 S ₁ Nuclease digestion of DNA	30
4.2 The colorimetric diphenylamine reaction	31
4.2.1 DNA standard graph	31
4.2.2 Percentage of DNA hydrolysis	32
4.2.3 Determination of 100% DNA hydrolyzed	32
IV. Investigation of the Mechanism of Bilirubin Interacted with Metal Ion Influenced DNA Degradation <i>in vitro</i>	32
1. Standard graph of malondialdehyde (MDA)	33
2. Free radical generation resulted from the interaction of bilirubin with metal ions	33
3. Investigation of free radical generated by bilirubin degradation <i>in vitro</i> by the MDA assay	34

CHAPTER III: RESULTS

I.	Spectrophotometric Determination of Bilirubin Absorption Spectra	36
II.	Investigation of the Absorption Spectra of Bilirubin in the Presence of Effectors by Spectrophotometric Method	37
	1. The effect of transition metal ions on bilirubin absorption spectra	37
	1.1 Types of transition metal ions	37
	1.2 The effect of transition metal ion concentrations on the absorption spectra of bilirubin	39
	2. Confirmation of bilirubin binding with interaction metal ion	42
	3. The effect of albumin on bilirubin-transition metal ion complex on the absorption spectra of bilirubin	43
III.	Investigation of the Effect of Bilirubin interacted with Transition metal ions on DNA Degradation <i>in vitro</i>	48
	1. Investigation of the effect of bilirubin and transition metal ions concentrations on DNA degradation <i>in vitro</i>	48
	1.1 Quantitation of DNA degradation	48
	1.2 The effect of bilirubin concentrations on the degradation of DNA <i>in vitro</i> with fixed transition metal ion concentration	49
	1.3 The effect of transition metal ions concentrations on DNA degradation <i>in vitro</i> at fixed 500 μ M bilirubin concentrations	52
	2. Time dependent kinetic of DNA degradation	56
IV.	Investigation of the Mechanism of Bilirubin Interacted with Metal Ion Influenced DNA Degradation <i>in vitro</i>	59
	1. Standard graph of malondialdehyde (MDA)	59
	2. Free radical generation resulted by the interaction of bilirubin with metal ions	61
	2.1 The inhibition effect of free radical scavengers on bilirubin-CuCl ₂ complex	61

	Page
2.2 The inhibition effect of free radical scavengers on bilirubin – FeCl ₂ complex	64
2.3 The inhibition effect of free radical scavengers on bilirubin – ZnCl ₂ complex	67
3. Investigation of free radical generation by the MDA assay by spectrophotometric scanning	73
CHAPTER IV: DISCUSSION AND CONCLUSION	
A. DISCUSSION	76
B. CONCLUSION	85
REFERENCES	86
APPENDICE	96
I. Appendix A. Reagent Preparation	97
II. Appendix B. Confirmation of Cu(I) Production by Bathocuproine	102
III. Appendix C. Absorption spectra of bilirubin-Cu(II) complex with And without scavenger (thiourea)	103
CURRICULUM VITAE	104

LIST OF TABLES

Table	Page
1. Inhibition of calf thymus DNA degradation after treatment with bilirubin and transition metal ions (200 μ M each) in the presence of free radical scavengers (50 mM in all case).	72

LIST OF FIGURES

Figure	Page
1 Enzymatic mechanism of bilirubin formation.	2
2 Proposed physiologic role of bilirubin: Oxidation-reduction cycles for bilirubin and GSH.	4
3 The absorption spectra of 50 μM (2.9 mg %) bilirubin dissolved in 10 mM Tris-HCl buffer, pH 7.5 in a Shimadzu UV -2450 Spectrophotometer.	36
4 Effect of 3 types of transition metal ions on bilirubin absorption spectra.	38
5 The effect of $\text{CuCl}_2[\text{Cu(II)}]$ on the absorption spectrum of bilirubin.	40
6 The effect of transition metal ion concentrations on the absorption spectra of bilirubin (500 μM).	41
7 The absorbance peaks of bilirubin or quercetin interacted with Cu(II) were compared with reference bilirubin absorption spectrum.	42
8 The effect of albumin on the absorption spectra of bilirubin-transition metal ion complex. Albumin as a molar ratio of bilirubin; 0.5:1.0, on the selected optimal bilirubin concentrations (500 μM).	44
9 The effect of albumin on the absorption spectra of bilirubin-transition metal ion complex. Albumin as a molar ratio of bilirubin; 1.0:1.0, on the selected optimal bilirubin concentrations (500 μM).	46
10 The effect of albumin on the absorption spectra of bilirubin-transition metal ion complex. Albumin as a molar ratio of bilirubin; 1.5:1.0, on the selected optimal bilirubin concentrations (500 μM).	47
11 Standard graph of DNA by Diphenylamine reaction.	48
12 The effect of increasing bilirubin concentrations interacted with fixed concentration of transition metal ion on DNA degradation <i>in vitro</i> .	50

Figure	Page
13 The acid soluble DNA hydrolyzed (%) estimated by the DPA reaction (resulted by the interactions of increasing with bilirubin fixed concentration of transition metal ion).	51
14 The effect of increasing interaction ion concentrations interacted with fixed concentration of bilirubin caused DNA degradation <i>in vitro</i> .	53
15 The acid soluble DNA hydrolyzed (%) estimated by the DPA reaction (resulted by the interactions of increasing transition metal ion with fixed concentration of bilirubin).	54
16 Degradation of calf thymus DNA by the bilirubin-Cu (II) complex detected by DPA reaction.	55
17 Time dependent kinetic of DNA degradations caused by the interactions of bilirubin with transition metal ions.	57
18 The time dependent kinetic of DNA degradations by the bilirubin-Cu(II) complex.	58
19 Standard graph of MDA assay.	60
20 The typical absorbance peak of MDA by TBA reaction.	60
21 The inhibition effect of free radical scavengers on DNA degradation caused by bilirubin-CuCl ₂ interaction.	62
22 The decrease in percentage of DNA hydrolysis by the inhibition effect of free radical scavengers (200 μM bilirubin-200 μM CuCl ₂).	63
23. The inhibition effect of free radical scavengers on DNA degradation caused by bilirubin-FeCl ₂ interaction.	65
24. The decrease in percentage of DNA hydrolysis by the inhibition effect of free radical scavengers(200 μM bilirubin-200 μM FeCl ₂).	66
25 The inhibition effect of free radical scavengers on DNA degradation caused by bilirubin-ZnCl ₂ interaction.	68
26 The decrease in percentage of DNA hydrolysis by the inhibition effect of free radical scavengers(200 μM bilirubin-200 μM FeCl ₂).	69

Figure	Page
27 Inhibition of calf thymus DNA degradation after treatment with bilirubin and transition metal ions in the presence of free radical scavengers (thiourea, sodium azide and mannitol).	71
28 The scanning of the reaction of bilirubin in the presence and absence of CuCl_2 .	74
29 The scanning of the reaction containing bilirubin and DNA in the presence and absence of CuCl_2 .	75
30 Detection of bilirubin induced Cu(I) production by bathocuproine.	102
31 The scanning of the reaction of bilirubin- CuCl_2 in the presence and absence of scavenger (thiourea).	103

ABBREVIATIONS

%	percentage
μg	microgram
μl	microlitre
μM	micromolar
Abs	absorbance
ATP	adenosine tri-phosphate
BSA	bovine serum albumin
CuCl ₂	cupric chloride
CO	carbon monoxide
CO ₂	carbon dioxide
Cu (II)	cupric ion
Cu	copper
dL	deciliter
DNA	deoxyribonucleic acid
DPA	diphenylamine
EDTA	ethylene diamine tetra-acetic acid
FeCl ₂	ferrous chloride
Fe(II)	ferrous ion
g	gram
HCl	hydrochloric acid
L	litre

LDH	lactate dehydrogenase
M	molar
MDA	malondialdehyde
mg	milligram
mL	millitre
mM	millimolar
N	Normal
N ₂	nitrogen
NaOH	sodium hydroxide
nm	nanometer
NO	nitric oxide
O ₂	oxygen
°C	degree Celsius
PARP	poly ADP ribose polymerase
RNA	ribonucleic acid
S	sodium azide
T	thiourea
TAE	Tris Acetate EDTA
TBA	thiobarbituric acid
TD	reaction contained DNA dissolved in Tris-HCl
TD500BC	reaction in Tris-HCl contained 500 µg DNA, bilirubin and cupric chloride
TDB	reaction in Tris-HCl contained DNA and bilirubin

TDBC	reaction in Tris-HCl contained DNA, bilirubin and cupric chloride
TDBCM	reaction in Tris-HCl contained DNA, bilirubin, cupric chloride and mannitol
TDBCS	reaction in Tris-HCl contained DNA, bilirubin, cupric chloride and sodium azide
TDBCT	reaction in Tris-HCl contained DNA, bilirubin, cupric chloride and thiourea
TDBF	reaction in Tris-HCl contained DNA, bilirubin and ferrous chloride
TDBFM	reaction in Tris-HCl contained DNA, bilirubin, ferrous chloride and mannitol
TDBFS	reaction in Tris-HCl contained DNA, bilirubin, ferrous chloride and sodium azide
TDBFT	reaction in Tris-HCl contained DNA, bilirubin, ferrous chloride and thiourea
TDBM	reaction in Tris-HCl contained DNA, bilirubin and mannitol
TDBS	reaction in Tris-HCl contained DNA, bilirubin and sodium azide
TDBT	reaction in Tris-HCl contained DNA, bilirubin and thiourea
TDBZ	reaction in Tris-HCl contained DNA, bilirubin and zinc chloride
TDBZM	reaction in Tris-HCl contained DNA, bilirubin, zinc chloride and mannitol
TDBZS	reaction in Tris-HCl contained DNA, bilirubin, zinc chloride and sodium azide
TDBZT	reaction in Tris-HCl contained DNA, bilirubin, zinc chloride and thiourea

TDC	reaction in Tris-HCl contained DNA and cupric chloride
TDM	reaction in Tris-HCl contained DNA and mannitol
TDS	reaction in Tris-HCl contained DNA and sodium azide
TDT	reaction in Tris-HCl contained DNA and thiourea
TDZ	reaction in Tris-HCl contained DNA and zinc chloride
TDZM	reaction in Tris-HCl contained DNA, zinc chloride and mannitol
TDZS	reaction in Tris-HCl contained DNA, zinc chloride and sodium azide
TDZT	reaction in Tris-HCl contained DNA, zinc chloride and thiourea
UCB	unconjugated bilirubin
UGT1A1	UDP glucuronosyltransferase 1 family, polypeptide A1
UV	ultraviolet
ZnCl ₂	zinc chloride
Zn	zinc