

REFERENCES

1. Weiss H-R, Goodall D. The treatment of adolescent idiopathic scoliosis (AIS) according to present evidence. a systematic review. *Eur J Phys Rehabil Med.* 2008; 44: 177-93.
2. Asher M, Burton D. Adolescent idiopathic scoliosis: natural history and long term treatment effects. *Scoliosis.* 2006.
3. Mirtz T, Thompson M, Greene L, Wyatt L, Akagi C. Adolescent idiopathic scoliosis screening for school, community, and clinical health promotion practice utilizing the PRECEDE-PROCEED model. *Chiropr Osteopat.* 2005; 13: 25.
4. Soutanis K, Payatakes A, Chouliaras V, Mandellos G, Pyrovolou N, Pliarchopoulou F, et al. Rare causes of scoliosis and spine deformity: experience and particular features. *Scoliosis.* 2007.
5. Wever DJ, Veldhuizen AG, Klein JP, Webb PJ, Nijenbanning G, Cool JC, et al. A biomechanical analysis of the vertebral and rib deformities in structural scoliosis. *Eur Spine J.* 1999; 8: 252-60.
6. Smyth RJ, Chapman KR, Wright TA, Crawford JS, Rebuck AS. Pulmonary function in adolescents with mild idiopathic scoliosis. *Thorax.* 1984; 39: 901-4.
7. Newton PO, editor. *Adolescent Idiopathic Scoliosis.* Rosemont: American Academy of Orthopaedic Surgeons; 2004.
8. Newton PO, Faro FD, Gollogly S, Betz RR, Lenke LG, Lowe TG. Results of preoperative pulmonary function testing of adolescents with idiopathic

- scoliosis. a study of six hundred and thirty-one patients. *J Bone Joint Surg Am.* 2005; 87: 1937-46.
9. Barrios C, Perez-Encinas C, Maruenda JI. Significant ventilatory functional restriction in adolescent with mild or moderate scoliosis during maximal exercise tolerance test. *Spine.* 2005; 30: 1619-5.
 10. Boyer J, Amin N, Taddonio R, Dozor AJ. Evidence of airway obstruction in children with idiopathic scoliosis. *Chest.* 1996; 109: 1532-5.
 11. Szeinberg A, Cannym GJ, Rashed N, Veneruso G, Levion H. Forced vital capacity and maximal respiratory pressure in patients with mild and moderate scoliosis. *Pediatr Pulmonol.* 1988; 4: 8-12.
 12. Alves VLS, Stirbulov R, Avanzi O. Impact of a physical rehabilitation program on the respiratory function of adolescent with idiopathic scoliosis. *Chest.* 2006; 130: 500-5.
 13. Reamy BV, Slakey JB. Adolescent idiopathic scoliosis: review and current concepts. *Am Fam Physician.* 2001; 64: 111-6.
 14. Weinstein SL, Dolan L, Cheng JCY, Danielsson A, Morcuende JA. Adolescent idiopathic scoliosis. *Lancet.* 2008; 371: 1527-37.
 15. Roach JW. Adolescent idiopathic scoliosis. *Orthop Clin North Am.* 1999; 30: 353-65.
 16. Grivas TB, Vasiliadis E, Mouzakis V, Mihas C, Koufopoulos G. Association between adolescent idiopathic scoliosis prevalence and age at menarche in different geographic latitudes. *Scoliosis.* 2006.
 17. Scoliosis research society. 2008. "Type of scoliosis: adolescents." [Online]. Available <http://www.srs.org/patients/adolescent/> (6 November 2008)

18. Farady JA. Current principles in the nonoperative management of structural adolescent idiopathic scoliosis. *Phys Ther.* 1983; 63: 512-22.
19. Greiner KA. Adolescent idiopathic scoliosis: radiologic decision-making. *Am Fam Physician.* 2002; 65: 1817-22.
20. Chun EM, Suh SW, Modi HN, Kang EY, Hong SJ, Song H. The change in ratio of convex and concave lung volume in adolescent idiopathic scoliosis: a 3D CT scan based cross sectional study of effect of severity of curve on convex and concave lung volume in 99 cases. *Eur Spine J.* 2007; 17: 224-9.
21. Wasserman K, Hansen JE, Sue DY, Stringer WW, Whipp BJ, editors. *Principles of exercise testing and interpretation: including pathophysiology and clinical applications.* 4th ed. Philadelphia: Lippincott Williams & Wilkins; 2005. p. 10-59.
22. Bolton CF, Chen R, Wijdicks EFM, Zifko UA. Anatomy and physiology of nervous system control of respiration. *Neurology of breathing.* Philadelphia: Butterworth Heinemann; 2004. p. 20-32.
23. Miller MR, Hanskinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J.* 2005; 26: 319-38.
24. Powers SK, Howley ET. Respiration during exercise. *Exercise Physiology.* 6th ed. New York: McGraw Hill; 2007. p. 200-29.
25. Rosthstein JM, Roy SH, Wolf SL, editors. Pulmonary anatomy and pulmonary therapy. *The rehabilitation specialist's handbook.* 3rd ed. Philadelphia; 2005. p.436.

26. Astrand PO, Rodahl K, Dahl HA, Stromme SB. Respiration. Textbook of work physiology. Physiological bases of exercise. 4th ed. Ontario: Human Kinetics; 2003. p. 177-212.
27. ATS/ERS statement on respiratory muscle testing. Am J Respir Crit Care Med. 2002; 166: 518-624.
28. Chen H, Kuo C. Relationship between respiratory muscle function and age, sex and other factors. News Physiol Sci. 1989; 66: 943-8.
29. Clanton T, Diaz PT. Clinical assessment of the respiratory muscles. Phys Ther. 1995; 75: 983-94.
30. Arena R, Myers J, Williams MA, Gulati M, Kligfield P, Balady GJ, et al. Assessment of functional capacity in clinical and research setting: a scientific statement from the American Heart Association committee on exercise, rehabilitation and prevention of the council on clinical cardiology and the council on cardiovascular nursing. Circulation. 2007; 116: 329-43.
31. ATS statement: guidelines for the six-minute walk test. Am J Respir Crit Care Med. 2002; 166: 111-7.
32. Fleg JL, Pina IL, Balady GJ, Chaitman BR, Fletcher B, Lavie C, et al. Assessment of functional capacity in clinical and research applications: an advisor from the committee on exercise, rehabilitation, and prevention, council on clinical cardiology, American Heart Association. Circulation. 2000; 102: 1591-7.
33. Carter R, Holiday DB, Nwasuruba C, Stocks J, Grothues C, Tjep B. 6-minute walk work for assessment of functional capacity in patients with COPD. Chest. 2003; 123: 1408-15.

34. Cooper KH. A means of assessing maximal oxygen intake: correlation between field and treadmill testing. *JAMA*. 1968; 203: 201-4.
35. Reybrouck T. Clinical usefulness and limitations of the 6-minute walk test in patients with cardiovascular or pulmonary disease. *Chest*. 2003; 123: 325-7.
36. Solway S, Brooks D, Lacasse Y, Thomas S. A qualitative systematic overview of the measurement properties of functional walk tests used in the cardiorespiratory domain. *Chest*. 2001; 119: 256-70.
37. Luxton N, Alison JA, Wu J, Mckey MG. Relationship between field walking tests and incremental cycle ergometry in COPD. *Respirology*. 2008; 13: 856-62.
38. Taylor MJ, Mazzone M, Wrotniak BH. Outcome of an exercise and education intervention for children who are over weight. *Pediatr Phys Ther*. 2005: 180-8.
39. Casas A, Vilaro J, Rabinovich R, Mayer A, Barbera AJ, Rodriguez-Roisin R, et al. Encouraged 6-min walking test indicates maximum sustainable exercise in COPD patients. *Chest*. 2005; 128: 55-61.
40. Enright PL. The six-minute walk test. *Respir Care*. 2003; 48: 783-5.
41. Catherine D, Robert M, Abdissa N, Salim Y. Reliability, validity and responsiveness of the six-minute walk test in patients with heart failure clinical investigations. *Am Heart J*. 2001; 142: 698-703.
42. Li AM, Yin L, Yu CCW, Tsang T, So HK, Wong E, et al. The six-minute walk test in healthy children:reliability and validity. *Eur Respir J*. 2005; 25: 1057-60.

43. Rostago C, Gensini GF. Six minute walk test: a simple and useful test to evaluate functional capacity in patients with heart failure. *Intern Emerg Med.* 2007; 3: 205-12.
44. Kervio G, Ville NS, Leclercq C, Daubert J, Carre F. Intensity and daily reliability of the six-minute walk test in moderate chronic heart failure patients. *Arch Phys Med Rehabil* 2004; 85: 1513-8.
45. Cahalin LP, Mathier MA, Semigran MJ, Dec GW, Disalvo TG. The six-minute walk test predicts peak oxygen uptake and survival in patients with advanced heart failure. *Chest.* 1996; 110: 325-32.
46. Li AM, Yin L, Au JT, So HK, Tsang T, Wong E, et al. Standard reference for the six-minute walk test in healthy children aged 7 to 16 years. *Am J Respir Crit Care Med.* 2007; 176: 174-6.
47. Lammers AE, Hislop AA, Flynn Y, Haworth SG. The 6-minute walk test: normal values for children of 4-11 years of age. *Arch Dis Child.* 2008; 93: 464-8.
48. Geiger R, Strasak A, Treml B, Gasser K, Kleinsaser A, Fischer V, et al. Six-minute walk test in children and adolescents. *J Pediatr.* 2007; 150: 395-9.
49. Hassan J, van der Net J, Helders PJM, Prakken BJ, Takken T. Six-minute walk test in children with chronic conditions. *Br J Sports Med.* 2008.
50. Mattiello R, Sarria EE, Stein R, Fischer GB, Mocelin HT, Barreto SSM, et al. Functional capacity assessment in children and adolescents with post-infectious broncholitis obliterans. *J Pediatr.* 2008; 84: 337-43.
51. Trooster T, Gosselink R, Decramer M. Six minute walking distance in healthy elderly subjects. *Eur Respir J.* 1999; 14: 270-4.

52. Wu G, Sanderson B, Bittner V, Ala B. The 6-minute walk test: how important is the learning effect? *Am Heart J.* 2003; 146: 129-33.
53. Eaton T, Young P, Milne D, Wells AU. Six-minute walk, maximal exercise tests reproducibility in fibrotic interstitial pneumonia. *Am J Respir Crit Care Med.* 2005; 171: 1150-7.
54. Kervio G, Carre F, Ville N. Reliability and intensity of the six-minute walk test in healthy elderly subjects. *Med Sci Sports Exercise.* 2003; 35: 169-74.
55. Brooks D, Solway S, Weinacht K, Wang D, Thomas S. Comparison between an indoor and an outdoor 6-minute walk test among individuals with chronic obstructive pulmonary disease. *Arch Phys Med Rehabil* 2003; 84: 873-6.
56. Lipkin DP, Scriven AJ, Crake T, Poole-wilson PA. Six-minute walking test for assessing exercise capacity in chronic heart failure. *Br Med J.* 1986; 292: 653-5.
57. Davies D. *Growth. Child development.* New York: The Guilford Press; 1999. p. 40-42.
58. Bansal R, Simakajornboon N, Cangiamella S, Beckerman RC, Kiernan M. The measurement of respiratory muscle function in adolescent idiopathic scoliosis. *Chest-Poster presentation.* 2005: 354s-5s.
59. Smyth RJ, Rebuck AS. Maximal inspiratory and expiratory pressure in adolescents. Normal values. *Chest.* 1984; 86: 568-72.
60. Jones RS, Kennedy JD, Hasham F, Owen R, Taylor JF. Mechanical inefficiency of the thoracic cage in scoliosis. *Thorax.* 1981; 36: 456-61.

61. Weinstein SL, Zavala DC, Ponseti IV. Idiopathic scoliosis: long-term follow-up and prognosis in untreated patients. *J Bone Joint Surg Am.* 1981; 63: 702-12.
62. Weinstein SL, Dolan LA, Spratt KF. Health and function of patients with untreated idiopathic scoliosis: A 50 year natural history study. *JAMA.* 2003; 289: 559-67.
63. Whaley MH, Brubaker PH, Otto RM, editors. *ACSM's guidelines for exercise testing and prescription* 7th ed: Lippincott Williams & Wilkins; 2006. p. 80.
64. Milbrandt TA, Sucato DJ. The position of the aorta relative to the spine in patients with left thoracic scoliosis: a comparison with normal patients. *Spine.* 2007; 32: E348-53.
65. Sucato DJ, Duchene C. The position of the aorta relative to the spine: a comparison of patients with and without idiopathic scoliosis. *J Bone Joint Surg Am.* 2003; 85: 1461-9.
66. Sevastik B, Xiong B, Hedlund R, Sevastik J. The position of the aorta in relation to the vertebra in patients with idiopathic thoracic scoliosis. *Surg Radiol Anat.* 1996; 18: 51-6.