

## REFERENCES

1. Wood O. Clayton's electrotherapy. In: Kitchen S, Bazin S, editors. Sensory and motor nerve activation. 10th ed. London: WB Saunders company Ltd; 1996. p. 77-8.
2. Popovic MR., Thrasher TA, Zivanovic V, Takaki J, Hajek V. Neuroprostheses for retraining reaching and grasping functions in severe hemiplegic patients. Neuromodul. 2005;8:58-72.
3. O'Sullivan SB. Examination of motor function: Motor control and motor learning. In: O'Sullivan SB, Thomas SJ, editors. Physical rehabilitation. 5 th ed. Philadelphia: F.A Davis Company; 2007. p. 233-4.
4. Nakayama H, Jørgensen HS, Raaschou HO, Olsen TS. Recovery of upper extremity function in stroke patients: The Copenhagen stroke study. Arch Phys Med Rehabil. 1994;75:394-8.
5. Hara Y. Neurorehabilitation with new functional electrical stimulation for hemiparetic upper extremity in stroke patients. J Nippon Med Sch. 2008;75:4-14.
6. Weinstein CJ, Rose DK, Tan SM, Lewthwaite R, Chui HC, Azen SP. A randomized controlled comparison of upper-extremity rehabilitation strategies in acute stroke: a pilot study of immediate and long-term outcomes. Arch Phys Med Rehabil. 2004;85:620-8.

7. Jette DU, Latham NK, Smout RJ, Gassaway J, Slavin MD, Horn SD. Physical therapy interventions for patients with stroke in inpatient rehabilitation facilities. *Phys Ther.* 2005;85:238-48.
8. Dickstein R, Hocherman S, Pillar T, Shaham R. Stroke rehabilitation. Three exercise therapy approaches. *Phys Ther.* 1986;66:1233-8.
9. Taub.E, Morris.D.M. Constraint-induced movement therapy to enhance recovery after stroke. *Curr Atheroscler Rep.* 2001;3:279-86.
10. Bonifer NM, Anderson KM, Arciniegas DB. Constraint-induced movement therapy after stroke: Efficacy for patients with minimal upper-extremity motor ability. *Arch Phys Med Rehabil.* 2005;86:1867-73.
11. Wolf SL, Blanton S, Baer H, Breshears J, Butler AJ. Repetitive task practice: A critical review of constraint-induced movement therapy in stroke. *Neurologist.* 2002;8:325-38.
12. Rathkolb O, Baykoushev S, Baykousheva V. Myobiofeedback in motor reeducation of wrist and fingers after hemispherical stroke. *Electromyogr Clin Neurophysiol.* 1990;30:89-92.
13. Huang H, Ingalls T, Olson L, Ganley K, Rikakis T, He J. Interactive multimodal biofeedback for task-oriented neural rehabilitation. *Conf Proc IEEE Eng Med Biol Soc.* 2005;3:2547-50.
14. Hesse S, Schulte-Tigges G, Konrad M, Bardeleben A, Werner C. Robot-assisted arm trainer for the passive and active practice of bilateral forearm and wrist movements in hemiparetic subjects. *Arch Phys Med Rehabil.* 2003;84:915-20.

15. Masiero S, Celia A, Rosati G, Armani M. Robotic-assisted rehabilitation of the upper limb after acute stroke. *Arch Phys Med Rehabil.* 2007;88:142-9.
16. de Kroon JR, IJzerman MJ, Lankhorst GJ, Zilvold G. Electrical stimulation of the upper limb in stroke: Stimulation of the extensors of the hand vs. alternate stimulation of flexors and extensors. *Am J Phys Med Rehabil.* 2004;83:592-600.
17. Alon G, Levitt AF, McCarthy PA. Functional electrical stimulation enhancement of upper extremity functional recovery during stroke rehabilitation: A pilot study. *Neurorehabil Neural Repair.* 2007;21:207-15.
18. Powell J, Pandyan AD, Granat M, Cameron M, Stott DJ. Electrical stimulation of wrist extensors in poststroke hemiplegia. *Stroke.* 1999;30:1384-9.
19. Ring H, Rosenthal N. Controlled study of neuroprosthetic functional electrical stimulation in sub-acute post-stroke rehabilitation. *J Rehabil Med.* 2005;37:32-6.
20. Sullivan JE, Hedman LD. Effects of home-based sensory and motor amplitude electrical stimulation on arm dysfunction in chronic stroke. *Clin Rehabil.* 2007;21:142-50.
21. Muraoka Y. Development of an EMG recording device from stimulation electrodes for functional electrical stimulation. *Front Med Biol Eng.* 2002;11:323-33.
22. Weingarden HP, Zeilig G, Heruti R, et al. Hybrid functional electrical stimulation orthosis system for the upper limb: Effects on spasticity in chronic stable hemiplegia. *Am J Phys Med Rehabil.* 1998;77:276-81.

23. Dunning K, Berberich A, Albers B, et al. A four-week, task-specific neuroprosthesis program for a person with no active wrist or finger movement because of chronic stroke. *Phys Ther.* 2008;88:397-405.
24. Hara Y, Ogawa S, Tsujiuchi K, Muraoka Y. A home-based rehabilitation program for the hemiplegic upper extremity by power-assisted functional electrical stimulation. *Disabil Rehabil.* 2008;30:296-304.
25. Sullivan JE, Hedman LD. A home program of sensory and neuromuscular electrical stimulation with upper-limb task practice in a patient 5 years after a stroke. *Phys Ther.* 2004;84:1045-54.
26. Alon G, Levitt AF, McCarthy PA. Functional electrical stimulation (FES) may modify the poor prognosis of stroke survivors with severe motor loss of the upper extremity: A preliminary study. *Am J Phys Med Rehabil.* 2008;87:627-36.
27. Hashimoto H, Kuriyama Y, Imakita S, Sawada T, Naritomi H. Regional cerebral hemodynamics among hypertensive patients with lacunar infarctions during blood pressure control in subacute and chronic phases. *J Stroke Cerebrovasc Dis.* 1997;6:319-24.
28. Caplan LR. Caplan's stroke: A clinical approach. 3rd ed. Boston: Butterworth-Heinemann; 2000.
29. O'Sullivan SB. stroke. In: O'Sullivan SB, Thomas SJ, editors. Physical rehabilitation. 5 th ed. Philadelphia: F.A. Davis company; 2007. p. 720.
30. Parker VM, Wade DT, Langton RH. Loss of arm function after stroke: Measurement, frequency and recovery. *Int Rehabil Med.* 1986;8:69-73.

31. de Kroon JR, IJzerman MJ, Chae J, Lankhorst GJ, Zilvold G. Relation between stimulation characteristics and clinical outcome in studies using electrical stimulation to improve motor control of the upper extremity in stroke. *J Rehabil Med.* 2005;37:65-74.
32. Bohannon RW. Relationship between active range of motion deficits and muscle strength and tone at the elbow in patients with hemiparesis. *Clin Rehabil.* 1991;5:219-24.
33. Canning CG, Ada L, O'Dwyer N. Slowness to develop force contributes to weakness after stroke. *Arch Phys Med Rehabil.* 1999;80:66-70.
34. MacComas AJ, Silica REP, Upton ARM, aguilera W. Functional changes in motor neuron of hemiparetic patients. *J Neurol Neurosurg Psychiatry.* 1973;36:183-93.
35. Lance JW. Symposium synopsis. In: Fildman RG, Young RR, Koella WP, editors. *Spasticity: disordered motor control.* Miami: Symposia specialists; 1980. p. 485-94.
36. Mukul Mukherjee P, Lisa K. McPeak, MD, John B. Redford, PhD, Chao Sun, MD, Wen Liu, PhD. The effect of electro-acupuncture on spasticity of the wrist joint in chronic stroke survivors wrist joint in chronic stroke survivors. *Arch Phys Med Rehabil.* 2007;88:159-66.
37. Julius PA. Sensorimotor neurophysiology and the basis of neurofacilitation therapeutic techniques. In: Brandstater ME, Basmajian JV, editors. *Stroke rehabilitation.* Baltimore: Williams and Wilkins; 1987. p. 109-78.

38. Davies PM. Steps to follow: A guild to the treatment of adult hemiplegia. Berlin: Springer-Verlag; 1985.
39. Lin FM, Sabbahi M. Correlation of spasticity with hyperactive stretch reflexes and motor dysfunction in hemiplegia. *Arch Phys Med Rehabil.* 1999;80:526-30.
40. Musampa NK, Mathieu PA, Levin M, F. Relationship between stretch reflex thresholds and voluntary arm muscle activation in patients with spasticity. *Exp Brain Res.* 2007;181:579-93.
41. Gillen G. Upper extremity function and management. In: Gillen G, Burkhardt A, editors. *Stroke rehabilitation: A functional-based approach.* St Luris Baltimore: Mosby; 1998.
42. Hendricks HT, IJzerman MJ, de Kroon JR, in 't Groen FA, Zilvold G. Functional electrical stimulation by means of the 'Ness Handmaster Orthosis' in chronic stroke patients: an exploratory study. *Clin Rehabil.* 2001;15:217-20.
43. Mayer NH, Whyte J, Wannstedt G, Ellis CA. Comparative impact of 2 botulinum toxin injection techniques for elbow flexor hypertonia. *Arch Phys Med Rehabil.* 2008;89:982-7.
44. Raine S. Defining the Bobath concept using the Delphi technique. *Physiother Res Int.* 2006;11:4-13.
45. Aprile I, Di Stasio E, Romitelli F, et al. Effects of rehabilitation on quality of life in patients with chronic stroke. *Brain Inj.* 2008;22:451-6.
46. Duncan PW. Stroke disability. *Phys Ther.* 1994;74:399-407.

47. Manimmanakorn N, Vichiansiri R, Nuntharuksa C, Permsirivanich W, Kuptniratsaikul V. Quality of life after stroke rehabilitation among urban vs. rural patients in Thailand. *J Med Assoc Thai.* 2008;91:394-9.
48. Bach-y-Rita P. Process of recovery from stroke. In: Brandstater ME, Basmajian JV, editors. *Stroke rehabilitation.* Baltimore: Williams and Wilkins; 1987. p. 80-108.
49. Bennett SE, Karnes JL. *Neurological disabilities: assessment and treatment.* Philadelphia: Lippincott; 1998.
50. Carey JR, Kimberley TJ, Lewis SM, Auerbach EJ, Dorsey L, Rundquist P. Analysis of fMRI and finger tracking training in subjects with chronic stroke. *Brain.* 2002;125:773-88.
51. Nelles G, Jentzen W, Juueptner M, Muller S, Diener H. Arm training induced brain plasticity in stroke studied with serial positron emission tomography. *2001;13:1146-54.*
52. Gritsenko V, Prochazka A. A functional electric stimulation-assisted exercise therapy system for hemiplegic hand function. *Arch Phys Med Rehabil.* 2004;85:881-5.
53. Wang RY, Yang YR, Tsai MW, Wang WTJ, Chan RC. Effects of functional electric stimulation on upper limb motor function and shoulder range of motion in hemiplegic patients. *Am J Phys Med Rehabil* 2002;81:283–90.
54. Faghri PD, Rodgers MM, Glaser RM, Bors JG, Ho C, Akuthota P. The effects of functional electrical stimulation on shoulder subluxation, arm function

- recovery, and shoulder pain in hemiplegic stroke patients. *Arch Phys Med Rehabil.* 1994;75:73-9.
55. Powell J, Pandyan AD, Granat M, Cameron M, Stott DJ. Electrical stimulation of wrist extensors in poststroke hemiplegia. *Stroke.* 1999;30(7):1384-9.
56. Alon G, Alan LF, Patricia MA. Functional electrical stimulation enhancement of upper extremity functional recovery during stroke rehabilitation: A pilot study. *Neurorehabil Neural Repair.* 2007;21:207-15.
57. Thrasher TA, Popovic MR. Functional electrical stimulation of walking: Function, exercise and rehabilitation. *Ann Readapt Med Phys.* 2008;51:452-60.
58. Glanz M, Klawansky S, Stason W, Berkey C, Chalmers TC. Functional electrostimulation in poststroke rehabilitation: A meta-analysis of the randomized controlled trials. *Arch Phys Med Rehabil.* 1996;77:549-53.
59. Vitenzon AS, Mironov EM, Petrushanskaia KA. Functional electrostimulation of muscles as a method for restoring motor functions. *Neurosci Behav Physiol.* 2005;35:709-14.
60. Hummelsheim H, Maier-Loth ML, Eickhof C. The functional value of electrical muscle stimulation for the rehabilitation of the hand in stroke patients. *Scand J Rehabil Med.* 1997;29:3-10.
61. Kobayashi H, Onishi H, Ihashi K, Yagi R, Handa Y. Reduction in subluxation and improved muscle function of the hemiplegic shoulder joint after therapeutic electrical stimulation. *J Electromyogr Kinesiol.* 1999;9:327-36.

62. Hara Y, Ogawa S, Muraoka Y. Hybrid power-assisted functional electrical stimulation to improve hemiparetic upper-extremity function. *Am J Phys Med Rehabil.* 2006;85:977-85.
63. Ring H, Rosenthal N. Controlled study of neuroprosthetic functional electrical stimulation in sub-acute post-stroke rehabilitation. *J Rehabil Med.* 2005;37:32-6.
64. Rossini PM, Calautti C, Pauri F, Baron JC. Post-stroke plastic reorganisation in the adult brain. *Lancet Neurol.* 2003;2:493-502.
65. Wu CW, van Gelderen P, Hanakawa T, Yaseen Z, Cohen LG. Enduring representational plasticity after somatosensory stimulation. *Neuroimage.* 2005;27:872-84.
66. Wu CW, Hyae-Jung S, Cohen LG. Influence of electric somatosensory stimulation on paretic-hand function in chronic stroke. *Arch Phys Med Rehabil.* 2006;87:351-7.
67. Tyson S. Use of transcutaneous nerve stimulation to treat sensory loss after stroke. *Physiother Res Int.* 2003;8:53-7.
68. Wu CW, Hyae-Jung S, Cohen LG. Influence of electric somatosensory stimulation on paretic-hand function in chronic stroke. *Arch Phys Med Rehabil.* 2006;87:351-7.
69. Elovic EP, Brashear A, Kaelin D, et al. Repeated treatments with botulinum toxin type a produce sustained decreases in the limitations associated with focal upper-limb poststroke spasticity for caregivers and patients. *Arch Phys Med Rehabil.* 2008;89:799-806.

70. Hooker DN. Therapeutic modalities for allied health professionals. In: Prentice WE, editor. Electrical stimulating currents. New York: McGraw-Hill; 1998. p. 73-117.
71. Scott O. Stimilative effects. In: Kitchen S, editor. Electrotherapy: Evidence-based practice. 11th ed. Edinburgh: Churchill Livingstone; 2002. p. 113-28.
72. Enoka RM. Neural adaptations with chronic physical activity. J Biomech. 1997;30:447-55.
73. Feys H, De Weerd W, Verbeke G. Early and repetitive stimulation of the arm can substantially improve the long-term outcome after stroke: 5-year follow-up study of a randomized trial. Stroke. 2004;4:924-9.
74. Carr JH, Shepherd RB. A motor relearning programme for stroke. Oxford: Heinemann Medical Books; 1987.
75. Ang JH, Man DW. The discriminative power of the Wolf motor function test in assessing upper extremity functions in persons with stroke. Int J Rehabil Res. 2006;29:357-61.
76. Wolf SL, Catlin PA, Ellis M, Archer AL, Morgan B, Piacentino A. Assessing Wolf motor function test as outcome measure for research in patients after stroke. Stroke. 2001;32:1635-9.
77. Whitall J, Savin DN, Jr Harris-Love M, Waller SM. Psychometric properties of a modified Wolf motor function test for people with mild and moderate upper-extremity hemiparesis. Arch Phys Med Rehabil. 2006;87:656-60.

78. Morris DM, Usatte G, Crago JE, Cook III EW, Taub E. The reliability of the Wolf motor function test for assessing upper extremity function after stroke. *Arch Phys Med Rehabil.* 2001;82:750-5.
79. Lin KC, Hsieh YW, Wu CY, Chen CL, Jang Y, Liu JS. Minimal detectable change and clinically important difference of the Wolf Motor Function Test in stroke patients. *Neurorehabil Neural Repair.* 2009;23:429-34.
80. Schmitt JS, Di Fabio RP. Reliable change and minimum important difference (MID) proportions facilitated group responsiveness comparisons using individual threshold criteria. *J Clin Epidemiol.* 2004;57:1008-18.
81. Johnson L, Selfe J. Measurement of mobility following stroke: A comparison of the Modified Rivermead Mobility Index and the Motor Assessment Scale. *Physiother.* 2004;90:132-8.
82. Hsueh IP, Hsieh CL. Responsiveness of two upper extremity function instruments for stroke inpatients receiving rehabilitation. *Clin Rehabil.* 2002;16:617-24.
83. Poole JL, Whitney SL. Motor assessment scale for stroke patients: Concurrent validity and interrater reliability. *Arch Phys Med Rehabil.* 1988;69:195-7.
84. Blackburn M, van Vliet P, Mockett SP. Reliability of measurements obtained with the modified Ashworth scale in the lower extremities of people with stroke. *Phys Ther.* 2002;82:25-34.
85. Wallace D, Duncan PW, Lai SM. Comparison of the responsiveness of the Barthel Index and the Motor Component of the Functional Independence

- Measure in stroke : The impact of using different methods for measuring responsiveness. *J Clin Epidemiol.* 2002;55:922-8.
86. Green J, Forster A, Young J. A test-retest reliability study of the Barthel Index, the Rivermead Mobility Index, the Nottingham Extended Activities of Daily Living Scale and the Frenchay Activities Index in stroke patients. *Disabil Rehabil.* 2001;23:670-6.
  87. Schultz-Larsen K, Lomholt RK, Kreiner S. Mini-Mental Status Examination: A short form of MMSE was as accurate as the original MMSE in predicting dementia. *J Clin Epidemiol.* 2007;60:260-7.
  88. Sithinamsuwan P, Niyasom S, Nidhinandana S, Supasyndh O. Dementia and depression in end stage renal disease: Comparison between hemodialysis and continuous ambulatory peritoneal dialysis. *J Med Assoc Thai.* 2005;88:s141-7.
  89. Dickinson C, Shim M. The influence of manual dexterity on reading speed with a hand-held magnifier. *Invest Ophthalmol Vis Sci.* 2007;48:4368-74.
  90. Suksriwan S. Circuit upper arm training in patients with hemiplegia [MS thesis]: Chiang Mai University; 2006.
  91. Thielman GT, Dean CM, Gentile AM. Rehabilitation of reaching after stroke: task-related training versus progressive resistive exercise. *Arch Phys Med Rehabil.* 2004;85:1613-8.
  92. Kwakkel G, Kollen BJ, van der Grond J, Prevo AJ. Probability of regaining dexterity in the flaccid upper limb: Impact of severity of paresis and time since onset in acute stroke. *Stroke.* 2003;34:2181-6.

93. Wolf SL, McJunkin JP, Swanson ML, Weiss PS. Pilot normative database for the Wolf Motor Function Test. *Arch Phys Med Rehabil.* 2006;87:443-5.
94. Hedman LD, Sullivan JE, Hilliard MJ, Brown DM. Neuromuscular electrical stimulation during task-oriented exercise improves arm function for an individual with proximal arm dysfunction after stroke. *Am J Phys Med Rehabil.* 2007;86:592-6.
95. Hill-Hermann V, Strasser A, Albers B, et al. Task-specific, patient-driven neuroprosthesis training in chronic stroke: results of a 3-week clinical study. *Am J Occup Ther.* 2008;62:466-72.
96. Pang MY, Harris JE, Eng JJ. A community-based upper-extremity group exercise program improves motor function and performance of functional activities in chronic stroke: A randomized controlled trial. *Arch Phys Med Rehabil.* 2006;87:1-9.
97. Andrews K, Brocklehurst JC, Richards B, Laycock PJ. The rate of recovery from stroke and its measurements. *Int Rehabil Med.* 1981;3:155-61.
98. Skibbeck CE, Wade DT, Hewer RL, Woud VA. Recovery after stroke. *J Neurol Neurosurg Psychiatry.* 1983;46:5-8.
99. Sabari JS, Lim AL, Velozo CA, Lehman L, Kieran O, Lai JS. Assessing arm and hand function after stroke: A validity test of the hierarchical scoring system used in the motor assessment scale for stroke. *Arch Phys Med Rehabil.* 2005;86:1609-15.
100. Roberts JM, Wilson K. Effect of stretching duration on active and passive range of motion in the lower extremity. *Br J Sports Med.* 1999;33:259-63.

101. Barker RN, Brauer S, Carson R. Training-induced changes in the pattern of triceps to biceps activation during reaching tasks after chronic and severe stroke. *Exp Brain Res.* 2009;196:483-96.
102. Chang JJ, Tung WL, Wu WL, Huang MH. Effects of robot-aided bilateral force-induced isokinetic arm training combined with conventional rehabilitation on arm motor function in patients with chronic stroke. *Arch Phys Med Rehabil.* 2007;88:1332-8.
103. Peurala SH, Pitkänen K, Sivenius J, Tarkka IM. Cutaneous electrical stimulation may enhance sensorimotor recovery in chronic stroke. *Clin Rehabil.* 2002;16:709-16.
104. Bütefisch C, Hummelsheim H, Denzler P, Mauritz KH. Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. *J Neurol Sci.* 1995;130:59-68.