

REFERENCES

- Abbasova, S.G., Shcheprova Zh, M., Laman, A.G., Shepelyakovskaya, A.O., Baidakova, L.K., Rodionov, I.L. and Kushlinskii, N.E. (2007) Epitope mapping of human Fas using peptide phage display. *Bull Exp Biol Med* 144, 515-9.
- Afanasieva, T.A., Wittmer, M., Vitaliti, A., Ajmo, M., Neri, D. and Klemenž, R. (2003) Single-chain antibody and its derivatives directed against vascular endothelial growth factor: application for antiangiogenic gene therapy. *Gene Ther* 10, 1850-9.
- Altruda, F., Cervella, P., Gaeta, M.L., Daniele, A., Giancotti, F., Tarone, G., Stefanuto, G. and Silengo, L. (1989) Cloning of cDNA for a novel mouse membrane glycoprotein (gp42): shared identity to histocompatibility antigens, immunoglobulins and neural-cell adhesion molecules. *Gene* 85, 445-51.
- Alvarez, R.D., Barnes, M.N., Gomez-Navarro, J., Wang, M., Strong, T.V., Arafat, W., Arani, R.B., Johnson, M.R., Roberts, B.L., Siegal, G.P. and Curiel, D.T. (2000) A cancer gene therapy approach utilizing an anti-erbB-2 single-chain antibody-encoding adenovirus (AD21): a phase I trial. *Clin Cancer Res* 6, 3081-7.
- Azzazy, H.M. and Highsmith, W.E., Jr. (2002) Phage display technology: clinical applications and recent innovations. *Clin Biochem* 35, 425-45.
- Baneyx, F. and Mujacic, M. (2004) Recombinant protein folding and misfolding in *Escherichia coli*. *Nat Biotechnol* 22, 1399-408.

- Barbas, C.F.3rd., Burton, D.R., Scott, J.K., Silverman, G.J. (2001) Phage display: a laboratory manual. Cold Spring Harbor Laboratory Press, NY.
- Bardwell, J.C., McGovern, K. and Beckwith, J. (1991) Identification of a protein required for disulfide bond formation *in vivo*. Cell 67, 581-9.
- Barouch, D.H. and Nabel, G.J. (2005) Adenovirus vector-based vaccines for human immunodeficiency virus type 1. Hum Gene Ther 16, 149-56.
- Bass, S., Greene, R. and Wells, J.A. (1990) Hormone phage: an enrichment method for variant proteins with altered binding properties. Proteins 8, 309-14.
- Baum, C., Dullmann, J., Li, Z., Fehse, B., Meyer, J., Williams, D.A. and von Kalle, C. (2003) Side effects of retroviral gene transfer into hematopoietic stem cells. Blood 101, 2099-114.
- Bedzyk, W.D., Weidner, K.M., Denzin, L.K., Johnson, L.S., Hardman, K.D., Pantoliano, M.W., Asel, E.D. and Voss, E.W., Jr. (1990) Immunological and structural characterization of a high affinity anti-fluorescein single-chain antibody. J Biol Chem 265, 18615-20.
- Beerli, R.R., Wels, W. and Hynes, N.E. (1994) Autocrine inhibition of the epidermal growth factor receptor by intracellular expression of a single-chain antibody. Biochem Biophys Res Commun 204, 666-72.
- Berditchevski, F., Chang, S., Bodorova, J. and Hemler, M.E. (1997) Generation of monoclonal antibodies to integrin-associated proteins. Evidence that alpha3beta1 complexes with EMMPRIN/basigin/OX47/M6. J Biol Chem 272, 29174-80.

Bergelson, J.M., Cunningham, J.A., Droguett, G., Kurt-Jones, E.A., Krithivas, A., Hong, J.S., Horwitz, M.S., Crowell, R.L. and Finberg, R.W. (1997) Isolation of a common receptor for Coxsackie B viruses and adenoviruses 2 and 5. *Science* 275, 1320-3.

Berk, A.J. (2007) Adenoviridae: The viruses and their replication. In "Fields Virology" (D. M. Knipe, and P. M. Howley, 5th Eds.) Vol. II. Lippincott Williams & Wilkins, a Wolters Kluwer Business, Philadelphia.

Berntzen, G., Andersen, J.T., Ustgard, K., Michaelsen, T.E., Mousavi, S.A., Qian, J.D., Kristiansen, P.E., Lauvrak, V. and Sandlie, I. (2009) Identification of a high affinity FcγRIIA-binding peptide that distinguishes FcγRIIA from FcγRIIB and exploits FcγRIIA-mediated phagocytosis and degradation. *J Biol Chem* 284, 1126-35.

Beyer, F., Doebis, C., Busch, A., Ritter, T., Mhashilkar, A., Marasco, W.M., Laube, H., Volk, H.D. and Seifert, M. (2004) Decline of surface MHC I by adenoviral gene transfer of anti-MHC I intrabodies in human endothelial cells-new perspectives for the generation of universal donor cells for tissue transplantation. *J Gene Med* 6, 616-23.

Biswas, C., Zhang, Y., DeCastro, R., Guo, H., Nakamura, T., Kataoka, H. and Nabeshima, K. (1995) The human tumor cell-derived collagenase stimulatory factor (renamed EMMPRIN) is a member of the immunoglobulin superfamily. *Cancer Res* 55, 434-9.

- Bitto, E. and McKay, D.B. (2002) Crystallographic structure of SurA, a molecular chaperone that facilitates folding of outer membrane porins. *Structure* 10, 1489-98.
- Bitto, E. and McKay, D.B. (2003) The periplasmic molecular chaperone protein SurA binds a peptide motif that is characteristic of integral outer membrane proteins. *J Biol Chem* 278, 49316-22.
- Bloomfield, V.A. (1996) DNA condensation. *Curr Opin Struct Biol* 6, 334-41.
- Boldicke, T. (2007) Blocking translocation of cell surface molecules from the ER to the cell surface by intracellular antibodies targeted to the ER. *J Cell Mol Med* 11, 54-70.
- Boldicke, T., Weber, H., Mueller, P.P., Barleon, B. and Bernal, M. (2005) Novel highly efficient intrabody mediates complete inhibition of cell surface expression of the human vascular endothelial growth factor receptor-2 (VEGFR-2/KDR). *J Immunol Methods* 300, 146-59.
- Bukau, B. (1993) Regulation of the *Escherichia coli* heat-shock response. *Mol Microbiol* 9, 671-80.
- Bukau, B. and Horwich, A.L. (1998) The Hsp70 and Hsp60 chaperone machines. *Cell* 92, 351-66.
- Cao, T. and Heng, B.C. (2005) Intracellular antibodies (intrabodies) versus RNA interference for therapeutic applications. *Ann Clin Lab Sci* 35, 227-9.
- Carlson, J.R. (1988) A new means of inducibly inactivating a cellular protein. *Mol Cell Biol* 8, 2638-46.

- Cascales, E., Gavioli, M., Sturgis, J.N. and Lloubes, R. (2000) Proton motive force drives the interaction of the inner membrane TolA and outer membrane pal proteins in *Escherichia coli*. *Mol Microbiol* 38, 904-15.
- Casey, J.L., Coley, A.M. and Foley, M. (2008) Phage display of peptides in ligand selection for use in affinity chromatography. *Methods Mol Biol* 421, 111-24.
- Cattaneo, R. (2004) Four viruses, two bacteria, and one receptor: membrane cofactor protein (CD46) as pathogens' magnet. *J Virol* 78, 4385-8.
- Cesareni, G. (1992) Peptide display on filamentous phage capsids: A new powerful tool to study protein-ligand interaction. *FEBS Lett* 307, 66-70.
- Chen, X., Lin, J., Kanekura, T., Su, J., Lin, W., Xie, H., Wu, Y., Li, J., Chen, M. and Chang, J. (2006) A small interfering CD147-targeting RNA inhibited the proliferation, invasiveness, and metastatic activity of malignant melanoma. *Cancer Res* 66, 11323-30.
- Chen, X., Su, J., Chang, J., Kanekura, T., Li, J., Kuang, Y.H., Peng, S., Yang, F., Lu, H. and Zhang, J.L. (2008) Inhibition of CD147 Gene Expression via RNA Interference Reduces Tumor Cell Proliferation, Activation, Adhesion, and Migration Activity in the Human Jurkat T-Lymphoma Cell Line. *Cancer Invest*, 1.
- Chiampanichayakul, S., Peng-In, P., Khunkaewla, P., Stockinger, H. and Kasinrer, W. (2006) CD147 contains different bioactive epitopes involving the regulation of cell adhesion and lymphocyte activation. *Immunobiology* 211, 167-78.

- Choi, J.H. and Lee, S.Y. (2004) Secretory and extracellular production of recombinant proteins using *Escherichia coli*. *Appl Microbiol Biotechnol* 64, 625-35.
- Cohen, C.J., Shieh, J.T., Pickles, R.J., Okegawa, T., Hsieh, J.T. and Bergelson, J.M. (2001) The coxsackievirus and adenovirus receptor is a transmembrane component of the tight junction. *Proc Natl Acad Sci U S A* 98, 15191-6.
- Curtin, K.D., Meinertzhagen, I.A. and Wyman, R.J. (2005) Basigin (EMMPRIN/CD147) interacts with integrin to affect cellular architecture. *J Cell Sci* 118, 2649-60.
- Danese, P.N. and Silhavy, T.J. (1997) The sigma(E) and the Cpx signal transduction systems control the synthesis of periplasmic protein-folding enzymes in *Escherichia coli*. *Genes Dev* 11, 1183-93.
- Danese, P.N., Snyder, W.B., Cosma, C.L., Davis, L.J. and Silhavy, T.J. (1995) The Cpx two-component signal transduction pathway of *Escherichia coli* regulates transcription of the gene specifying the stress-inducible periplasmic protease, DegP. *Genes Dev* 9, 387-98.
- Darby, N.J. and Creighton, T.E. (1995) Catalytic mechanism of DsbA and its comparison with that of protein disulfide isomerase. *Biochemistry* 34, 3576-87.
- de Gier, J.W. and Luirink, J. (2001) Biogenesis of inner membrane proteins in *Escherichia coli*. *Mol Microbiol* 40, 314-22.
- de Keyzer, J., van der Does, C. and Driessen, A.J. (2003) The bacterial translocase: a dynamic protein channel complex. *Cell Mol Life Sci* 60, 2034-52.

- De Las Penas, A., Connolly, L. and Gross, C.A. (1997) The sigmaE-mediated response to extracytoplasmic stress in *Escherichia coli* is transduced by RseA and RseB, two negative regulators of sigmaE. *Mol Microbiol* 24, 373-85.
- Dehecchi, M.C., Melotti, P., Bonizzato, A., Santacatterina, M., Chilosi, M. and Cabrini, G. (2001) Heparan sulfate glycosaminoglycans are receptors sufficient to mediate the initial binding of adenovirus types 2 and 5. *J Virol* 75, 8772-80.
- Deng, L.W., Malik, P. and Perham, R.N. (1999) Interaction of the globular domains of pIII protein of filamentous bacteriophage fd with the F-pilus of *Escherichia coli*. *Virology* 253, 271-7.
- Deshane, J., Cabrera, G., Grim, J.E., Siegal, G.P., Pike, J., Alvarez, R.D. and Curiel, D.T. (1995a) Targeted eradication of ovarian cancer mediated by intracellular expression of anti-erbB-2 single-chain antibody. *Gynecol Oncol* 59, 8-14.
- Deshane, J., Siegal, G.P., Alvarez, R.D., Wang, M.H., Feng, M., Cabrera, G., Liu, T., Kay, M. and Curiel, D.T. (1995b) Targeted tumor killing via an intracellular antibody against erbB-2. *J Clin Invest* 96, 2980-9.
- Dobson, C.M. (2003) Protein folding and misfolding. *Nature* 426, 884-90.
- Doebis, C., Schu, S., Ladhoff, J., Busch, A., Beyer, F., Reiser, J., Nicosia, R.F., Broesel, S., Volk, H.D. and Seifert, M. (2006) An anti-major histocompatibility complex class I intrabody protects endothelial cells from an attack by immune mediators. *Cardiovasc Res* 72, 331-8.

Doerfler, W., Schubbert, R., Heller, H., Kammer, C., Hilger-Eversheim, K., Knoblauch, M. and Remus, R. (1997) Integration of foreign DNA and its consequences in mammalian systems. *Trends Biotechnol* 15, 297-301.

Driessen, A.J., Fekkes, P. and van der Wolk, J.P. (1998) The Sec system. *Curr Opin Microbiol* 1, 216-22.

Driskell, R.A. and Engelhardt, J.F. (2003) Current status of gene therapy for inherited lung diseases. *Annu Rev Physiol* 65, 585-612.

Emadi, S., Barkhordarian, H., Wang, M.S., Schulz, P. and Sierks, M.R. (2007) Isolation of a human single chain antibody fragment against oligomeric alpha-synuclein that inhibits aggregation and prevents alpha-synuclein-induced toxicity. *J Mol Biol* 368, 1132-44.

Erickson, J.W. and Gross, C.A. (1989) Identification of the sigma E subunit of *Escherichia coli* RNA polymerase: a second alternate sigma factor involved in high-temperature gene expression. *Genes Dev* 3, 1462-71.

Fadool, J.M. and Linser, P.J. (1993) 5A11 antigen is a cell recognition molecule which is involved in neuronal-gial interactions in avian neural retina. *Dev Dyn* 196, 252-62.

Fekkes, P., de Wit, J.G., van der Wolk, J.P., Kimsey, H.H., Kumamoto, C.A. and Driessen, A.J. (1998) Preprotein transfer to the *Escherichia coli* translocase requires the co-operative binding of SecB and the signal sequence to SecA. *Mol Microbiol* 29, 1179-90.

Fekkes, P. and Driessen, A.J. (1999) Protein targeting to the bacterial cytoplasmic membrane. *Microbiol Mol Biol Rev* 63, 161-73.

Fessler, S.P. and Young, C.S. (1998) Control of adenovirus early gene expression during the late phase of infection. *J Virol* 72, 4049-56.

Fleischli, C., Verhaagh, S., Havenga, M., Sirena, D., Schaffner, W., Cattaneo, R., Greber, U.F. and Hemmi, S. (2005) The distal short consensus repeats 1 and 2 of the membrane cofactor protein CD46 and their distance from the cell membrane determine productive entry of species B adenovirus serotype 35. *J Virol* 79, 10013-22.

Fossum, S., Mallett, S. and Barclay, A.N. (1991) The MRC OX-47 antigen is a member of the immunoglobulin superfamily with an unusual transmembrane sequence. *Eur J Immunol* 21, 671-9.

Frech, C., Wunderlich, M., Glockshuber, R. and Schmid, F.X. (1996) Preferential binding of an unfolded protein to DsbA. *EMBO J* 15, 392-98.

Fulford, W. and Model, P. (1984) Gene X of bacteriophage f1 is required for phage DNA synthesis. Mutagenesis of in-frame overlapping genes. *J Mol Biol* 178, 137-53.

Gabison, E.E., Mourah, S., Steinfels, E., Yan, L., Hoang-Xuan, T., Watsky, M.A., De Wever, B., Calvo, F., Mauviel, A. and Menashi, S. (2005) Differential expression of extracellular matrix metalloproteinase inducer (CD147) in normal and ulcerated corneas: role in epithelio-stromal interactions and matrix metalloproteinase induction. *Am J Pathol* 166, 209-19.

Gaggar, A., Shayakhmetov, D.M. and Lieber, A. (2003) CD46 is a cellular receptor for group B adenoviruses. *Nat Med* 9, 1408-12.

Gaggar, A., Shayakhmetov, D.M., Liszewski, M.K., Atkinson, J.P. and Lieber, A. (2005) Localization of regions in CD46 that interact with adenovirus. *J Virol* 79, 7503-13.

Gailus, V., Ramsperger, U., Johner, C., Kramer, H. and Rasched, I. (1994) The role of the adsorption complex in the termination of filamentous phage assembly. *Res Microbiol* 145, 699-709.

Gallagher, S.M., Castorino, J.J., Wang, D. and Philp, N.J. (2007) Monocarboxylate transporter 4 regulates maturation and trafficking of CD147 to the plasma membrane in the metastatic breast cancer cell line MDA-MB-231. *Cancer Res* 67, 4182-9.

Gamer, J., Multhaup, G., Tomoyasu, T., McCarty, J.S., Rudiger, S., Schonfeld, H.J., Schirra, C., Bujard, H. and Bukau, B. (1996) A cycle of binding and release of the DnaK, DnaJ and GrpE chaperones regulates activity of the Escherichia coli heat shock transcription factor sigma32. *EMBO J* 15, 607-17.

Glasgow, J.N., Bauerschmitz, G.J., Curiel, D.T. and Hemminki, A. (2004) Transductional and transcriptional targeting of adenovirus for clinical applications. *Curr Gene Ther* 4, 1-14.

Goenaga, A.L., Zhou, Y., Legay, C., Bougherara, H., Huang, L., Liu, B., Drummond, D.C., Kirpotin, D.B., Auclair, C., Marks, J.D. and Poul, M.A. (2007) Identification and characterization of tumor antigens by using antibody phage display and intrabody strategies. *Mol Immunol* 44, 3777-88.

Goncalves, M.A. (2005) Adeno-associated virus: from defective virus to effective vector. *Virol J* 2, 43.

- Goodrum, F.D. and Ornelles, D.A. (1999) Roles for the E4 orf6, orf3, and E1B 55-kilodalton proteins in cell cycle-independent adenovirus replication. *J Virol* 73, 7474-88.
- Greber, U.F. (2002) Signalling in viral entry. *Cell Mol Life Sci* 59, 608-26.
- Griesenbach, U., Geddes, D.M. and Alton, E.W. (2006) Gene therapy progress and prospects: cystic fibrosis. *Gene Ther* 13, 1061-7.
- Griffin, H., Elston, R., Jackson, D., Ansell, K., Coleman, M., Winter, G. and Doorbar, J. (2006) Inhibition of papillomavirus protein function in cervical cancer cells by intrabody targeting. *J Mol Biol* 355, 360-78.
- Guo, H., Zucker, S., Gordon, M.K., Toole, B.P. and Biswas, C. (1997) Stimulation of matrix metalloproteinase production by recombinant extracellular matrix metalloproteinase inducer from transfected Chinese hamster ovary cells. *J Biol Chem* 272, 24-7.
- Hacein-Bey-Abina, S., von Kalle, C., Schmidt, M., Le Deist, F., Wulffraat, N., McIntyre, E., Radford, I., Villeval, J.L., Fraser, C.C., Cavazzana-Calvo, M. and Fischer, A. (2003) A serious adverse event after successful gene therapy for X-linked severe combined immunodeficiency. *N Engl J Med* 348, 255-6.
- Halbert, D.N., Cutt, J.R. and Shenk, T. (1985) Adenovirus early region 4 encodes functions required for efficient DNA replication, late gene expression, and host cell shutoff. *J Virol* 56, 250-7.
- Hamilton, S., Odili, J., Wilson, G.D. and Kupsch, J.M. (2002) Reducing renal accumulation of single-chain Fv against melanoma-associated proteoglycan by coadministration of L-lysine. *Melanoma Res* 12, 373-9.

- Hammond, C. and Helenius, A. (1995) Quality control in the secretory pathway. *Curr Opin Cell Biol* 7, 523-9.
- Han, J., Sabbatini, P., Perez, D., Rao, L., Modha, D. and White, E. (1996) The E1B 19K protein blocks apoptosis by interacting with and inhibiting the p53-inducible and death-promoting Bax protein. *Genes Dev* 10, 461-77.
- Hao, J., Serohijos, A.W., Newton, G., Tassone, G., Wang, Z., Sgroi, D.C., Dokholyan, N.V. and Bacion, J.P. (2008) Identification and rational redesign of peptide ligands to CRIP1, a novel biomarker for cancers. *PLoS Comput Biol* 4, e1000138.
- Harlow, E., Lane, D. (1998) *Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, NY.
- Harms, N., Koningstein, G., Dontje, W., Muller, M., Oudega, B., Luirink, J. and de Cock, H. (2001) The early interaction of the outer membrane protein phoe with the periplasmic chaperone Skp occurs at the cytoplasmic membrane. *J Biol Chem* 276, 18804-11.
- Hartl, F.U. and Hayer-Hartl, M. (2002) Molecular chaperones in the cytosol: from nascent chain to folded protein. *Science* 295, 1852-8.
- Hay, R.T., Freeman, A., Leith, I., Monaghan, A. and Webster, A. (1995) Molecular interactions during adenovirus DNA replication. *Curr Top Microbiol Immunol* 199 (Pt 2), 31-48.
- He, T.C., Zhou, S., da Costa, L.T., Yu, J., Kinzler, K.W. and Vogelstein, B. (1998) A simplified system for generating recombinant adenoviruses. *Proc Natl Acad Sci U S A* 95, 2509-14.

Hearing, P., Samulski, R.J., Wishart, W.L. and Shenk, T. (1987) Identification of a repeated sequence element required for efficient encapsidation of the adenovirus type 5 chromosome. *J Virol* 61, 2555-8.

Hell, R.C., Amim, P., de Andrade, H.M., de Avila, R.A., Felicori, L., Oliveira, A.G., Oliveira, C.A., Nascimento, E., Tavares, C.A., Granier, C. and Chavez-Olortegui, C. (2009) Immunodiagnosis of human neurocysticercosis using a synthetic peptide selected by phage-display. *Clin Immunol.*

Heller, M., von der Ohe, M., Kleene, R., Mohajeri, M.H. and Schachner, M. (2003) The immunoglobulin-superfamily molecule basigin is a binding protein for oligomannosidic carbohydrates: an anti-idiotypic approach. *J Neurochem* 84, 557-65.

Heng, B.C., Kemeny, D.M., Liu, H. and Cao, T. (2005) Potential applications of intracellular antibodies (intrabodies) in stem cell therapeutics. *J Cell Mol Med* 9, 191-5.

Heppner, K.J., Matrisian, L.M., Jensen, R.A. and Rodgers, W.H. (1996) Expression of most matrix metalloproteinase family members in breast cancer represents a tumor-induced host response. *Am J Pathol* 149, 273-82.

Hertveldt, K., Belien, T. and Volckaert, G. (2009) General M13 phage display: M13 phage display in identification and characterization of protein-protein interactions. *Methods Mol Biol* 502, 321-39.

Hiniker, A. and Bardwell, J.C. (2003) Disulfide bond isomerization in prokaryotes. *Biochemistry* 42, 1179-85.

Holliger, P. and Hudson, P.J. (2005) Engineered antibody fragments and the rise of single domains. *Nat Biotechnol* 23, 1126-36.

Holliger, P. and Riechmann, L. (1997) A conserved infection pathway for filamentous bacteriophages is suggested by the structure of the membrane penetration domain of the minor coat protein g3p from phage fd. *Structure* 5, 265-75.

Holmgren, A. (1989) Thioredoxin and glutaredoxin systems. *J Biol Chem* 264, 13963-6.

Hong, S.S., Karayan, L., Tournier, J., Curiel, D.T. and Boulanger, P.A. (1997) Adenovirus type 5 fiber knob binds to MHC class I alpha2 domain at the surface of human epithelial and B lymphoblastoid cells. *EMBO J* 16, 2294-306.

Hoogenboom, H.R., de Bruine, A.P., Hufton, S.E., Hoet, R.M., Arends, J.W. and Roovers, R.C. (1998) Antibody phage display technology and its applications. *Immunotechnology* 4, 1-20.

Hoyer, W., Gronwall, C., Jonsson, A., Stahl, S. and Hard, T. (2008) Stabilization of a beta-hairpin in monomeric Alzheimer's amyloid-beta peptide inhibits amyloid formation. *Proc Natl Acad Sci U S A* 105, 5099-104.

http://www.alt.med-rz.uni-sb.de/med_fak/biochemie/MartinJ/Eninformation.html

http://www.athenaes.com/tech_brief_ACESyebf.php

Huang, S., Kamata, T., Takada, Y., Ruggeri, Z.M. and Nemerow, G.R. (1996)

Adenovirus interaction with distinct integrins mediates separate events in cell entry and gene delivery to hematopoietic cells. *J Virol* 70, 4502-8.

Hudson, P.J. (1998) Recombinant antibody fragments. *Curr Opin Biotechnol* 9, 395-402.

Igakura, T., Kadomatsu, K., Taguchi, O., Muramatsu, H., Kaname, T., Miyauchi, T., Yamamura, K., Arimura, K. and Muramatsu, T. (1996) Roles of basigin, a member of the immunoglobulin superfamily, in behavior as to an irritating odor, lymphocyte response, and blood-brain barrier. *Biochem Biophys Res Commun* 224, 33-6.

Intasai, N., Arooncharus, P., Kasinrerak, W. and Tayapiwatana, C. (2003) Construction of high-density display of CD147 ectodomain on VCSM13 phage via gpVIII: effects of temperature, IPTG, and helper phage infection-period. *Protein Expr Purif* 32, 323-31.

Intasai, N., Mai, S., Kasinrerak, W. and Tayapiwatana, C. (2006) Binding of multivalent CD147 phage induces apoptosis of U937 cells. *Int Immunol* 18, 1159-69.

Jager, S., Jahnke, A., Wilmes, T., Adebahr, S., Vogtle, F.N., Delima-Hahn, E., Pfeifer, D., Berg, T., Lubbert, M. and Trepel, M. (2007) Leukemia-targeting ligands isolated from phage-display peptide libraries. *Leukemia* 21, 411-20.

Jannot, C.B., Beerli, R.R., Mason, S., Gullick, W.J. and Hynes, N.E. (1996) Intracellular expression of a single-chain antibody directed to the EGFR leads to growth inhibition of tumor cells. *Oncogene* 13, 275-82.

Jendreyko, N., Popkov, M., Beerli, R.R., Chung, J., McGavern, D.B., Rader, C. and Barbas, C.F., 3rd. (2003) Intradiabodies, bispecific, tetravalent antibodies for

the simultaneous functional knockout of two cell surface receptors. *J Biol Chem* 278, 47812-9.

Jendreyko, N., Popkov, M., Rader, C. and Barbas, C.F., 3rd. (2005) Phenotypic knockout of VEGF-R2 and Tie-2 with an intradiabody reduces tumor growth and angiogenesis in vivo. *Proc Natl Acad Sci U S A* 102, 8293-8.

Jia, L., Xu, H., Zhao, Y., Jiang, L., Yu, J. and Zhang, J. (2008) Expression of CD147 mediates tumor cells invasion and multidrug resistance in hepatocellular carcinoma. *Cancer Invest* 26, 977-83.

Jones, C.H., Danese, P.N., Pinkner, J.S., Silhavy, T.J. and Hultgren, S.J. (1997) The chaperone-assisted membrane release and folding pathway is sensed by two signal transduction systems. *EMBO J* 16, 6394-406.

Jung, D., Neron, S., Drouin, M. and Jacques, A. (2005) Efficient gene transfer into normal human B lymphocytes with the chimeric adenoviral vector Ad5/F35. *J Immunol Methods* 304, 78-87.

Kabat, E.A., Wu, T.T. and Bilofsky, H. (1976a) Attempts to locate residues in complementarity-determining regions of antibody combining sites that make contact with antigen. *Proc Natl Acad Sci U S A* 73, 617-9.

Kabat, E.A., Wu, T.T. and Bilofsky, H. (1976b) Some correlations between specificity and sequence of the first complementarity-determining segments of human kappa light chains. *Proc Natl Acad Sci U S A* 73, 4471-3.

Kabat, E.A., Wu, T.T., Bilofsky, H. and National Institutes of Health (U.S.). Division of Research Resources. (1976c) Variable regions of immunoglobulin chains :

tabulations and analyses of amino acid sequences. Medical Computer Systems, Cambridge, Mass.

Kannabiran, C., Morris, G.F. and Mathews, M.B. (1999) Dual action of the adenovirus E1A 243R oncoprotein on the human proliferating cell nuclear antigen promoter: repression of transcriptional activation by p53. *Oncogene* 18, 7825-33.

Kaplan, J.M., Armentano, D., Scaria, A., Woodworth, L.A., Pennington, S.E., Wadsworth, S.C., Smith, A.E. and Gregory, R.J. (1999) Novel role for E4 region genes in protection of adenovirus vectors from lysis by cytotoxic T lymphocytes. *J Virol* 73, 4489-92.

Karlsson, F., Borrebaeck, C.A., Nilsson, N. and Malmberg-Hager, A.C. (2003) The mechanism of bacterial infection by filamentous phages involves molecular interactions between TolA and phage protein 3 domains. *J Bacteriol* 185, 2628-34.

Kasinrerk, W., Fiebiger, E., Stefanova, I., Baumruker, T., Knapp, W. and Stockinger, H. (1992) Human leukocyte activation antigen M6, a member of the Ig superfamily, is the species homologue of rat OX-47, mouse basigin, and chicken HT7 molecule. *J Immunol* 149, 847-54.

Kasinrerk, W., Tokrasinwit, N. and Phunpae, P. (1999) CD147 monoclonal antibodies induce homotypic cell aggregation of monocytic cell line U937 via LFA-1/ICAM-1 pathway. *Immunology* 96, 184-92.

Kato-Takagaki, K., Mizukoshi, Y., Yoshizawa, Y., Akazawa, D., Torii, Y., Ono, K., Tanimura, R., Shimada, I. and Takahashi, H. (2009) Structural and interaction

analysis of glycoprotein VI-binding peptide selected from phage display library. *J Biol Chem*.

Kaufman, R.J. (2002) Orchestrating the unfolded protein response in health and disease. *J Clin Invest* 110, 1389-98.

Kay, M.A., Glorioso, J.C. and Naldini, L. (2001) Viral vectors for gene therapy: the art of turning infectious agents into vehicles of therapeutics. *Nat Med* 7, 33-40.

Keiler, K.C., Silber, K.R., Downard, K.M., Papayannopoulos, I.A., Biemann, K. and Sauer, R.T. (1995) C-terminal specific protein degradation: activity and substrate specificity of the Tsp protease. *Protein Sci* 4, 1507-15.

Khokhlova, O.V. and Nesmeianova, M.A. (2003) [Interaction of SecB and SecA with the N-terminal region of mature alkaline phosphatase on its secretion in *Escherichia coli*]. *Mol Biol (Mosk)* 37, 712-8.

Khunkeawla, P., Moonsom, S., Staffler, G., Kongtawelert, P. and Kasinrerak, W. (2001) Engagement of CD147 molecule-induced cell aggregation through the activation of protein kinases and reorganization of the cytoskeleton. *Immunobiology* 203, 659-69.

Kiefhaber, T., Rudolph, R., Kohler, H.H. and Buchner, J. (1991) Protein aggregation *in vitro* and *in vivo*: a quantitative model of the kinetic competition between folding and aggregation. *Biotechnology (N Y)* 9, 825-9.

Kim, D.J., Chung, J.H., Ryu, Y.S., Rhim, J.H., Kim, C.W., Suh, Y. and Chung, H.K. (2002) Production and characterisation of a recombinant scFv reactive with human gastrointestinal carcinomas. *Br J Cancer* 87, 405-13.

- Kim, J., Chen, C.P. and Rice, K.G. (2005) The proteasome metabolizes peptide-mediated nonviral gene delivery systems. *Gene Ther* 12, 1581-90.
- Kioi, M., Seetharam, S. and Puri, R.K. (2008) Targeting IL-13R α 2-positive cancer with a novel recombinant immunotoxin composed of a single-chain antibody and mutated *Pseudomonas* exotoxin. *Mol Cancer Ther* 7, 1579-87.
- Kirsch, A.H., Diaz, L.A., Jr., Bonish, B., Antony, P.A. and Fox, D.A. (1997) The pattern of expression of CD147/neurothelin during human T-cell ontogeny as defined by the monoclonal antibody 8D6. *Tissue Antigens* 50, 147-52.
- Kleerebezem, M., Crielaard, W. and Tommassen, J. (1996) Involvement of stress protein PspA (phage shock protein A) of *Escherichia coli* in maintenance of the protonmotive force under stress conditions. *EMBO J* 15, 162-71.
- Kobayashi, T., Kishigami, S., Sone, M., Inokuchi, H., Mogi, T. and Ito, K. (1997) Respiratory chain is required to maintain oxidized states of the DsbA-DsbB disulfide bond formation system in aerobically growing *Escherichia coli* cells. *Proc Natl Acad Sci U S A* 94, 11857-62.
- Koch, C., Staffler, G., Huttinger, R., Hilgert, I., Prager, E., Cerny, J., Steinlein, P., Majdic, O., Horejsi, V. and Stockinger, H. (1999) T cell activation-associated epitopes of CD147 in regulation of the T cell response, and their definition by antibody affinity and antigen density. *Int Immunol* 11, 777-86.
- Kontermann, R., Dübel, S. (2001) *Antibody engineering*. Springer.
- Kontermann, R.E. (2004) Intrabodies as therapeutic agents. *Methods* 34, 163-70.
- Korn, T., Nettelbeck, D.M., Volkel, T., Muller, R. and Kontermann, R.E. (2004) Recombinant bispecific antibodies for the targeting of adenoviruses to CEA-

expressing tumour cells: a comparative analysis of bacterially expressed single-chain diabody and tandem scFv. *J Gene Med* 6, 642-51.

Kwong, K.Y., Baskar, S., Zhang, H., Mackall, C.L. and Rader, C. (2008) Generation, affinity maturation, and characterization of a human anti-human NKG2D monoclonal antibody with dual antagonistic and agonistic activity. *J Mol Biol* 384, 1143-56.

Lai, C.M., Lai, Y.K. and Rakoczy, P.E. (2002) Adenovirus and adeno-associated virus vectors. *DNA Cell Biol* 21, 895-913.

Lazdunski, C.J., Bouveret, E., Rigal, A., Journet, L., Llobes, R. and Benedetti, H. (1998) Colicin import into *Escherichia coli* cells. *J Bacteriol* 180, 4993-5002.

Lazzaroni, J.C., Germon, P., Ray, M.C. and Vianney, A. (1999) The Tol proteins of *Escherichia coli* and their involvement in the uptake of biomolecules and outer membrane stability. *FEMS Microbiol Lett* 177, 191-7.

Lewin, B., Cassimeris, L., Lingappa, V.F. and Plopper, G. (2007) *Cells*. Jones and Bartlett Publishers Ontario.

Li, E., Brown, S.L., Stupack, D.G., Puente, X.S., Cheresch, D.A. and Nemerow, G.R. (2001) Integrin alpha(v)beta1 is an adenovirus coreceptor. *J Virol* 75, 5405-9.

Li, H., Cybulsky, M.I., Gimbrone, M.A., Jr. and Libby, P. (1993) An atherogenic diet rapidly induces VCAM-1, a cytokine-regulatable mononuclear leukocyte adhesion molecule, in rabbit aortic endothelium. *Arterioscler Thromb* 13, 197-204.

Li, Q.Q., Wang, W.J., Xu, J.D., Cao, X.X., Chen, Q., Yang, J.M. and Xu, Z.D. (2007) Up-regulation of CD147 and matrix metalloproteinase-2, -9 induced by P-

glycoprotein substrates in multidrug resistant breast cancer cells. *Cancer Sci* 98, 1767-74.

Liberek, K., Galitski, T.P., Zylicz, M. and Georgopoulos, C. (1992) The DnaK chaperone modulates the heat shock response of *Escherichia coli* by binding to the sigma 32 transcription factor. *Proc Natl Acad Sci U S A* 89, 3516-20.

Lim, M., Martinez, T., Jablons, D., Cameron, R., Guo, H., Toole, B., Li, J.D. and Basbaum, C. (1998) Tumor-derived EMMPRIN (extracellular matrix metalloproteinase inducer) stimulates collagenase transcription through MAPK p38. *FEBS Lett* 441, 88-92.

Liu, D. and Knapp, J.E. (2001) Hydrodynamics-based gene delivery. *Curr Opin Mol Ther* 3, 192-7.

Liu, F., Lu, Q., Ye, X., Fang, C., Zhao, Y., Liang, M., Hu, F., Lieber, A. and Chen, H.Z. (2008) Cancer gene therapy of adenovirus-mediated anti-4-1BB scFv in immunocompetent mice. *Cancer Biol Ther* 7, 448-53.

Liu, M.Y., Han, W., Ding, Y.L., Zhou, T.H., Tian, R.Y., Yang, S.L., Liu, H. and Gong, Y. (2005) Generation and characterization of C305, a murine neutralizing scFv antibody that can inhibit BlyS binding to its receptor BCMA. *Acta Biochim Biophys Sin (Shanghai)* 37, 415-20.

Lobato, M.N. and Rabbitts, T.H. (2004) Intracellular antibodies as specific reagents for functional ablation: future therapeutic molecules. *Curr Mol Med* 4, 519-28.

Lowman, H.B. (1997) Bacteriophage display and discovery of peptide leads for drug development. *Annu Rev Biophys Biomol Struct* 26, 401-24.

- Lowman, H.B., Bass, S.H., Simpson, N. and Wells, J.A. (1991) Selecting high-affinity binding proteins by monovalent phage display. *Biochemistry* 30, 10832-8.
- Lowman, H.B. and Wells, J.A. (1993) Affinity maturation of human growth hormone by monovalent phage display. *J Mol Biol* 234, 564-78.
- Luo, J., Deng, Z.L., Luo, X., Tang, N., Song, W.X., Chen, J., Sharff, K.A., Luu, H.H., Haydon, R.C., Kinzler, K.W., Vogelstein, B. and He, T.C. (2007) A protocol for rapid generation of recombinant adenoviruses using the AdEasy system. *Nat Protoc* 2, 1236-47.
- Lutz, P., Rosa-Calatrava, M. and Kedinger, C. (1997) The product of the adenovirus intermediate gene IX is a transcriptional activator. *J Virol* 71, 5102-9.
- Lynch, S.M., Zhou, C. and Messer, A. (2008) An scFv intrabody against the nonamyloid component of alpha-synuclein reduces intracellular aggregation and toxicity. *J Mol Biol* 377, 136-47.
- Majhen, D. and Ambriovic-Ristov, A. (2006) Adenoviral vectors--how to use them in cancer gene therapy? *Virus Res* 119, 121-33.
- Majmudar, G., Nelson, B.R., Jensen, T.C. and Johnson, T.M. (1994) Increased expression of matrix metalloproteinase-3 (stromelysin-1) in cultured fibroblasts and basal cell carcinomas of nevoid basal cell carcinoma syndrome. *Mol Carcinog* 11, 29-33.
- Makrides, S.C. (1996) Strategies for achieving high-level expression of genes in *Escherichia coli*. *Microbiol Rev* 60, 512-38.

- Mao, S., Gao, C., Lo, C.H., Wirsching, P., Wong, C.H. and Janda, K.D. (1999) Phage-display library selection of high-affinity human single-chain antibodies to tumor-associated carbohydrate antigens sialyl Lewisx and Lewisx. *Proc Natl Acad Sci U S A* 96, 6953-8.
- Marasco, W.A., Chen, S., Richardson, J.H., Ramstedt, U. and Jones, S.D. (1998) Intracellular antibodies against HIV-1 envelope protein for AIDS gene therapy. *Hum Gene Ther* 9, 1627-42.
- Marasco, W.A., Haseltine, W.A. and Chen, S.Y. (1993) Design, intracellular expression, and activity of a human anti-human immunodeficiency virus type 1 gp120 single-chain antibody. *Proc Natl Acad Sci U S A* 90, 7889-93.
- Martin, J.L., Bardwell, J.C. and Kuriyan, J. (1993) Crystal structure of the DsbA protein required for disulphide bond formation *in vivo*. *Nature* 365, 464-8.
- McCafferty, J., Griffiths, A.D., Winter, G. and Chiswell, D.J. (1990) Phage antibodies: filamentous phage displaying antibody variable domains. *Nature* 348, 552-4.
- Menashi, S., Serova, M., Ma, L., Vignot, S., Mourah, S. and Calvo, F. (2003) Regulation of extracellular matrix metalloproteinase inducer and matrix metalloproteinase expression by amphiregulin in transformed human breast epithelial cells. *Cancer Res* 63, 7575-80.
- Mergulhao, F.J., Summers, D.K. and Monteiro, G.A. (2005) Recombinant protein secretion in *Escherichia coli*. *Biotechnol Adv* 23, 177-202.

- Messer, A. and McLear, J. (2006) The therapeutic potential of intrabodies in neurologic disorders: focus on Huntington and Parkinson diseases. *BioDrugs* 20, 327-33.
- Mhashilkar, A.M., Bagley, J., Chen, S.Y., Szilvay, A.M., Helland, D.G. and Marasco, W.A. (1995) Inhibition of HIV-1 Tat-mediated LTR transactivation and HIV-1 infection by anti-Tat single chain intrabodies. *EMBO J* 14, 1542-51.
- Mhashilkar, A.M., Doebis, C., Seifert, M., Busch, A., Zani, C., Soo Hoo, J., Nagy, M., Ritter, T., Volk, H.D. and Marasco, W.A. (2002) Intrabody-mediated phenotypic knockout of major histocompatibility complex class I expression in human and monkey cell lines and in primary human keratinocytes. *Gene Ther* 9, 307-19.
- Millimaggi, D., Mari, M., D'Ascenzo, S., Carosa, E., Jannini, E.A., Zucker, S., Carta, G., Pavan, A. and Dolo, V. (2007) Tumor vesicle-associated CD147 modulates the angiogenic capability of endothelial cells. *Neoplasia* 9, 349-57.
- Missiakas, D., Georgopoulos, C. and Raina, S. (1993) Identification and characterization of the *Escherichia coli* gene dsbB, whose product is involved in the formation of disulfide bonds *in vivo*. *Proc Natl Acad Sci U S A* 90, 7084-8.
- Missiakas, D. and Raina, S. (1997) Protein misfolding in the cell envelope of *Escherichia coli*: new signaling pathways. *Trends Biochem Sci* 22, 59-63.
- Missiakas, D., Schwager, F., Betton, J.M., Georgopoulos, C. and Raina, S. (1996) Identification and characterization of HsIV HsIU (ClpQ ClpY) proteins

involved in overall proteolysis of misfolded proteins in *Escherichia coli*.
EMBO J 15, 6899-909.

Miyauchi, T., Kanekura, T., Yamaoka, A., Ozawa, M., Miyazawa, S. and Muramatsu, T. (1990) Basigin, a new, broadly distributed member of the immunoglobulin superfamily, has strong homology with both the immunoglobulin V domain and the beta-chain of major histocompatibility complex class II antigen. J Biochem 107, 316-23.

Miyauchi, T., Masuzawa, Y. and Muramatsu, T. (1991) The basigin group of the immunoglobulin superfamily: complete conservation of a segment in and around transmembrane domains of human and mouse basigin and chicken HT7 antigen. J Biochem 110, 770-4.

Mizuguchi, H. and Hayakawa, T. (2002) Adenovirus vectors containing chimeric type 5 and type 35 fiber proteins exhibit altered and expanded tropism and increase the size limit of foreign genes. Gene 285, 69-77.

Mizuguchi, H. and Hayakawa, T. (2004) Targeted adenovirus vectors. Hum Gene Ther 15, 1034-44.

Mizuguchi, H., Kay, M.A. and Hayakawa, T. (2001) Approaches for generating recombinant adenovirus vectors. Adv Drug Deliv Rev 52, 165-76.

Mohrluder, J., Stangler, T., Hoffmann, Y., Wiesehan, K., Mataruga, A. and Willbold, D. (2007) Identification of calreticulin as a ligand of GABARAP by phage display screening of a peptide library. FEBS J 274, 5543-55.

Mori, H. and Ito, K. (2001) The Sec protein-translocation pathway. Trends Microbiol 9, 494-500.

Mori, H. and Ito, K. (2003) Biochemical characterization of a mutationally altered protein translocase: proton motive force stimulation of the initiation phase of translocation. *J Bacteriol* 185, 405-12.

Mullen, L.M., Nair, S.P., Ward, J.M., Rycroft, A.N. and Henderson, B. (2006) Phage display in the study of infectious diseases. *Trends Microbiol* 14, 141-7.

Munro, S. and Pelham, H.R. (1987) A C-terminal signal prevents secretion of luminal ER proteins. *Cell* 48, 899-907.

Muramatsu, T. and Miyauchi, T. (2003) Basigin (CD147): a multifunctional transmembrane protein involved in reproduction, neural function, inflammation and tumor invasion. *Histol Histopathol* 18, 981-7.

Muraoka, K., Nabeshima, K., Murayama, T., Biswas, C. and Kono, M. (1993) Enhanced expression of a tumor-cell-derived collagenase-stimulatory factor in urothelial carcinoma: its usefulness as a tumor marker for bladder cancers. *Int J Cancer* 55, 19-26.

Nabeshima, K., Iwasaki, H., Koga, K., Hojo, H., Suzumiya, J. and Kikuchi, M.

(2006a) Emmprin (basigin/CD147): matrix metalloproteinase modulator and multifunctional cell recognition molecule that plays a critical role in cancer progression. *Pathol Int* 56, 359-67.

Nabeshima, K., Iwasaki, H., Nishio, J., Koga, K., Shishime, M. and Kikuchi, M.

(2006b) Expression of emmprin and matrix metalloproteinases (MMPs) in peripheral nerve sheath tumors: emmprin and membrane-type (MT)1-MMP expressions are associated with malignant potential. *Anticancer Res* 26, 1359-67.

- Nam, C.H., Lobato, M.N., Appert, A., Drynan, L.F., Tanaka, T. and Rabbitts, T.H. (2008) An antibody inhibitor of the LMO2-protein complex blocks its normal and tumorigenic functions. *Oncogene*.
- Neering, S.J., Hardy, S.F., Minamoto, D., Spratt, S.K. and Jordan, C.T. (1996) Transduction of primitive human hematopoietic cells with recombinant adenovirus vectors. *Blood* 88, 1147-55.
- Nehme, C.L., Cesario, M.M., Myles, D.G., Koppel, D.E. and Bartles, J.R. (1993) Breaching the diffusion barrier that compartmentalizes the transmembrane glycoprotein CE9 to the posterior-tail plasma membrane domain of the rat spermatozoon. *J Cell Biol* 120, 687-94.
- Neri, D., Petrucci, H. and Roncucci, G. (1995) Engineering recombinant antibodies for immunotherapy. *Cell Biophys* 27, 47-61.
- Nevins, J.R., DeGregori, J., Jakoi, L. and Leone, G. (1997) Functional analysis of E2F transcription factor. *Methods Enzymol* 283, 205-19.
- Nilsson, M., Ljungberg, J., Richter, J., Kiefer, T., Magnusson, M., Lieber, A., Widegren, B., Karlsson, S. and Fan, X. (2004) Development of an adenoviral vector system with adenovirus serotype 35 tropism; efficient transient gene transfer into primary malignant hematopoietic cells. *J Gene Med* 6, 631-41.
- Nishiyama, K., Fukuda, A., Morita, K. and Tokuda, H. (1999) Membrane deinsertion of SecA underlying proton motive force-dependent stimulation of protein translocation. *EMBO J* 18, 1049-58.

- Nissim, A., Hoogenboom, H.R., Tomlinson, I.M., Flynn, G., Midgley, C., Lane, D. and Winter, G. (1994) Antibody fragments from a 'single pot' phage display library as immunochemical reagents. *EMBO J* 13, 692-8.
- Olafsen, T., Tan, G.J., Cheung, C.W., Yazaki, P.J., Park, J.M., Shively, J.E., Williams, L.E., Raubitschek, A.A., Press, M.F. and Wu, A.M. (2004) Characterization of engineered anti-p185HER-2 (scFv-CH3)₂ antibody fragments (minibodies) for tumor targeting. *Protein Eng Des Sel* 17, 315-23.
- Paetzel, M., Karla, A., Strynadka, N.C. and Dalbey, R.E. (2002) Signal peptidases. *Chem Rev* 102, 4549-80.
- Pantoliano, M.W., Bird, R.E., Johnson, S., Asel, E.D., Dodd, S.W., Wood, J.F. and Hardman, K.D. (1991) Conformational stability, folding, and ligand-binding affinity of single-chain Fv immunoglobulin fragments expressed in *Escherichia coli*. *Biochemistry* 30, 10117-25.
- Pavoni, E., Flego, M., Dupuis, M.L., Barca, S., Petronzelli, F., Anastasi, A.M., D'Alessio, V., Pelliccia, A., Vaccaro, P., Monteriu, G., Ascione, A., De Santis, R., Felici, F., Cianfriglia, M. and Minenkova, O. (2006) Selection, affinity maturation, and characterization of a human scFv antibody against CEA protein. *BMC Cancer* 6, 41.
- Peng, J.L., Wu, S., Zhao, X.P., Wang, M., Li, W.H., Shen, X., Liu, J., Lei, P., Zhu, H.F. and Shen, G.X. (2007) Downregulation of transferrin receptor surface expression by intracellular antibody. *Biochem Biophys Res Commun* 354, 864-71.

- Pingmuang, P. (2008) Construction of adenovirus expression systems for comparison of single chain Fv production in cytoplasmic compartment and endoplasmic reticulum and as a secreting molecule. Department of Medical Technology, Faculty of Associated Medical Sciences, Master of Science. Chiang Mai University.
- Pogliano, J., Lynch, A.S., Belin, D., Lin, E.C. and Beckwith, J. (1997) Regulation of *Escherichia coli* cell envelope proteins involved in protein folding and degradation by the Cpx two-component system. *Genes Dev* 11, 1169-82.
- Poller, W., Schneider-Rasp, S., Liebert, U., Merklein, F., Thalheimer, P., Haack, A., Schwaab, R., Schmitt, C. and Brackmann, H.H. (1996) Stabilization of transgene expression by incorporation of E3 region genes into an adenoviral factor IX vector and by transient anti-CD4 treatment of the host. *Gene Ther* 3, 521-30.
- Popkov, M., Jendreyko, N., McGavern, D.B., Rader, C. and Barbas, C.F., 3rd. (2005) Targeting tumor angiogenesis with adenovirus-delivered anti-Tie-2 intrabody. *Cancer Res* 65, 972-81.
- Pushkarsky, T., Yurchenko, V., Vanpouille, C., Brichacek, B., Vaisman, I., Hatakeyama, S., Nakayama, K.I., Sherry, B. and Bukrinsky, M.I. (2005) Cell surface expression of CD147/EMMPRN is regulated by cyclophilin 60. *J Biol Chem* 280, 27866-71.
- Quemener, C., Gabison, E.E., Naimi, B., Lescaille, G., Bougatef, F., Podgorniak, M.P., Labarchede, G., Lebbe, C., Calvo, F., Menashi, S. and Mourah, S. (2007) Extracellular matrix metalloproteinase inducer up-regulates the

urokinase-type plasminogen activator system promoting tumor cell invasion. *Cancer Res* 67, 9-15.

Raina, S., Missiakas, D. and Georgopoulos, C. (1995) The *rpoE* gene encoding the sigma E (sigma 24) heat shock sigma factor of *Escherichia coli*. *EMBO J* 14, 1043-55.

Rainov, N.G. and Ren, H. (2003) Clinical trials with retrovirus mediated gene therapy--what have we learned? *J Neurooncol* 65, 227-36.

Rao, L., Perez, D. and White, E. (1996) Lamin proteolysis facilitates nuclear events during apoptosis. *J Cell Biol* 135, 1441-55.

Rea, D., Havenga, M.J., van Den Assem, M., Suttmuller, R.P., Lemckert, A., Hoeben, R.C., Bout, A., Melief, C.J. and Offringa, R. (2001) Highly efficient transduction of human monocyte-derived dendritic cells with subgroup B fiber-modified adenovirus vectors enhances transgene-encoded antigen presentation to cytotoxic T cells. *J Immunol* 166, 5236-44.

Rebel, V.I., Hartnett, S., Denham, J., Chan, M., Finberg, R. and Sieff, C.A. (2000) Maturation and lineage-specific expression of the coxsackie and adenovirus receptor in hematopoietic cells. *Stem Cells* 18, 176-82.

Reimers, N., Zafrakas, K., Assmann, V., Egen, C., Riethdorf, L., Riethdorf, S., Berger, J., Ebel, S., Janicke, F., Sauter, G. and Pantel, K. (2004) Expression of extracellular matrix metalloproteases inducer on micrometastatic and primary mammary carcinoma cells. *Clin Cancer Res* 10, 3422-8.

- Reinman, M., Jantti, J., Alfthan, K., Keranen, S., Soderlund, H. and Takkinen, K. (2003) Functional inactivation of the conserved Sem1p in yeast by intrabodies. *Yeast* 20, 1071-84.
- Ren, Z. and Black, L.W. (1998) Phage T4 SOC and HOC display of biologically active, full-length proteins on the viral capsid. *Gene* 215, 439-44.
- Richardson, J.H. and Marasco, W.A. (1995) Intracellular antibodies: development and therapeutic potential. *Trends Biotechnol* 13, 306-10.
- Riethdorf, S., Reimers, N., Assmann, V., Kornfeld, J.W., Terracciano, L., Sauter, G. and Pantel, K. (2006) High incidence of EMMPRIN expression in human tumors. *Int J Cancer* 119, 1800-10.
- Rietsch, A., Belin, D., Martin, N. and Beckwith, J. (1996) An *in vivo* pathway for disulfide bond isomerization in *Escherichia coli*. *Proc Natl Acad Sci U S A* 93, 13048-53.
- Robbins, P.D., Tahara, H. and Ghivizzani, S.C. (1998) Viral vectors for gene therapy. *Trends Biotechnol* 16, 35-40.
- Roberts, B.L., Markland, W., Ley, A.C., Kent, R.B., White, D.W., Guterman, S.K. and Ladner, R.C. (1992) Directed evolution of a protein: selection of potent neutrophil elastase inhibitors displayed on M13 fusion phage. *Proc Natl Acad Sci U S A* 89, 2429-33.
- Robles, Y., Gonzalez, E., Govezensky, T., Mungia, M.E., Acero, G., Bobes, R.J., Gevorkian, G. and Manoutcharian, K. (2005) Isolation of the *Taenia crassiceps* antigens from a phage display cDNA library and evaluation of their use for diagnosis of neurocysticercosis. *Clin Immunol* 116, 265-70.

Rothlisberger, D., Honegger, A. and Pluckthun, A. (2005) Domain interactions in the Fab fragment: a comparative evaluation of the single-chain Fv and Fab format engineered with variable domains of different stability. *J Mol Biol* 347, 773-89.

Russel, M., Linderoth, N.A. and Sali, A. (1997) Filamentous phage assembly: variation on a protein export theme. *Gene* 192, 23-32.

Russel, M., Whirlow, H., Sun, T.P. and Webster, R.E. (1988) Low-frequency infection of F- bacteria by transducing particles of filamentous bacteriophages. *J Bacteriol* 170, 5312-6.

Russell, W.C. (2000) Update on adenovirus and its vectors. *J Gen Virol* 81, 2573-604.

Sakurai, F., Mizuguchi, H., Yamaguchi, T. and Hayakawa, T. (2003) Characterization of *in vitro* and *in vivo* gene transfer properties of adenovirus serotype 35 vector. *Mol Ther* 8, 813-21.

Sakurai, F., Murakami, S., Kawabata, K., Okada, N., Yamamoto, A., Seya, T., Hayakawa, T. and Mizuguchi, H. (2006) The short consensus repeats 1 and 2, not the cytoplasmic domain, of human CD46 are crucial for infection of subgroup B adenovirus serotype 35. *J Control Release* 113, 271-8.

Santini, C., Brennan, D., Mennuni, C., Hoess, R.H., Nicosia, A., Cortese, R. and Luzzago, A. (1998) Efficient display of an HCV cDNA expression library as C-terminal fusion to the capsid protein D of bacteriophage lambda. *J Mol Biol* 282, 125-35.

Saul, F.A., Arie, J.P., Vulliez-le Normand, B., Kahn, R., Betton, J.M. and Bentley, G.A. (2004) Structural and functional studies of FkpA from *Escherichia coli*, a

cis/trans peptidyl-prolyl isomerase with chaperone activity. *J Mol Biol* 335, 595-608.

Schlosshauer, B. and Herzog, K.H. (1990) Neurothelin: an inducible cell surface glycoprotein of blood-brain barrier-specific endothelial cells and distinct neurons. *J Cell Biol* 110, 1261-74.

Schrama, D., Reisfeld, R.A. and Becker, J.C. (2006) Antibody targeted drugs as cancer therapeutics. *Nat Rev Drug Discov* 5, 147-59.

Schroers, R., Hildebrandt, Y., Hasenkamp, J., Glass, B., Lieber, A., Wulf, G. and Piesche, M. (2004) Gene transfer into human T lymphocytes and natural killer cells by Ad5/F35 chimeric adenoviral vectors. *Exp Hematol* 32, 536-46.

Schubert, U., Anton, L.C., Gibbs, J., Norbury, C.C., Yewdell, J.W. and Bennink, J.R. (2000) Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* 404, 770-4.

Schuster, V.L., Lu, R., Kanai, N., Bao, Y., Rosenberg, S., Prie, D., Ronco, P. and Jennings, M.L. (1996) Cloning of the rabbit homologue of mouse 'basigin' and rat 'OX-47': kidney cell type-specific expression, and regulation in collecting duct cells. *Biochim Biophys Acta* 1311, 13-9.

Segerman, A., Atkinson, J.P., Marttila, M., Dennerquist, V., Wadell, G. and Arnberg, N. (2003) Adenovirus type 11 uses CD46 as a cellular receptor. *J Virol* 77, 9183-91.

Saulberger, H., Lottspeich, F. and Risau, W. (1990) The inducible blood--brain barrier specific molecule HT7 is a novel immunoglobulin-like cell surface glycoprotein. *EMBO J* 9, 2151-8.

- Shayakhmetov, D.M., Papayannopoulou, T., Stamatoyannopoulos, G. and Lieber, A. (2000) Efficient gene transfer into human CD34(+) cells by a retargeted adenovirus vector. *J Virol* 74, 2567-83.
- Shokri, A., Sanden, A.M. and Larsson, G. (2003) Cell and process design for targeting of recombinant protein into the culture medium of *Escherichia coli*. *Appl Microbiol Biotechnol* 60, 654-64.
- Sidhu, S.S. (2000) Phage display in pharmaceutical biotechnology. *Curr Opin Biotechnol* 11, 610-6.
- Sidhu, S.S. (2001) Engineering M13 for phage display. *Biomol Eng* 18, 57-63.
- Sidhu, S.S., Mengistab, A.T., Tauscher, A.N., LaVail, J. and Basbaum, C. (2004) The microvesicle as a vehicle for EMMPRIN in tumor-stromal interactions. *Oncogene* 23, 956-63.
- Smith, G.P. (1985) Filamentous fusion phage: novel expression vectors that display cloned antigens on the virion surface. *Science* 228, 1315-7.
- Smith, G.P. and Petrenko, V.A. (1997) Phage Display. *Chem Rev* 97, 391-410.
- Snyder, W.B., Davis, L.J., Danese, P.N., Cosma, C.L. and Silhavy, T.J. (1995) Overproduction of NlpE, a new outer membrane lipoprotein, suppresses the toxicity of periplasmic LacZ by activation of the Cpx signal transduction pathway. *J Bacteriol* 177, 4216-23.
- Staffler, G. and Stockinger, H. (2000) Cd147. *J Biol Regul Homeost Agents* 14, 327-30.
- Staffler, G., Szekeres, A., Schutz, G.J., Saemann, M.D., Prager, E., Zeyda, M., Drbal, K., Zlabinger, G.J., Stulnig, T.M. and Stockinger, H. (2003) Selective

inhibition of T cell activation via CD147 through novel modulation of lipid rafts. *J Immunol* 171, 1707-14.

Steinberger, P., Andris-Widhopf, J., Buhler, B., Torbett, B.E. and Barbas, C.F., 3rd.

(2000a) Functional deletion of the CCR5 receptor by intracellular immunization produces cells that are refractory to CCR5-dependent HIV-1 infection and cell fusion. *Proc Natl Acad Sci U S A* 97, 805-10.

Steinberger, P., Sutton, J.K., Rader, C., Elia, M. and Barbas, C.F., 3rd. (2000b)

Generation and characterization of a recombinant human CCR5-specific antibody. A phage display approach for rabbit antibody humanization. *J Biol Chem* 275, 36073-8.

Stirnemann, K., Romero, J.F., Baldi, L., Robert, B., Cesson, V., Besra, G.S.,

Zauderer, M., Wurm, F., Corradin, G., Mach, J.P., Macdonald, H.R. and Donda, A. (2008) Sustained activation and tumor targeting of NKT cells using a CD1d-anti-HER2-scFv fusion protein induce antitumor effects in mice. *J Clin Invest* 118, 994-1005.

Stockinger, H. (1997) Interaction of GPI-anchored cell surface proteins and complement receptor type 3. *Exp Clin Immunogenet* 14, 5-10.

Stockinger H, Ebel T, Hansmann C, Koch C., Majdic O., Prager E., Patel D.D., Fox D.A., Horejsi V., Sagawa K., and Shen D.C. CD147 (neurothelin/basigin) workshop panel report. In Kishimoto T., Kikutani H., von dem Borne A.E.G.K., Goyert S>M., Mason D.Y., Miyasaka M., Moretta L., Okumura K., Shaw S.,cSpringer T.A., Sugamura K., and Zola H., eds, *Leukocyte Typing VI*, p. 760. Garland Publishing, New York. 1997:760.

Stocks, M.R. (2004) Intrabodies: production and promise. *Drug Discov Today* 9, 960-6.

Su, Y.C., Lim, K.P. and Nathan, S. (2003) Bacterial expression of the scFv fragment of a recombinant antibody specific for *Burkholderia pseudomallei* exotoxin. *J Biochem Mol Biol* 36, 493-8.

Sun, J. and Hemler, M.E. (2001) Regulation of MMP-1 and MMP-2 production through CD147/extracellular matrix metalloproteinase inducer interactions. *Cancer Res* 61, 2276-81.

Suzuki, S., Sato, M., Senoo, H. and Ishikawa, K. (2004) Direct cell-cell interaction enhances pro-MMP-2 production and activation in co-culture of laryngeal cancer cells and fibroblasts: involvement of EMMPRIN and MT1-MMP. *Exp Cell Res* 293, 259-66.

Tamm, L.K., Hong, H. and Liang, B. (2004) Folding and assembly of beta-barrel membrane proteins. *Biochim Biophys Acta* 1666, 250-63.

Tanaka, T., Williams, R.L. and Rabbitts, T.H. (2007) Tumour prevention by a single antibody domain targeting the interaction of signal transduction proteins with RAS. *Embo J* 26, 3250-9.

Tang, J., Wu, Y.M., Zhao, P., Yang, X.M., Jiang, J.L. and Chen, Z.N. (2008) Overexpression of HAb18G/CD147 promotes invasion and metastasis via alpha3beta1 integrin mediated FAK-paxillin and FAK-PI3K-Ca²⁺ pathways. *Cell Mol Life Sci* 65, 2933-42.

Tang, W., Chang, S.B. and Hemler, M.E. (2004a) Links between CD147 function, glycosylation, and caveolin-1. *Mol Biol Cell* 15, 4043-50.

- Tang, Y., Kesavan, P., Nakada, M.T. and Yan, L. (2004b) Tumor-stroma interaction: positive feedback regulation of extracellular matrix metalloproteinase inducer (EMMPRIN) expression and matrix metalloproteinase-dependent generation of soluble EMMPRIN. *Mol Cancer Res* 2, 73-80.
- Tang, Y., Nakada, M.T., Kesavan, P., McCabe, F., Millar, H., Rafferty, P., Bugelski, P. and Yan, L. (2005) Extracellular matrix metalloproteinase inducer stimulates tumor angiogenesis by elevating vascular endothelial cell growth factor and matrix metalloproteinases. *Cancer Res* 65, 3193-9.
- Taraboletti, G., D'Ascenzo, S., Borsotti, P., Giavazzi, R., Pavan, A. and Dolo, V. (2002) Shedding of the matrix metalloproteinases MMP-2, MMP-9, and MT1-MMP as membrane vesicle-associated components by endothelial cells. *Am J Pathol* 160, 673-80.
- Tavladoraki, P., Benvenuto, E., Trinca, S., De Martinis, D., Cattaneo, A. and Galeffi, P. (1993) Transgenic plants expressing a functional single-chain Fv antibody are specifically protected from virus attack. *Nature* 366, 469-72.
- Tavladoraki, P., Girotti, A., Donini, M., Arias, F.J., Mancini, C., Morea, V., Chiaraluce, R., Consalvi, V. and Benvenuto, E. (1999) A single-chain antibody fragment is functionally expressed in the cytoplasm of both *Escherichia coli* and transgenic plants. *Eur J Biochem* 262, 617-24.
- Tayapiwatana, C., Arooncharus, P. and Kasinrer, W. (2003) Displaying and epitope mapping of CD147 on VCSM13 phages: influence of *Escherichia coli* strains. *J Immunol Methods* 281, 177-85.

- Tayapiwatana, C., Chotpadiwetkul, R. and Kasinrerak, W. (2006) A novel approach using streptavidin magnetic bead-sorted *in vivo* biotinylated survivin for monoclonal antibody production. *J Immunol Methods* 317, 1-11.
- Tollefson, A.E., Ryerse, J.S., Scaria, A., Hermiston, T.W. and Wold, W.S. (1996) The E3-11.6-kDa adenovirus death protein (ADP) is required for efficient cell death: characterization of cells infected with adp mutants. *Virology* 220, 152-62.
- Toole, B.P. (2003) Emmprin (CD147), a cell surface regulator of matrix metalloproteinase production and function. *Curr Top Dev Biol* 54, 371-89.
- Tuve, S., Chen, B.M., Liu, Y., Cheng, T.L., Toure, P., Sow, P.S., Feng, Q., Kiviat, N., Strauss, R., Ni, S., Li, Z.Y., Roffler, S.R. and Lieber, A. (2007) Combination of tumor site-located CTL-associated antigen-4 blockade and systemic regulatory T-cell depletion induces tumor-destructive immune responses. *Cancer Res* 67, 5929-39.
- Ura, T., Yoshida, A., Xin, K.Q., Yoshizaki, S., Yashima, S., Abe, S., Mizuguchi, H. and Okuda, K. (2008) Designed recombinant adenovirus type 5 vector induced envelope-specific CD8(+) cytotoxic T lymphocytes and cross-reactive neutralizing antibodies against human immunodeficiency virus type 1. *J Gene Med.*
- Uren, A.G., Kool, J., Berns, A. and van Lohuizen, M. (2005) Retroviral insertional mutagenesis: past, present and future. *Oncogene* 24, 7656-72.

- Walton, T.A. and Sousa, M.C. (2004) Crystal structure of Skp, a prefoldin-like chaperone that protects soluble and membrane proteins from aggregation. *Mol Cell* 15, 367-74.
- Wang, L., Ku, X.M., Li, Y., Bian, H.J., Zhang, S.H., Ye, H., Yao, X.Y., Li, B.H., Yang, X.M., Liao, C.G. and Chen, Z.N. (2006) Regulation of matrix metalloproteinase production and tumor cell invasion by four monoclonal antibodies against different epitopes of HAb18G/CD147 extracellular domain. *Hybridoma (Larchmt)* 25, 60-7.
- Wang, L., Miller, A. and Kendall, D.A. (2000) Signal peptide determinants of SecA binding and stimulation of ATPase activity. *J Biol Chem* 275, 10154-9.
- Wang, W.J., Li, Q.Q., Xu, J.D., Cao, X.X., Li, H.X., Tang, F., Chen, Q., Yang, J.M., Xu, Z.D. and Liu, X.P. (2008) Interaction between CD147 and P-glycoprotein and their regulation by ubiquitination in breast cancer cells. *Chemotherapy* 54, 291-301.
- Weiner, L. and Model, P. (1994) Role of an *Escherichia coli* stress-response operon in stationary-phase survival. *Proc Natl Acad Sci U S A* 91, 2191-5.
- Wheeler, Y.Y., Chen, S.Y. and Sane, D.C. (2003a) Intrabody and intrakine strategies for molecular therapy. *Mol Ther* 8, 355-66.
- Wheeler, Y.Y., Kute, T.E., Willingham, M.C., Chen, S.Y. and Sane, D.C. (2003b) Intrabody-based strategies for inhibition of vascular endothelial growth factor receptor-2: effects on apoptosis, cell growth, and angiogenesis. *FASEB J* 17, 1733-5.


- Whitelegg, N.R. and Rees, A.R. (2000) WAM: an improved algorithm for modelling antibodies on the WEB. *Protein Eng* 13, 819-24.
- Wickham, T.J. (2000) Targeting adenovirus. *Gene Ther* 7, 110-4.
- Wickham, T.J., Mathias, P., Cheresch, D.A. and Nemerow, G.R. (1993) Integrins alpha v beta 3 and alpha v beta 5 promote adenovirus internalization but not virus attachment. *Cell* 73, 309-19.
- Wiethoff, C.M. and Middaugh, C.R. (2003) Barriers to nonviral gene delivery. *J Pharm Sci* 92, 203-17.
- Williams, B.R. and Zhu, Z. (2006) Intrabody-based approaches to cancer therapy: status and prospects. *Curr Med Chem* 13, 1473-80.
- Wilson, M.C., Meredith, D. and Halestrap, A.P. (2002) Fluorescence resonance energy transfer studies on the interaction between the lactate transporter MCT1 and CD147 provide information on the topology and stoichiometry of the complex in situ. *J Biol Chem* 277, 3666-72.
- Wilson, M.R. and Easterbrook-Smith, S.B. (2000) Clusterin is a secreted mammalian chaperone. *Trends Biochem Sci* 25, 95-8.
- Winter, G., Griffiths, A.D., Hawkins, R.E. and Hoogenboom, H.R. (1994) Making antibodies by phage display technology. *Annu Rev Immunol* 12, 433-55.
- Witty, M., Sanz, C., Shah, A., Grossmann, J.G., Mizuguchi, K., Perham, R.N. and Luisi, B. (2002) Structure of the periplasmic domain of *Pseudomonas aeruginosa* TolA: evidence for an evolutionary relationship with the TonB transporter protein. *EMBO J* 21, 4207-18.

- Woodbury, R.L., Topping, T.B., Diamond, D.L., Suci, D., Kumamoto, C.A., Hardy, S.J. and Randall, L.L. (2000) Complexes between protein export chaperone SecB and SecA. Evidence for separate sites on SecA providing binding energy and regulatory interactions. *J Biol Chem* 275, 24191-8.
- Worn, A. and Pluckthun, A. (2001) Stability engineering of antibody single-chain Fv fragments. *J Mol Biol* 305, 989-1010.
- Wright, M., Grim, J., Deshane, J., Kim, M., Strong, T.V., Siegal, G.P. and Curiel, D.T. (1997) An intracellular anti-erbB-2 single-chain antibody is specifically cytotoxic to human breast carcinoma cells overexpressing erbB-2. *Gene Ther* 4, 317-22.
- Wu, E. and Nemerow, G.R. (2004) Virus yoga: the role of flexibility in virus host cell recognition. *Trends Microbiol* 12, 162-9.
- Yamaoka, T., Yonemitsu, Y., Komori, K., Baba, H., Matsumoto, T., Onohara, T. and Maehara, Y. (2005) Ex vivo electroporation as a potent new strategy for nonviral gene transfer into autologous vein grafts. *Am J Physiol Heart Circ Physiol* 289, H1865-72.
- Yan, L., Zucker, S. and Toole, B.P. (2005) Roles of the multifunctional glycoprotein, emmprin (basigin; CD147), in tumour progression. *Thromb Haemost* 93, 199-204.
- Yang, H., Zou, W., Li, Y., Chen, B. and Xin, X. (2007a) Bridge linkage role played by CD98hc of anti-tumor drug resistance and cancer metastasis on cisplatin-resistant ovarian cancer cells. *Cancer Biol Ther* 6, 942-7.

- Yang, J.M., Xu, Z., Wu, H., Zhu, H., Wu, X. and Hait, W.N. (2003) Overexpression of extracellular matrix metalloproteinase inducer in multidrug resistant cancer cells. *Mol Cancer Res* 1, 420-7.
- Yang, W., Meng, L., Wang, H., Chen, R., Wang, R., Ma, X., Xu, G., Zhou, J., Wang, S., Lu, Y. and Ma, D. (2006) Inhibition of proliferative and invasive capacities of breast cancer cells by arginine-glycine-aspartic acid peptide *in vitro*. *Oncol Rep* 15, 113-7.
- Yang, Y., Yang, S., Ye, Z., Jaffar, J., Zhou, Y., Cutter, E., Lieber, A., Hellstrom, I. and Hellstrom, K.E. (2007b) Tumor cells expressing anti-CD137 scFv induce a tumor-destructive environment. *Cancer Res* 67, 2339-44.
- Yang, Z.M., Li, E.M., Lai, B.C., Wang, Y.L. and Si, L.S. (2007c) Anti-CD3 scFv-B7.1 fusion protein expressed on the surface of HeLa cells provokes potent T-lymphocyte activation and cytotoxicity. *Biochem Cell Biol* 85, 196-202.
- Yotnda, P., Onishi, H., Heslop, H.E., Shayakhmetov, D., Lieber, A., Brenner, M. and Davis, A. (2001) Efficient infection of primitive hematopoietic stem cells by modified adenovirus. *Gene Ther* 8, 930-7.
- Yurchenko, V., Pushkarsky, T., Li, J.H., Dai, W.W., Sherry, B. and Bukrinsky, M. (2005) Regulation of CD147 cell surface expression: involvement of the proline residue in the CD147 transmembrane domain. *J Biol Chem* 280, 17013-9.
- Zdoroveac, A., Doebis, C., Laube, H., Brosel, S., Schmitt-Knosalla, I., Volk, H.D. and Seifert, M. (2008) Modulation of graft arteriosclerosis in a rat carotid transplantation model. *J Surg Res* 145, 161-9.

Zhang, Y. and Bergelson, J.M. (2005) Adenovirus receptors. *J Virol* 79, 12125-31.

Zou, W., Yang, H., Hou, X., Zhang, W., Chen, B. and Xin, X. (2007) Inhibition of CD147 gene expression via RNA interference reduces tumor cell invasion, tumorigenicity and increases chemosensitivity to paclitaxel in HO-8910pm cells. *Cancer Lett* 248, 211-8.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved