

เอกสารอ้างอิง

กองทันตสากรรณสุข กรมอนามัย กระทรวงสาธารณสุข. รายงานผลการสำรวจสภาวะทันตสุขภาพแห่งชาติ ครั้งที่ 5 พ.ศ. 2543-2544 : บริษัท สามเจริญพาณิชย์ (กรุงเทพ) จำกัด. 2545

Adonogianaki E, Mooney J, Kinane DF. The ability of gingival crevicular fluid acute phase proteins to distinguish healthy, gingivitis, periodontitis sites. *J Clin Periodontol* 1992; 19: 98-102

Akalin FA, Sengün D, Eratalay K, Renda N, Caglayan G. Hygroxyproline and total protein levels in gingiva and gingival crevicular fluid in patients with juvenile, rapidly progressive, and adult periodontitis. *J periodontol* 1993; 64: 323-329

Albandar JM, Brunelle JA, Kingman A. Destructive periodontal disease in adults 30 years od age and older in the United States, 1988-1994. *J Periodontol* 1999; 70: 13-29

Armitage GC. Periodontal disease: Diagnosis. *Ann Periodontol* 1996; 1: 37-215.

Armitage GC. Development of a classification system for periodontal disease and conditions. *Ann Periodontol* 1999; 4: 1-6

Armitage GC. Analysis of gingival crevicular fluid and risk of progression of periodontitis. *Periodontology 2000* 2004; 34: 109-119.

Baldwin PD, Pender N, Last KS. Effect on tooth movement of force delivery from nickel – titanium archwires. *Eur J Orthod* 1999; 21: 481-9.

Bartold PM. Proteoglycans of the periodontium: Structure, role and function. *J Periodont Res* 1987; 22: 431-444.

Bartold PM, Miki Y, McAllister B, Narayanan AS, Page RC. Glycosaminoglycans of human cementum. *J Periodontal Res*. 1988 Jan; 23(1): 13-7.

Bartold PM. A biochemical and immunohistochemical study of the proteoglycans of alveolar bone. *J Dental Res* 1990; 69: 7-19.

- Binder TA, Goodson JM, Socransky SS. Gingival fluid levels of acid and alkaline phosphatase. *J Periodont Res* 1987; 22: 14-19.
- Carranza FA, Rapley JW. Clinical features of gingivitis. In: Carranza's clinical periodontology. 9th edition. Philadelphia: W.B. Saunders company; 2002: 269-278
- Cox SW, Eley BM. Cathepsin B/L-, elastase-, tryptase, trypsin- and dipeptidyl peptidase IV-like activities in gingival crevicular fluid. A comparison of levels before and after basic periodontal treatment of chronic periodontitis patients. *J Clin Periodontol* 1992; 19: 333-339
- Curtis MA, Griffiths GS, Price SJ, Coulthurst SK, Johnson NW. The total proteins concentration of gingival crevicular fluid. Variation with sampling time and gingival inflammation. *J Clin Periodontol* 1988; 15: 628-632.
- Embery G, Oliver WM, Stanbury JB, Purvis JA. The electrophoretic detection of acidic glycosaminoglycans in human gingival sulcus fluid. *Arch Oral Biol* 1982; 27: 177-79.
- Embery G, Last KS. Biochemical markers of periodontal tissue destruction. *Dent Update* 1989; 2: 167-72.
- Embery G, Waddington RJ, Hall RC, Last KS. Connective tissue elements as diagnostic aids in periodontology. *Periodontology* 2000 2000; 24: 193-214
- Figueroedo CMS, Gustafsson A. Increased amount of laminin in GCF from untreated patients with periodontitis. *J Clin Periodontol* 2000; 27: 313-318
- Genco RJ, Zambon JJ, Murray PA. Serum and gingival fluid antibodies as adjuncts in the diagnosis of *Actinobacillus actinomycetemcomitans*-associated periodontal disease. *J Periodontol* 1985; 56 (suppl.): 41-50
- Giannobile WV, Riviere GR, Gorski JP, Tira DE, Cobb CM. Glycosaminoglycans and periodontal disease: Analysis of GCF by safranin O. *J Periodontol* 1993; 64: 186-190.

- Giannobile WV, Al-Shammary KF, Sarment DP. Matrix molecules and growth factors as indicators of periodontal disease activity. *Periodontology 2000* 2003; 31: 125-134
- Giannopoulou C, Demeurisse C, Cimasoni G. Elastase release from gingival crevicular and peripheral neutrophils in periodontitis and health. *Arch Oral Biol* 1994; 39: 741-745
- Haerian A, Adonogianaki E, Mooney J, Docherty JP, Kinane DF. Gingival crevicular stromelysin, collagenase and tissue inhibitor of metalloproteinase levels in healthy and disease sites. *J Clin Periodontol* 1995; 22: 505-509
- Jaito N, Jotikasthira D, Krisanaprakornkit S, Ong-chai S, Kongtawelert P. Monitoring of chondroitin-6-sulfate levels in gingival crevicular fluid during orthodontic canine movement. Biological Mechanisms of Tooth Eruption, Resorption and Movement Edited by Davidovitch Z, Mah J and Suthanarak S 2006: 289-295
- Kagayama M, Sanano Y, Mizoguchi I, Kamo N, Takahashi I, Mitani H. Localization of glycosaminoglycans in periodontal ligament during physiological and experimental tooth movement . *J Periodont Res* 1996; 31: 229-34.
- Kido J, Nakamura T, Asahara Y, Sawa T, Kohri K, Nagata T. Osteopontin in gingival crevicular fluid. *J Periodont Res* 2001; 36: 328-333
- Kunimatsu K, Yamamoto K, Ichimaru E, Kato Y, Kato I. Cathepsins B, H and L activities in gingival crevicular fluid from chronic adult periodontitis patients and experimental gingivitis subjects. *J Periodont Res* 1990; 25: 69-73
- Kunimatsu K, Mataki S, Tanaka H, et al. A cross-sectional study on osteocalcin levels in gingival crevicular fluid from periodontal patients. *J Periodontol* 1993; 64: 865-869
- Lamster IB, Holmes LG, Gross KBW, Oshrain RL, Cohen DW et al. The relationship of β -glucuronidase activity in crevicular fluid to probing attachment loss in patients with adult periodontitis. Findings from a multicenter study. *J Clin Periodontol* 1995; 22: 36-44

- Last KS, Stanbury JB, Embery G. Glycosaminoglycans in human gingival fluids as indicators of active periodontal disease. *Arch Oral Biol* 1985; 30: 275-281.
- Last KS, Donkin C, Embery G. Glycosaminoglycans in human gingival crevicular fluid during orthodontic movement. *Arch Oral Biol* 1988; 33(12): 907-12.
- Lee AJ, Walsh TN, Hodges SJ, Rawlinson A. Gingival crevicular fluid osteocalcin in adult periodontitis. *J Clin Periodontol* 1999; 26: 252-256
- Lisgarten MA, Schifter CC, Lester L. 3-years longitudinal study of the periodontal status of adult population with gingivitis. *J Clin Periodontol* 1985; 12: 225-238.
- Lisgarten MA. Nature of periodontal disease: Pathogenic mechanisms. *J Periodont Res* 1987; 22: 172-178.
- Löe H. The gingival index, the plaque index and the retention index systems. *J Periodontol* 1967; 38: 610-616
- Löe H, Anerud A, Boysen H, Morrison E. Natural history of periodontal disease in man. *J Clin Periodontol* 1986; 13: 431-440
- Lopatin DE, Caffesse ER, Bye FL, Caffesse RG. Concentration of fibronectin in sera and crevicular fluid in various stages of periodontal disease. *J Clin Periodontol* 1989; 16: 359-364
- Mäkinen KK, Sewón L, Mäkinen P-L. Analysis in gingival crevicular fluid of two oligopeptides derived from human hemoglobin β -chain. *J Periodont Res* 1996; 31:43-46
- Nakamura T, Kido Jun-ichi, Kido R, Ohishi K, Yamauchi N, Kataoka M, Nagata T. The association of calprotectin level in gingival crevicular fluid with gingival index and the activities of collagenase and aspartate aminotransferase in adult periodontitis patients. *J Periodontol* 2000; 71: 361-367
- Nakashima K, Rochrich N, Cimasoni G. Osteocalcin, prostaglandin E2 and alkaline phosphatase in gingival crevicular fluid: their relations to periodontal status. *J Clin Periodontol* 1994; 21: 327-333

- Newman MG. Classification and epidemiology of periodontal diseases. In: Carranza's clinical periodontology. 9th edition. Philadelphia: W.B. Saunders company; 2002: 64-73
- Offenbacher S, Odie BM, Van Dyke TE. The use of crevicular fluid prostaglandin E₂ levels as a predictor of periodontal attachment loss. *J Periodont Res* 1986; 21: 101-112
- Okasaki J, Kamada A, Matsukawa F, Sakaki T, Gonda Y. Disaccharide analysis of chondroitin sulphate in human gingival crevicular fluid using high-performance liquid chromatography. *Arch Oral Biol* 1995; 40: 777-779.
- Page RC and Schroeder HE. Periodontitis in humans. In: Page RC and Schroeder HE eds. Periodontitis in man and other animals. A comparative review. Newyork: Karger; 1982: 1-57
- Page RC, Altman LC, Ebersole JL, Vandesteen GE, Dahlberg WH, William BL, Osterberg SK. Rapidly progressive periodontitis; A distinct clinical condition. *J Periodontol* 1983; 53: 197-209.
- Pender N, Samuel RHA, Last KS. The monitor of orthodontic tooth movement over a 2-year period by analysis of gingival crevicular fluid. *Eur J Orthod* 1994; 16: 511-20.
- Peanmaneesuk S, Trianrong C, Kongtawelert P, Ong-chai S. Chondroitin sulphate epitopes in sera of normal and osteoarthritic horses. *Chiangmai Veterinary Journal* 2003; 1: 3-10.
- Pearson CH, Gibson JG. Sulfated galactosaminoglycans of bovine periodontal ligament. Evidence for the presence of two major types of hybrids but no chondroitin sulfate. *Connect Tissue Res.* 1982; 10(2): 161-71.
- Persson GR, Page RC. Diagnostic characteristics of crevicular fluid aspartate aminotransferase (AST) levels associated with periodontal disease activity. *J Clin Periodontol* 1992; 19: 43-48
- Pothacharoen P., Teekachunhateen S, Louthrenoo W, Yingsung W, Ong-Chai S, Hardingham T, Kongtawelert P. Brief report. Raised chondroitin sulfate epitopes and hyaluronan in serum from rheumatoid arthritis and osteoarthritis patients. *OsteoArthritis and Cartilage* 2006; 14: 299-301

- Refaie A, Anuksaksathiem O, Singh G, Moran J, Dolby AE. Antibody to collagen type I in gingival crevicular fluid. *J Periodontol* 1990; 61: 289-292
- Shibutani T, Murahashi Y, Iwayama Y. Immunohistochemical localization of chondroitin sulfate and dermatan sulfate proteoglycan in human gingival connective tissue. *J Periodont Res* 1989; 24: 310-313
- Shibutani T, Nishino W, Shiraki M and Iwayama. ELISA detection of glycosaminoglycan (GAG)-linked proteoglycans in gingival crevicular fluid. *J Periodont Res* 1993; 28: 17-20
- Smith AJ, Addy M, Embrey G. Gingival crevicular fluid glycosaminoglycan levels in patients with chronic adult periodontitis. *J Clin Periodontol* 1995; 22: 355-361
- Talonpoika J, Heino J, Larjava H, Häkkinen L, Paunio K. Gingival crevicular fluid fibronectin degradation in periodontal health and disease. *Scand J Dent Res* 1989; 97: 415-421
- Talonpoika JT, Hämäläinen MM. Collagen I carboxyterminal propeptide in human gingival crevicular fluid before and after periodontal treatment. *Scand J Dent Res* 1993; 101: 154-158
- Talonpoika JT, Paunio K, Söderling E. Molecular forms and concentration of fibronectin and fibrin in gingival crevicular fluid before and after treatment. *Scand J Dent Res* 1993; 101: 375-381
- Talonpoika JT, Hämäläinen MM. Type I collagen carboxyterminal telopeptide in human gingival crevicular fluid in different clinical conditions and after periodontal treatment. *J Clin Periodontol* 1994; 21: 320-326
- Teng YT, Sodek J, McCulloch CAG. Gingival crevicular fluid gelatinase and its relationship to periodontal disease in human subjects. *J Periodont Res* 1992; 27: 544-552
- Tiengburanatam N. Production and characterization of monoclonal antibody against chondroitin-6-sulphate. *Graduate school, Chiangmai University, Thailand* 1996.

- Tsai C-C, Ho Y-P, Chen C-C. Levels of interleukin-1 β and interleukin-8 in gingival crevicular fluids in adult periodontitis. *J Periodontol* 1995; 66: 852-859
- Tynelius-Bratthall G, Ericson D, Araujo HM. Fibronectin in saliva and gingival crevices. *J Periodont Res* 1986; 21: 563-568
- Uitto VJ. Gingival crevicular fluid – an introduction. *Periodontol 2000* 2003; 31: 9-11.
- Villela B, Cogen RB, Bartolucci AA, Birkedal-Hansen H. Crevicular fluid collagenase activity in healthy, gingivitis, chronic adult periodontitis and localized juvenile periodontitis patients. *J Periodont Res* 1987; 22: 209-211
- Waddington RJ, Embery G , Last KS. Glycosaminoglycans of human alveolar bone. *Arch Oral Biol* 1989; 34: 587-9.
- Waddington RJ, Embery G. Structural characterization of human alveolar bone proteoglycans. *Arch Oral Biol* 1991; 36: 859-66.
- Waddintong RJ, Embery G, Samuels RH. Characterization of proteoglycan metabolites in human gingival crevicular fluid during orthodontic tooth movement. *Arch Oral Biol* 1994; 39: 361-8.
- Waddington RJ, Langley MS, Guida L, Iurorio G, Labella R et al. Relationship of sulphated glycosaminoglycans in human gingival crevicular fluid with active periodontal disease. *J Periodont Res* 1996; 31: 168-170
- Waddington RJ, Embery G, Smith AJ. Immunochemical detection of the proteoglycans decorin and biglycan in human gingival crevicular fluids from sites of advance periodontitis. *Arch Oral Biol* 1998; 43: 287-95.
- Yan F, Marshall R, Wynne S, Xiao Y, Bartold PM. Glycosaminoglycan in gingival crevicular fluid of patients with periodontal class II furcation involvement before and after guided tissue regeneration. A pilot study. *J Periodontol* 2000; 71: 1-7.