

## ชื่อเรื่องวิทยานิพนธ์

การประเมินพันธุพิษของน้ำเสียและตะกอนดินในบางท้องที่ในจังหวัดเชียงใหม่และลำพูน โดยวิธีการตรวจสอบความผิดปกติของโครโนโซมหัวหอมในระบบน้ำเสีย – โลเพส

## ผู้เขียน

นางสาวนันทนิจ จากรุศรีภิญ

## ปริญญา

วิทยาศาสตรมหาบัณฑิต (ชีววิทยา)

## คณะกรรมการที่ปรึกษาวิทยานิพนธ์

รศ. ดร. พิพัฒน์ ภารตะศิลปิน

ประธานกรรมการ

รศ. ดร. ศิริเพ็ญ ตรัยไชยพร

กรรมการ

## บทคัดย่อ

การประเมินพันธุพิษของน้ำเสียและตะกอนดิน/ตะกอน จากบ่ออน้ำเข้าและออกของระบบบำบัดน้ำเสียของโรงงานกระดาษสา นิคมอุตสาหกรรมภาคเหนือและโรงกำจัดน้ำเสียมหาวิทยาลัยเชียงใหม่ โดยใช้หัวหอมแดง (*Allium ascalonicum* Linn.) เป็นพืชทดสอบอาชีวบิช *Allium* anaphase-telophase chromosome aberration assay (*Allium-AA* test) และการเกิด micronucleus ในเซลล์ระยะ interphase ได้ตรวจวิเคราะห์มาปริมาณโลหะหนักแคลเมียม โครเมี่ยม ( $6^{+}$ ) และทองแดง ซึ่งเป็นสารก่อการแตกหักของโครโนโซมในน้ำเสียและตะกอนดิน/ตะกอนทุกตัวอย่างค้าย ผลที่ได้จากการ *Allium-AA* test แสดงให้เห็นว่า�้ำเสียทุกตัวอย่างและตะกอนดิน/ตะกอนเกือบทุกตัวอย่าง สามารถทำให้เกิดพันธุพิษในระดับที่รุนแรงเทียบเท่าผลของ ethylmethanesulfonate (EMS) อย่างนี้นัยสำคัญทางสถิติ ( $P > 0.05$ ) ปริมาณโลหะหนักทั้ง 3 ชนิดในน้ำเสียทุกตัวอย่างมีค่าอยู่ในช่วง 0.0013-0.0551 mg/L ซึ่งมีค่าต่ำกว่ามาตรฐานน้ำทึ้งจากโรงงานอุตสาหกรรม ปริมาณโลหะหนักในตะกอนดิน/ตะกอนทุกตัวอย่างมีค่าอยู่ในช่วง 0.44-254.45 mg/Kg และมีค่าต่ำกว่าค่ามาตรฐานปริมาณโลหะหนักในภาคตะกอนน้ำเสียของสหราชอาณาจักรและสหภาพยุโรป ที่ยอมให้มีมากที่สุด เมื่อนำไปใช้ในการเกษตร ดังนั้นอาจกล่าวได้ว่าน้ำจะเป็นปัจจัยอันที่ไม่ได้ตรวจวิเคราะห์ที่มีผลต่อการเกิดพันธุพิษ

**Thesis Title** Genotoxic Assessment of Wastewater and Sediment in Some Areas in Chiang Mai and Lamphun Provinces Using the *Allium* Anaphase – Telophase Chromosome Aberration Assay

**Author** Miss Nanthnit Jaruseranee

**Degree** Master of Science (Biology)

**Thesis Advisory Committee** Assoc. Prof. Dr. Thipmani Paratasilpin Chairperson  
Assoc. Prof. Dr. Siripen Traichaiyaporn Member

## ABSTRACT

Genotoxic assessment of wastewater and sediment/sludge samples from the influent and effluent ponds of paper mill plant, Northern Region Industrial Estate and Chiang Mai university wastewater treatment plant was carried out. Shallot bulbs (*Allium ascalonicum* Linn.) were used in the *Allium* anaphase-telophase chromosome aberration assay (*Allium-AA* test) and micronucleus induction in interphase cells. All samples were also analyzed for 3 clastogenic heavy metal contents, Cd, Cr (6<sup>+</sup>) and Cu. The result from the *Allium-AA* test showed that a harmful level of genotoxicity was reached in all wastewater samples and most of sediment/sludge samples, with the effect comparable to that of ethylmethanesulfonate (EMS) at statistically significant level ( $P > 0.05$ ). The 3 heavy metal contents in all wastewater samples ranged from 0.0013-0.0551 mg/L which were under the industrial effluent standard. Those in all sediment/sludge samples ranged from 0.44-254.45 mg/Kg and were under the standard of heavy metals in sludge of the United States of America and the European Union at the maximum value allowing for agricultural use. Therefore it could be assumed that other factors unanalyzed might be engaged in inducing genotoxicity.