

REFERENCES

- Alam, M., and Pearson, M.J., 1990, Biscadinanes in oils from the Surma basin, Bangladesh, *Organic Geochemistry*, V.15, pp. 461-464.
- Ammosov, I.I., 1968, Coal organic matter as a parameter of degree of sedimentary rock lithification, *Proceedings of the 23rd International geological Congress*, Section 11, Prague, pp. 23-30.
- Bal A.A., Burgisser H.M., Harris D.K., Herber M.A., Rigby S.M., Thumprasertwong S., and Winkler F.J., 1992, The Tertiary Phitsanulok lacustrine basin, Thailand, In Piencharoen C. (Ed), *Proceedings of the National Conference on Geologic Resource of Thailand: Potential for Future Development*, Department of Mineral Resources, Bangkok, pp. 247-258.
- Baum, F., Braun, E.V., Hess, A., and Koch, K.E., 1982, *Geological map of Northern Thailand 1:250,000*, Federal Institute for Geosciences and Natural Resources, Hannover.
- Belay, G., 1992, *Distribution and characteristics of petroleum source rocks in oil well IF 30 O3S Ban Nong Yao, Fang basin, Northern Thailand*, unpublished M.Sc. Thesis, Chiang Mai University, 165p.
- Berner, R.A., 1989, Biogeochemical cycles of carbon and sulfur and the effect on atmospheric oxygen over Phanerozoic time, *Palaeogeography, Palaeoclimate, Palaeoecology*, (Global Planetary Change), V. 75, pp. 97-122.
- Berner, R.A., and Raiswell, R., 1983, Burial of organic carbon and pyrite sulfur in sediments over Phanerozoic time: a new theory, *Geochimica et Cosmochimica Acta*, V. 47, pp. 855-862.
- _____, 1984, C/S method for distinguishing freshwater from marine sedimentary rocks, *Geology*, V. 12, pp. 365-368.
- Boonyarat, N., 2001, *Geochemistry of formation water from the Mae Sot Formation, Fang Oil Field, Changwat Chiang Mai*, Unpublished M.Sc. Thesis, Chiang Mai University, 107 p.

- Bordenave, M.L., Espitalie, P., Leplat, P., Oudin, J.L., and Vandenbroucke, M., 1993, Screening techniques for source rock evaluation, In: Bordenave, M.L. (Ed.), Applied Petroleum Geochemistry, Éditions Technip, Paris, pp. 219-278.
- Bostick, N.H., 1979, Microscopic measurement of the level of catagenesis of solid organic matter in sedimentary rocks to aid exploration for petroleum and determine former burial temperature—a review, Society of Economic Paleontologists and Mineralogists, V. 26, pp. 17-43.
- Bray, E.E., and Evans, E.D., 1961, Distribution of n-paraffins as a clue to recognition of source beds, *Geochimica et Cosmochimica Acta*, V. 22, pp. 2-15.
- Brooks, J.D., Gould, K., and Smith, J.D., 1969, Isoprenoid hydrocarbons in coal and petroleum, *Nature*, V. 222, pp. 257-259.
- Bunopas, S., 1976, Geological map of Changwat Phitsanulok scale 1:250,000, Sheet NE 47-15, Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand.
- Bunopas, S., 1977, Geological map of Changwat Suphanburi scale 1:250,000, Sheet ND 47-7, Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand.
- Bunopas, S. and Vella, P., 1983. Opening of the gulf of Thailand: rifting of continental S.E. Asia and Late Cenozoic Tectonics, *Journal of Geological Society of Thailand*, V. 6, pp. 1–12.
- Carroll, A.R., and Bohacs, K.M., 2001, Lake-type controls on petroleum source rock potential in nonmarine basins, *American Association of Petroleum Geologists Bulletin*, V. 85, pp. 1033-1053.
- Chaodumrong, P., Janmaha, S., Leow, N.S., Praditan, S., Snansieng, S., and Ukakimapan, Y., 1983, A review of the Tertiary sedimentary rocks of Thailand, Workshop on Stratigraphic Correlation of Thailand and Malaysia, pp. 159-187.
- Chinbunchorn, N., Praditan, S., and Sattayarak, N., 1989, Petroleum Potential of Tertiary intermountane basins in Thailand, in Thanasuthipitak, T. and Ounchanum, P. (Eds.), *Proceeding of the International Symposium on Intermontane Basins: Geology & Resources*, Chiang Mai University, Thailand, pp. 29-42.

- Cook, A.C., and Sherwood, N.R., 1991, Classification of oil shales, coal and other organic-rich rocks, *Organic Geochemistry*, V. 17, pp. 211-222.
- Didyk, B.M., Simoneit, B.R.T., Brassell, S.C., and Eglinton, G., 1978, Organic geochemical indicators of palaeoenvironmental conditions of sediment, *Nature*, V. 272, pp. 216-222.
- Enminger, A., Albrecht, P., Ourisson, G., and Tissot, B., 1977, Evaluation of the polycyclic alkanes under the effect of burial (Early Toarcian shales, Paris Basin); In: Campos, R. and Goni, J. (Ed), *Advances in Organic Geochemistry 1975*, Enadimsa, Madrid, pp. 45-52.
- Flint, S., Steart, D.J., Hyde, T., Gevers, E.C.A., Dubrule, O.R.F., and Van Riessen, E.D., 1988, Aspects of reservoir geology and production behavior of Sirikit oil field, Thailand: an integrated study using well and 3-D seismic data, *American Association of Petroleum Geologists Bulletin*, V. 72, pp. 1254-1269.
- Foopathanakamol, A., Ratanasthien, B., Petersen, H.I., Uttamo, W., and Wongpornchai, P., 2008, Composition and petroleum potential of lake facies in the FA-MS-48-73 well, Mae Soon structure, Fang oilfield, Northern Thailand, *Journal of Petroleum Geology*, V. 31, pp. 317-325.
- Gibling, M., and Ratanasthien, B. (1980), Cenozoic basins of Thailand and their coal deposit, A preliminary report, *Bulletin of the Geological Society of Malaysia*, V. 13, pp 27-42.
- Gransch, J.A., and Posthuma, J., 1974, On the origin of sulphur in crudes, In, Tissot, B., and Bienner, F. (Eds.), *Advances of Organic Geochemistry, 1973*, Éditions Technip, Paris, pp. 727-739.
- Grantham, P. J., Posthuma, J. and Baak, A., 1983, Triterpanes in a number of Far-Eastern crude oils, In Bjoroy, M. (Ed.), *Advances in Organic Geochemistry 1981*, pp. 675-683.
- Hoffmann C.F., Mackenzie, A.S., Lewis, C.A., Maxwell, J.R., Oudin, J.L., Durand, B., and Vanden-Broucke, M., 1984, A biological marker study of coals, shales and oils from the Mahakam Delta, Kalimantan, Indonesia, *Chemical Geology*, V. 42, pp. 1-23.

- Hood, A., Gutjahr, C.C.M., and Heacock, R.L., 1975, Organic metamorphism and generation of petroleum, American Association of Petroleum Geologists Bulletin, V. 56, pp. 986-996.
- Huang, W.Y., and Meinschein, W.G., 1979, Sterols as ecological indicators, Geochimica et Cosmochimica Acta, V. 43, pp. 739-745.
- Hunt, J.M., 1996, Petroleum geochemistry and geology, W.H. Freeman and Company, New York, 743 p.
- Hutton, A.C., 1987. Petrographic classification of oil shales, International Journal of Coal Geology, V. 8, pp. 203-231.
- Hutton, A.C., Bharati, S., and Robl, T., 1994, Chemical and petrographic classification of kerogen/macerals, Energy and Fuels, V 8, pp. 1478-1488.
- Jitapunkul, S., 1992, Geology of Ban Pa Kha Coal Deposit, Li basin, Lamphun, In: Piancharoen, C. (Ed.), Proceedings of the National Conference on Geologic Resources of Thailand: Potential for Future Development, Department of Mineral Resources, Bangkok, Thailand, November 17-24, pp. 259-272.
- John, R.B., 1986, Biological markers in the sedimentary record, Elsevier, New York, 364 p.
- Khantaprab, C. and Kaewsang, K., 1989. Sedimentary facies and petroleum potential of the Fang Basin. In: Thanasuthipitak, T. and Ounchanum, P. (Eds.), Proceedings of the International Symposium on Intermontane Basins: Geology and Resources, January 30 to February 2, Chiang Mai, Thailand, pp. 125-126.
- Kolaczowska, E., Slougui, N.E., Watt, D.S., Marcura, R.E., and Moldowan, J.M., 1990, Thermodynamic stability of various alkylated, dealkylated, and rearranged 17α - and 17β -hopane isomers using molecular mechanics calculations, Organic Geochemistry, V. 16, pp. 1033-1038.
- Knox, G. J., and Wakefield, L. L., 1983, An introduction to the Geology of the Phitsanulok basin, In: Proceedings of Conference on Geology and Mineral Resources of Thailand, Department of Mineral Resources, Bangkok, 19-28 November, pp. 1-9.
- Lawwongngam, K., and Philp, R.P., 1991, Geochemical characteristic of oils from the Sirikit oil field, Phisanulok basin, Thailand, Chemical Geology, V. 93, pp. 129-146.

- Leco corporation, 2007, Carbon and Sulfur determination-Leco induction furnace instruments, Leco corporation, Michigan, United State, 48 p.
- Legendre, S., Rich, T. H. V., Knox, G. J., Punyaprasiddhi, P., Trumpy, D. M., Wahlert, J., and Napawongse N. P., 1988, Miocene fossilvertebrates from the Nong Hen-I (A) Exploration well of Thai Shell Exploration and Production Company Limited, Phitsanulok Basin, Thailand, *Journal of Vertebrate Paleontology*, V. 8, pp. 278-289.
- Lian, H.M., and Bradley, K., 1986, Exploration and development of natural gas, Pattani basin, In Chinbunchorn, N., Pradidtan, S., and Sattayarak, N. (Eds.), *Petroleum Potential of Tertiary intermountane basins in Thailand*, in Thanasuthipitak, T. and Ounchanum, P. (Eds.), *Proceeding of the International Symposium on Intermontane Basins: Geology & Resources*, Chiang Mai University, Thailand, pp. 29-42.
- Lijmbach, G.W.M., 1975, On the origin of petroleum, *Proceedings of the 9th World Petroleum Congress*, Applied Science Publishers, London, V. 2, pp. 357-369.
- Mackenzie, A.S., 1984, Application of biological markers in petroleum geochemistry, In Brooks, J., and Welte, D.H. (Eds.), *Advances in Petroleum Geochemistry*, Academic Press, London, V. 1, pp. 115-241.
- McKirdy, D.M., Cox, R.E., Volkman, J. K., and Howell, V. J., 1986, Botryococcane in new class of Australian nonmarine crude oils, *Nature*, V. 320, pp. 57-59.
- Miles, J.A., 1989, *Illustrated glossary of petroleum geochemistry*, Clarendon Press, Oxford, 137 p.
- Moldowan, J.M., and Seifert, W.K., 1980, First discovery of botryococcane in petroleum, *Journal of Chemical Society, Chemical Communications*, pp. 912-914.
- Morley, C.K., 2001, Combined escape tectonics and subduction rollback-back arc extension: a model for the evolution of Tertiary rift basins in Thailand, Malaysia and Laos, *Journal of the Geological Society*, V. 158, pp. 461-474.
- Mukhopadhyay, P.K., Hagemann, H.W., and Gormly, J.R., 1985, Characterization of kerogens as seen under the aspect of maturation and hydrocarbon generation, *Erdöl und Kohle-Erdgas Petrochemie vereinigt mit Brennstoff-Chemie*, V.38, pp 7-18.

- Nichols, G., and Uttamo, W., 2005, Sedimentation in a humid, interior, extensional basin: the Cenozoic Li basin, northern Thailand, *Journal of the Geological Society, London*, V. 162, pp. 333-347.
- Peters, K.E., 1986, Guidelines for evaluating petroleum source rock using programmed pyrolysis, *American Association of Petroleum Geologists Bulletin*, V. 70, pp. 318-329.
- Peters, K.E., and Moldowan, J.M., 1993, *The biomarker guide: interpreting molecular fossils in petroleum and ancient sediments*, Prentice Hall, Englewood Cliffs, New Jersey, 352 p.
- Peters, K.E., Fraser, T.H., Amris, W., Rustanto, B., and Hermanto, E., 1999, Geochemistry of crude oils from eastern Indonesia. *American Association of Petroleum Geologists Bulletin*, V. 83, pp. 1927-1942.
- Peters, K.E., Walters, C.C., and Moldowan, J.M., 2005, *The biomarker guide (Volume 1) biomarkers and isotopes in the environment and human history*, Cambridge University Press, 471p.
- Petersen, H.I., 2002, A re-consideration of the "oil window" for humic coal and kerogen type III source rocks, *Journal of Petroleum Geology*, V.25, pp.407-432.
- Petersen, H.I., Foopattanakamol, A. and Ratanasthien, B., 2006, Petroleum potential, thermal maturity and the oil window of oil shales and coals in Cenozoic rift-basins, central and northern Thailand, *Journal of Petroleum Geology*, V. 29, pp. 337–360.
- Petersen, H.I., Nytoft, H.P., Ratanasthien, B., and Foopattanakamol, A., 2007, Oils from Cenozoic rift-basins in central and Northern Thailand: Source and Thermal maturity, *Journal of Petroleum Geology*, V. 30, pp. 59–78.
- Phillips, S. and Bustin, R. M., 1996. Sulfur in the Changuinola peat deposit, Panama, as an indicator of the environments of deposition of peat and coal, *Journal of Sedimentary Research*, V. 66, pp. 184–196.

- Pisutha-Arnond, S., Ukkakimapan, J., Klan-Ngern, S., Pisutha-Arnond, V., and Taiyaqpt, M., 1997, Predicting oil and water interval in U-Thong field using thermal extraction-pyrolysis gas chromatography, In Dheerailok, P., Hinthong, C., Chaodumrong, P., Putthaphiban, P., Tansathien W., Utha-aroon, C., Sattarak, N., Nuchanong, T., and Techawan, S. (Eds.), Proceedings of the International Conference on Stratigraphy and Tectonic Evolution of Southeast Asia and the South Pacific and Associated Meetings of IGCP 359 and IGCP 383 (GEOTHAI'97), Bangkok, Thailand, August 19-24, pp. 543-559.
- Piyasin, S., 1974, Geological map of Changwat Uttaradit 1:250,000, Sheet NE47-11, Geological survey division, Department of mineral of Mineral Resources, Bangkok, Thailand
- Polachan, S., and Sattayarak, N., 1989, Strike-slip tectonics and the development of Tertiary basins in Thailand, In Thanasuthipitak, T. and Ounchanum, P. (Eds.) Proceedings of the International Symposium on Intermontane Basins: Geology & Resources, Chiang Mai University, Thailand, pp.243- 253.
- Powell, T.G., and McKirdy, D.M., 1973, Relationship between ratio of pristane to phytane, crude oil composition and geological environment in Australia, Nature, V. 243, pp. 37-39.
- PTTEP, 1987, Stratigraphic log of BP1-4 well, U Thong oilfield, Suphanburi basin.
- PTTEP, 1987, Stratigraphic log of BP1-W2 well, U Thong oilfield, Suphanburi basin.
- PTTEP, 2001, Mud log of LKU-E19 well, Sirikit oilfield, Phitsanulok basin.
- Ratanasthien, B., 1997, Algae type of oil source rocks in northern Thailand, In Dheerailok, P., Hinthong, C., Chaodumrong, P., Putthapiban, P., Tansathien, W., Utha-aroon, C., Sattayarak, N., Nuchanong, T. and Techawan, S. (Eds.), Proceedings of the International Conference on Stratigraphy and Tectonic Evolution of Southeast Asia and the South Pacific and Associated Meetings of IGCP 359 and IGCP 383 (GEOTHAI'97), Department of mineral Resources, Bangkok, Thailand, August 19-24, pp.606-612.
- _____, 1999, Association of oil source algae in some Tertiary basin in Northern Thailand, Journal of Asian Earth Science, V.17, pp.295-299.

- Ratanasthien, B., Kandharosa, W., Chompusri S., and Chartprasert, S., 1999, Liptinite in coal and oil source rocks in northern Thailand, *Journal of Asian Earth Science*, V.17, pp. 301-306.
- Riva, A., Salvatori, T., Cavaliere, R., Ricchiuto, T., and Novelli, L., 1986, Origin of oils in Po basin, Northern Italy, In Leythaeuser, D., and Rullkötter, J. (Eds.), *Advances in Organic Geochemistry*, 1985, pp. 391-400.
- Rodjanapo, S., 1998, Facies analysis of Tertiary rocks in Mae Soon and Nong Yao oilfields Fang basin, Northern Thailand, unpublished M.Sc. Thesis, Chiang Mai University, 282 p.
- Ronghe, S., and Surarat, K., 2002, Acoustic impedance interpretation for sand distribution adjacent to a rift boundary fault, Suphan Buri basin, Thailand, *American Association of Petroleum Geologists Bulletin*, V.86, No.10, pp.1753-1771.
- Scalan, R.S., and Smith, J.E., 1970, An improved measure of the odd-to-even predominance in the normal alkanes of sediment extracts and petroleum, *Geochimica et Cosmochimica Acta*, V. 34, pp. 611-620.
- Schiber, J., 1999, Microbial mats in terrigenous clastics: the challenge of identification in the rock record, *Palaios*, V.14, pp. 3-14.
- Schoell, M., Teschner, M., Wehner, H., Durand, B., and Oudin, J.L., (1983), Maturity related biomarker and stable isotope variations and their application to oil/source rock correlation in the Mahakam Delta, Kalimantan, In Bjorøy, M., Albrecht, C., and Cornford, C., (Eds.), *Advances in organic Geochemistry 1981*, John Wiley&Sons, New York, pp. 156-163.
- Seifert, W.K., and Moldowan, J.M., 1978, Applications of steranes, terpanes and monoaromatics to the maturation, migration and source of crude oils, *Geochimica et Cosmochimica Acta*, V.42, pp. 77-95.
- _____ 1980, The effect of thermal stress on source-rock quality as measured by hopane stereochemistry, *Physics and Chemistry of the earth*, V. 12, pp. 229-237.
- Sethakul, N., 1984, Pongnok oil field, In Thanasuthipitak, P. (Ed.), *Symposium on Cenozoic Basins of Thailand: Geology and Resources*, Chiang Mai University, 153 p.

- _____, 1985. Petroleum geology of Fang Basin Part I. Unpublished, report, Geology Division Oil Exploration Production, Defense Energy Department, 21 p.
- Shanmugam, G., 1985, Significance of coniferous rain forests and related organic matter in generating commercial quantities of oil, Gippsland basin, Australia, American Association of Petroleum Geologists Bulletin, V. 69, pp. 1241-1254.
- Sherwood, N.R., Cook, A.C., Gibling, M. and Tantisukrit, C., 1984, Petrology of a suite of sedimentary rocks associated with some coal-bearing basins in northwestern Thailand, International Journal of Coal Geology, V. 4, pp. 45–71.
- Shinbaoka, M., Bennet, A. J. R., and Goild, K.W., 1973, Diagenesis of organic matter and occurrence of hydrocarbons in some Australian sedimentary basins, Australian Petroleum Exploration Association Journal, V. 13, pp. 73-80.
- Simoneit, B.R.T., Grimalt, J.G., Wang, T.G., Cox, R.E., Hatcher, P.G., and Nissenbaum, A., 1986, Cyclic terpenoids of contemporary resinous plant detritus and of fossil woods, ambers and coal, Organic Geochemistry, V. 10, pp. 877-889.
- Snansieng, S., and Maneekut, N., 1985, Li basin, an analysis of the oldest Cenozoic basin of Thailand, Journal of Geological Society of Thailand, V. 8, pp. 29-35.
- Songtham, W., 2000, Palynology of Na Hong basin Amphoe Mae Chaem Changwat Chiang Mai, unpublished M.Sc Thesis, Chiang Mai University, 115 p.
- _____, 2003, Stratigraphic correlation of Tertiary basins in northern Thailand using algae pollen and spore, unpublished Ph.D. Thesis, Chiang Mai University, 280 p.
- Stach, E., 1975, Stach's Textbook of Coal Petrology, 2nd ed., Gebrüder Borntraeger, Berlin-Stuttgart, 535 p.
- Stach, E., Mackowsky, M. Th., Teichmüller, M., Taylor, G.H., Chandra, D., and Teichmüller, 1982, Stach's textbook of coal petrology, Gebrüder Borntraege, Berlin-Stuttgart, 535 p.

- Sukto, P., Suteethorn, V., and Boripatgosol, S., 1984, Geological map of Moulmein scale 1: 250,000, Sheet NE-47-14, Geological Survey Division, Department of Mineral Resources, Bangkok, Thailand.
- Tantisukrit, C., Gibling M., Uttamo, W., Thanasuthipitak, T., and Haraluck M., 1982, Oil shale strata in the Late Cenozoic basin at Mae Sot, Tak, In Thanasuthipitak, T. (Ed.), Annual Technical Meeting 1981, Department of Geological Sciences, Chiang Mai University, Thailand, pp.163-183.
- Taylor, G.H., Teichmuller, M., Davis, A., Diessel, C.F.K., Littke, R., and Robert, P., 1998, Organic petrology, Gebruder Borntraeger, Berlin, Stuttgart, 704 p.
- Thanomsap, S. and Sitahirun, S., 1992. The Mae Sot oil shale, In Piancharoen, C. (Ed.), Proceedings of the National Conference on Geologic Resources of Thailand: Potential for Future Development, Department of Mineral Resources, Bangkok, Thailand, November 17-24, pp. 676-691.
- Thompson, S., Cooper, B.S., Morley, R.J., and Barnard, P.C., 1985, Oil generation coals, In Thamas, B.M. (Ed.), Petroleum geochemistry in exploration of Norwegian Shelf, London, Graham and Trotman, pp. 59-73.
- Tissot, B.P., and Welte, D.H., 1978, Petroleum formation and occurrence, A new approach to oil and gas exploration, Springer-Verlag, Berlin Heidelberg New York, 538 p.
- Tissot, B.P., Pelet, R., and Ungerer, P.H., 1987, Thermal history of sedimentary basin, maturation indices, and kinetics of oil and gas generation, American Association of Petroleum Geologists Bulletin., V. 71, pp.1445-1466.
- Van Aarssen, B.G.K., Cox, H.C., Hoogendoorn, P., and De Leeuw, J.W., 1990, A cadinane biopolymer in fossil and extant demmar rasins as a source for cadinanes and biocadenane in crude oils from southeast Asia, *Geochimica et Cosmochimica Acta*, V. 56, pp. 3021-3031.
- Volkman, J.K., and Maxwell, J.R., 1986, Acyclic isoprenoids as biological markers; In Johns R.B., (Ed), Biological markers in sedimentary record, Elsevier, New York, pp. 1-42.

- Waples, D.W., and Machihara, T., 1991, Biomarkers for geologist: A Practical guide to the application of steranes and Triterpanes in petroleum geology, American Association of Petroleum Geologists, Methods in Exploration Series, No. 9, 91 p.
- Watanasak, M., 1989, Palynological zonation of Mid-Tertiary intermontane basins in northern Thailand, In Thanasuthipitak, T., and Ounchanum, P. (Eds.), Proceedings of the International Symposium on Intermontane Basins: Geology and Resources, Department of Geological Sciences, Chiang Mai University, Chiang Mai, Thailand, January 30-February 2, pp. 216-225.
- Water Resources Engineering Company, 1997, Exploration and assessment of potential of coal in Fang basin, Chiang Mai, Final Report, Department of Mineral Resources, 78 p.
- Welte, D.H., Hagemann, H.W., Hollerbach, A., Leytheuser, D., and Stahl, W., 1975, Correlation between petroleum and source rock, Proceedings of the 9th World Petroleum Congress, V. 2, pp. 179-191.
- Whitehead, E.V., 1974, The structure of petroleum pentacyclanes; In: Tissot, B., and Bienner, F. (Eds.), Advances in Organic Geochemistry 1973, Editions Technip, Paris, pp. 225-243.
- Wongpornchai, P., 1997, Origin of the formation in Nong Bua basin, Central Thailand, In Dheerailok, P., Hinthong, C., Chaodumrong, P., Putthapiban, P., Tansathien, W., Utha-aroon, C., Sattayarak, N., Nuchanong, T. and Techawan, S. (Eds.), Proceedings of the International Conference on Stratigraphy and Tectonic Evolution of Southeast Asia and the South Pacific and Associated Meetings of IGCP 359 and IGCP 383 (GEOTHAI'97), Department of Mineral Resources, Bangkok, Thailand, August 19-24, pp. 210-218.
- Workman, D.R., 1975, Tectonic evolution of Indochina, Journal of Geological Society of Thailand, V. 1, pp. 3-9.
- Zumberge, J.E., 1987, Terpenoid biomarker distributions in low maturity crude oil, Organic Geochemistry, V. 11, pp. 479-496.