
TABLE OF CONTENTS

Page
iii
iv
vii
xii
xiii
xv
1
5
19
29
51
60
78
80
81
84 85
85



ลิ<mark>ขสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

LIST OF TABLES

Tabl	e Que e	Page
1	The nutritional compositions of Spirogyra neglecta	7
2	Compositions of normal and high-fat diet	20
3	Details giving primer sequences and expected product sizes	27
	for the genes amplification	
4	Effect of Spirogyra neglecta extract on body weight, kidney	30
	weight per body weight ratio and biochemical plasma parameters	
	in experimental rats	

ลิ<mark>ปสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

LIST OF FIGURES

Figu	ire	Page
1	Fresh Spirogyra neglecta	5
2	Cell structure of Spirogyra neglecta	6
3	Diagram illustrates of several pathways that contribute	12
	to oxidative stress in response to increased glucose flux.	
4	Predicted transmembrane topology of OAT family	15
5	The classical model of the mechanism of basolateral	16
	organic anion transport system	
5	Modified predicted model for epidermal growth factor (EGF)	5 17
	stimulation on organic anion transporter 3 expression and function	
7	Diagram demonstrates the experimental design of this study	22
3	Effect of Spirogyra neglecta extract on lipid peroxidation	31
	in rat kidney tissues	
•	Para-aminohippurate transport mediated by rOat1 and rOat3	32
	in rat renal cortical tissues	
0	Estrone sulfate transport mediated by rOat3 using renal cortical slices	33
1	Effect of Spirogyra neglecta extract on insulin stimulated	35
	Oat1 and 3-mediated PAH transport	
2	Effect of Spirogyra neglecta extract on insulin stimulated	37
	Oat3 mediated ES transport	
3	Western blot analysis of rOat3 protein expression in rat kidneys	39
4	The amplification curve of antioxidant gene marker mRNA expression	41
	in renal tissues	
15	Antioxidant gene marker mRNA expressions in renal cortical tissues	44
6	Effect of Spirogyra neglecta extract on the expression of p65NF-kB	46
	in renal cortex	
17	Effect of <i>Spirogyra neglecta</i> extract on PKCα expression and activation	48
	in renal cortical tissues	
	in renal cortical tissues	

18	Effect of Spirogyra neglecta extract on the expression of	50
	PKCζ and p-PKCζ in renal cortical tissues	
19	The possible mechanisms of Spirogyra neglecta extract restores	57
	rOat1 and rOat3 functions and regulations in T2DM rat model	

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright[©] by Chiang Mai University All rights reserved

ABBREVIATIONS AND SYMBOLS

% %E °C α-KG μM μl ANOVA AGE BW cAMP Cd cGMP COX1 Cu Cu-Zn SOD DAG DM DMC DM+SN1000

DM+vit C

DNA DPM EDTA EGF EPP ERK1/2 Percentage Total energy in the diet Degree Celsius Alpha ketone glutarate Micromolar Microlitter Analysis of variance Advanced glycation end-products Body weight Cyclic adenosine monophosphate Cadmium Cyclic guanosine monophosphate Cyclooxygenase I Copper Copper-Zinc superoxide dismutase Diacylglycerol Diabetes mellitus Diabetic control Diabetic supplemented with SN at the dose of 1000 mg/kg BW Diabetic supplemented with vitamin C at the dose of 200 mg/kg BW Deoxyribonucleic acid Disintegrations per minute Ethylene dinitro tetra acetic acid Epidermal growth factor Ethyl phenylpropiolate Extracellular signal regulated kinase

ES ESRD ESRF FBG g GAE GPx HCI HDL HFD HFD

IDV JNK/SAPK

kcal/g kDa kg LDL Mn MAPK MATE1 MDA MDA-TBARS mRNA mg mg/dl mg/kg BW ml

mmol/l NaF Estrone sulfate End stage renal disease End stage renal failure Fasting blood glucose Gram Gallic acid equivalent unit Glutathione peroxidase Hydrochloric High density lipoprotein High fat diet Homeostasis model assessment of insulin resistance Integrated density value NH2-terminal Jun kinases/stress activated protein kinases Kilocalories per gram Kilodalton Kilogram Low density lipoprotein Manganese Mitogen-activated protein kinase Multidrug and toxin extrusion 1 Malondialdehyde Thiobarbiturate-malondialhyde Messenger ribonucleic acid Milligram Milligram per deciliter Milligram per kilogram body weight Milliliter Millimol per liter Sodium fluoride

NaOH Sodium hydroxide Na⁺-K⁺-ATPase Sodium potassium ATPase Normal control NC NC+SN1000 Normal with SN supplement at the dose of 1000 mg/kg BW NF-**k**B Nuclear factor keppa B Nanomol per milliliter nmol/ml NOS Nitric oxide synthase **NSAIDs** Non-steroidal anti-inflammatory drugs NTU Nephelometric turbidity units Oat1 Organic anion transporter 1 Oat3 Organic anion transporter 3 OCT Organic cation transporter $(O_2^{-\bullet})$ Superoxide anion (OH^{\bullet}) Hydroxyl radical PAH Para-amino hippurate PCR Polymerase chain reaction PI3K Phospho-inositide-3-kinase PGE2 Prostaglandin E2 PKC-α Protein kinase C-α ΡΚC-β Protein kinase C-β Protein kinase C-γ ΡΚC-γ ΡΚC-λ Protein kinase C-λ ΡΚC-ζ Protein kinase C-ζ PLA2 Phospholipase A2 PMA Phorbol ester p38 P38 mitogen-activated protein kinases PKA Protein kinase A ROS Reactive oxygen species **RT-PCR** Reverse transcription polymerase chain reaction

xvii

ลิยส์ Copyr A I I

SDS-PAGE	Sodium dodecyl sulfate polyacrylami
	gel electrophoresis
SE	Standard error of mean
SGLT	Sodium glucose transporter
SN	Spirogyra neglecta
SOD	Superoxide dismutase
STZ	Streptozotocin
TBARS	Thiobarbituric acid reactive substance
TEAC	Trolox equivalent antioxidant capacit
TMD	Transmembrane domains
TGFβ1	Tumor growth factor β 1
TNFα	Tumor necrosis factor α
T1DM	Type 1 diabetes mellitus
T2DM	Type 2 diabetes mellitus
T/M	Tissue to medium ratio
VEGF	Vascular endothelial growth factor
Zn	Zinc

ลิ<mark>ขสิทธิ์มหาวิทยาลัยเชียงใหม่</mark> Copyright[©] by Chiang Mai University All rights reserved

xviii