

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
ABSTRACT (THAI)	v
ABSTRACT (ENGLISH)	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xiv
CHAPTER 1 INTRODUCTION	1
1.1 Statement of the problem	1
1.2 Purpose and hypothesis of the study	3
1.3 Anticipated benefits	4
CHAPTER 2 LITURATURE REVIEW	5
2.1 Biomechanics for canine movement	5
2.2 Orthodontic elastomeric materials	12
2.3 Nickel-Titanium closed coil spring	15
2.4 Assessments of canine retraction	16
2.5 Gingival crevicular fluid (GCF) and enzyme-linked immunosorbent assay (ELISA)	22
CHAPTER 3 MATERIALS AND METHODS	25
3.1 Materials	25
3.2 Methods	28

CHAPTER 4 RESULTS	38
4.1 Clinical observations	38
4.2 Chondroitin sulfate (WF6 epitope) levels	43
CHAPTER 5 DISCUSSION	52
CHAPTER 6 CONCLUSIONS	58
BIBLIOGRAPHY	61
APPENDIX	68
Informed consent	72
CURRICULUM VITAE	74

LIST OF TABLES

Table	Page
4.1 Age (years) distribution by gender and number of the subjects (n) in each group in this study.	38
4.2 Range (Min, Max), Mean (\bar{x}), standard deviation (SD) of rate of mandibular canine movement (mm/month) by either a continuous or an interrupted force pattern.	39
4.3 Mean (\bar{x}), standard deviation (SD) and VAS scores of the patients' pain and discomfort resulting from either a continuous or an interrupted force pattern at the end of the 1 st and 5 th weeks during the loaded period.	40
A1 Raw data of CS (WF6 epitope) levels (ng/ μ g of total protein content) of continuous force pattern from fifteen subjects.	70
A2 Raw data of CS (WF6 epitope) levels (ng/ μ g of total protein content) of interrupted force pattern from fifteen subjects.	71

LIST OF FIGURES

Figure	Page
3.1 Orthodontic appliances (force-generating devices): Sentalloy® closed coil springs (100 gm) (a) and Dynaflex® elastomeric chains, short type (b).	26
3.2 Sample collection instruments: 1.5 ml Eppendorf tube (a) and Periopaper® Strip (b).	27
3.3 Push and pull force gauge.	28
3.4 Experimental diagram.	29
3.5 An elastomeric chain calibrated at 120 gm initial force was attached between the hook of the right mandibular first molar bracket and that of the right canine bracket (a). A Nickel-Titanium closed coil spring calibrated at 120 gm initial force was attached between the hook of the left mandibular first molar bracket and that of the left canine bracket (b).	31
3.6 Measuring the force generated by an elastomeric chain at the end of the 1 st to the 8 th weeks during the loaded period (a). Measuring the force generated by a Nickel-Titanium closed coil spring at the end of the 1 st to the 8 th weeks during the loaded period (b). A push and pull force gauge was used to measure the force generated by these force-generating devices.	32
3.7 GCF collection from experimental mandibular canines on both sides during the loaded period.	33
3.8 Measuring the distance between both reference points.	34

Figure

- 4.1 Boxplot graphs of forces, generated by elastomeric chains and measured at the beginning of the 1st week and the end of each week during the 8-week loaded period. 41
- 4.2 Profile of forces generated by either the 1st and the 2nd elastomeric chains or a Nickel-Titanium closed coil spring from a subject (S16) during the loaded period. 42
- 4.3 Boxplot graph of CS (WF6 epitope) levels of continuous force pattern at the end of each week during the 8-week unloaded period and the baseline data from the loaded period (collected at the beginning of the 1st week during the loaded period). 44
- 4.4 Boxplot graph of CS (WF6 epitope) levels of interrupted force pattern at the end of each week during the 8-week unloaded period and the baseline data from the loaded period (collected at the beginning of the 1st week during the loaded period). 45
- 4.5 Boxplot graphs of CS (WF6 epitope) levels at the beginning of the loaded period (baseline data) and at the 8-week loaded periods in the GCF of experimental mandibular canines. 46
- 4.6 GCF collections from a subject (S16) during the unloaded period. 47
- 4.7 Profile of CS (WF6 epitope) levels from a subject (S16) at the beginning of the 1st week and the end of each week during the 8-week unloaded period. 48

Figure

- 4.8 Profile of CS (WF6 epitope) levels from a subject (S16) at the beginning of the 1st week and the end of each week during the 8-week unloaded and the 8-week loaded periods. 49
- 4.9 Boxplot graphs of CS (WF6 epitope) levels by either a continuous or an interrupted force pattern show baseline data (collected at the beginning of the 1st week during the loaded period) and experimental data (collected at the end of each week during the 8-week loaded period). 51
- A1 The medians of CS (WF6 epitope) levels by a continuous force pattern: baseline data (collected at the beginning of the 1st week during the loaded period) and the end of each week during the 8-week loaded period. 68
- A2 The medians of CS (WF6 epitope) levels by an interrupted force pattern: baseline data (collected at the beginning of the 1st week during the loaded period) and the end of each week during the 8-week loaded period. 69



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University
All rights reserved