

## Table of Content

<b>ACKNOWLEDGEMENTS</b>	iii
<b>ABSTRACT (Eng)</b>	v
<b>ABSTRACT (Thai)</b>	x
<b>Table of Content</b>	xvi
<b>List of Tables</b>	xxii
<b>List of Illustrations</b>	xxxiii
<b>Introduction</b>	1
<b>Chapter 1 Literature review</b>	5
1.1 Rice Production	5
1.1.1 Rice production in Asia	5
1.1.2 Rice production in Africa	7
1.1.3 Rice production in other regions	8
1.2 Rice cultivation system	10
1.2.1 Irrigated rice ecosystem	12
1.2.2 Rainfed rice ecosystem	12
1.2.3 Deepwater and floating rice ecosystem	13
1.2.4 Upland rice ecosystem	13
1.3 Rainfed lowland rice ecosystem	14
1.3.1 Water soil conditions	14
1.3.2 Nutrient availability	17

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright © by Chiang Mai University

All rights reserved


1.4	Growth and development of rice plant	24
1.4.1	Vegetative growth stage	25
1.4.2	Reproductive growth stage	26
1.4.3	Ripening and senescence stage	27
1.5	The adaptation of plants to aerobic and anaerobic conditions	27
1.5.1	Root function in aerobic soil conditions	27
1.5.2	Plant in anaerobic conditions	28
1.5.3	Plant in alternate of aerobic and anaerobic conditions	29
1.6	Plant adaptation to phosphorus stress	30
1.7	The genotypic variation in responses to water regimes and phosphorus supply	31
1.8	Use of nutrient solution to simulate waterlogged and aerated soils for rice culture	33
<b>Chapter 2</b>	<b>Responses of upland and wetland rice cultivars to flooded and well drained soil water conditions</b>	<b>35</b>
2.1	Introduction	35
2.2	Materials and Methods	37
2.2.1	Experiment 1: Responses of rice in waterlogged and well drained soils	37
2.2.2	Experiment 2: Responses of rice in fluctuation of soil water regimes	38
2.2.3	Statistic analysis	39

2.3 Results	41
2.3.1 Experiment 1: Responses of rice in waterlogged and well drained soils	41
2.3.2 Experiment 2: Responses of rice in fluctuation of soil water regimes	54
2.4 Discussion	62
<b>Chapter 3 Solution culture: Method development for using stagnant nutrient solution culture</b>	66
3.1 Introduction	66
3.2 Materials and methods	68
3.2.1 Experiment 1: Comparing of rice growth in soil and nutrient solution culture	68
3.2.2 Experiment 2: Comparing of rice growth in stagnant and aerated nutrient solution culture	69
3.2.3 Experiment 3: Iron compound and concentration for rice growth in solution culture	70
3.2.4 Experiment 4: Comparing three levels of oxygen supply for rice growth in solution culture	71
3.2.5 Experiment 5: Comparing four levels of oxygen supply for rice growth in solution culture	72
3.2.6 Statistic analysis	72
3.3 Results	73

3.3.1	Experiment 1: Comparing of rice growth in soil and nutrient solution culture	73
3.3.2	Experiment 2: Comparing of rice growth in stagnant and aerated nutrient solution culture	80
3.3.3	Experiment 3: Iron compound and concentration for rice growth in solution culture	84
3.3.4	Experiment 4: Comparing three levels of oxygen supply for rice growth in solution culture	87
3.3.5	Experiment 5: Comparing four levels of oxygen supply for rice growth in solution culture	95
3.4	Discussion	99
<b>Chapter 4</b>	<b>Morphological and physiological responses of rice to limited phosphorus supply in aerated and stagnant solution culture</b>	<b>104</b>
4.1	Introduction	104
4.2	Materials and methods	106
4.2.1	Experiment 1 Australian rice in alteration of O <sub>2</sub> and P supplies in solution culture	106
4.2.2	Experiment 2: Short term responses of Australian rice to alteration of O <sub>2</sub> and P supplies	110
4.2.3	Experiment 3: Short term responses of Australian rice to alteration of O <sub>2</sub> and P supplies (Repeat)	111
4.2.4	Statistic analysis	112
4.3	Results	113

4.3.1	Experiment 1: Australian rice in alteration of O <sub>2</sub> and P supplies in solution culture	113
4.3.2	Experiment 2 and 3: Short term responses of Australian rice to alteration of O <sub>2</sub> and P supplies	123
4.4	Discussion	137
<b>Chapter 5</b>	<b>Genotypic variation in responses to phosphorus stress under aerobic and anaerobic conditions</b>	144
5.1	Introduction	144
5.2	Materials and methods	145
5.2.1	Experiment 1: Responses of Thai rice cultivars to P stress in solution culture	145
5.2.2	Experiment 2: Short term responses of RD7 to P stress in solution culture	146
5.2.3	Experiment 3: Phosphorus stress tolerance of three Australian rice cultivars	146
5.2.4	Statistic analysis	147
5.3	Results	147
5.3.1	Experiment 1: Responses of Thai rice cultivars to P stress in solution culture	147
5.3.2	Experiment 2: Short term responses of RD7 to P stress in solution culture	158
5.3.3	Experiment 3: Phosphorus stress tolerance of three Australian rice cultivars	167

5.4 Discussion	182
<b>Chapter 6 General discussion</b>	188
<b>Conclusion</b>	195
<b>References</b>	197
<b>Appendix</b>	216
<b>Curriculum Vitae</b>	224



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
 Copyright © by Chiang Mai University  
 All rights reserved

### List of Table

<b>Table</b>	<b>Page</b>
2.2.1.1 General characteristics of San Sai soil series used for all experiments.	40
2.3.1.1 The longest root length (cm) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for 2, 8 and 10 weeks.	43
2.3.1.2 Total root volume (ml) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for 10 weeks.	44
2.3.1.3 Root dry weight (mg/plant) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for 2 and 10 weeks. Values at 2 weeks are means of three replicates $\pm$ standard errors.	45
2.3.1.4 The maximum shoot length (cm) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for 10 weeks.	48
2.3.1.5 Shoot dry weight ( $\text{g plant}^{-1}$ ) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for four weeks.	49

- 2.3.1.6 Shoot dry weight ( $\text{g plant}^{-1}$ ) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for 10 weeks. 50
- 2.3.1.7 Root/shoot ratio of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for 10 weeks. Values are means of three replicates  $\pm$  standard errors. 51
- 2.3.1.8 Nitrogen (N), phosphorus (P) and potassium (K) contents ( $\text{mg plant}^{-1}$ ) of four rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions for eight weeks. 53
- 2.3.2.1 Maximum root length (cm) of three rice cultivars when grown in treatments including waterlogging 2-9 weeks (W++++) and drainage at 2-3 (W0+++), 4-5 (W+0++), 6-7 (W++0+), and 4-9 weeks (W+000), values for harvests at seven and nine weeks. 56
- 2.3.2.2 Total root volume (ml) of three rice cultivars when grown in when grown in treatments including waterlogging 2-7 weeks (W+++), and drainage at 2-3 (W0++), 4-5 (W+0+), 6-7 (W++0), and 4-7 weeks (W+00) soil conditions at seven weeks. 57
- 2.3.2.3 Root dry weight and shoot dry weight ( $\text{g plant}^{-1}$ ) of three rice cultivars when grown in treatments including waterlogging 2-5 weeks (W++) and drainage at 2-3 (W0+), 4-5 (W+0) in soil at five weeks. 58



- 2.3.2.4 Nitrogen concentration (%) of three rice cultivars when grown in when grown in treatments including waterlogging 2-7 weeks (W+++), and drainage at 2-3 (W0++), 4-5 (W+0+), 6-7 (W++0), and 4-9 weeks (W+00) soil conditions at seven weeks. 59
- 2.3.2.5 Phosphorus content (mg plant<sup>-1</sup>) of three rice cultivars when grown in waterlogged (W+) and well drained (W0) soil conditions at three weeks. 60
- 2.3.2.6 Potassium concentration (%) of three rice cultivars when grown in when grown in treatments including waterlogging 2-7 weeks (W+++), and drainage at 2-3 (W0++), 4-5 (W+0+), 6-7 (W++0), and 4-9 weeks (W+00) soil conditions at seven weeks. 61
- 3.3.1.1 Root, shoot and total plant dry weight (g plant<sup>-1</sup>) of KDML105 when grown in soil (waterlogged; W+, well drained; W0) and nutrient solution (Aerated; A and Stagnant; S) for two weeks. 74
- 3.3.1.2 A maximum shoot length (cm) and number of leaves of KDML105 when grown in soil (waterlogged; W+, well drained; W0) and nutrient solution (Aerated; A and Stagnant; S) cultures for three weeks. 75
- 3.3.1.3 The number of roots and shoot and root dry weights (g plant<sup>-1</sup>) of KDML105 when grown in soil (waterlogged; W+, well drained; W0) and nutrient solution (Aerated; A and Stagnant; S) for three weeks. 76

- 3.3.1.4 Root growth of KDML105 when grown in soil (waterlogged; W+, well drained; W0) and nutrient solution (Aerated; A and Stagnant; S) for four weeks. 77
- 3.3.1.5 Shoot growth of KDML105 when grown in soil (waterlogged; W+, well drained; W0) and nutrient solution (Aerated; A and Stagnant; S) cultures for four weeks. 78
- 3.3.1.6 Root and shoot growth of KDML105 when grown in soil (waterlogged; W+, well drained; W0) and nutrient solution (Aerated; A and Stagnant; S) cultures for five and seven weeks. 79
- 3.3.2.1 Root and shoot lengths (cm) of 15 rice cultivars when grown in aerated (A) and stagnant (S) nutrient solution for four weeks. 81
- 3.3.2.2 Root and leaf numbers of 15 rice cultivars when grown in aerated (A) and stagnant (S) nutrient solution for four weeks. 82
- 3.3.2.3 Root and shoot dry weight ( $\text{g plant}^{-1}$ ) of 15 rice cultivars when grown in aerated (A) and stagnant (S) nutrient solution for four weeks. 83
- 3.3.3.1 Growth of rice plants when grown in nutrient solution at 3 levels of Fe EDTA; 100, 150 and 200  $\mu\text{M}$  in no oxygen bubbling (0) or 3 hrs per day of air bubbling (3). 85
- 3.3.3.2 Dry weight ( $\text{g plant}^{-1}$ ) of rice plants when grown in nutrient solution culture at 3 levels of Fe EDTA; 100, 150 and 200  $\mu\text{M}$  in no oxygen bubbling (0) or 3 hrs per day of oxygen bubbling (3). 86

3.3.4.1	Maximum root length (cm) of three rice cultivars when grown in stagnant, still and aerated nutrient solution for four weeks.	89
3.3.4.2	Root numbers of three rice cultivars when grown in stagnant, still and aerated nutrient solution for four weeks.	90
3.3.4.3	Root dry weight (g/plant) of three rice cultivars when grown in stagnant, still and aerated nutrient solution for four weeks.	91
3.3.4.4	Shoot dry weight (g/plant) of three rice cultivars when grown in stagnant, still and aerated nutrient solution for four weeks.	92
3.3.4.5	Tiller numbers of three rice cultivar when grown in stagnant, still and aerated nutrient solution for four weeks.	93
3.3.4.6	SPAD values of three rice cultivars when grown in stagnant, still and aerated nutrient solution for four weeks.	94
3.3.5.1	Plant growth characteristics of Chainat 1 when grown in different culture conditions of oxygen concentration in nutrient solution for four weeks.	96
3.3.5.2	Plant growth characteristics of Chainat1 when grown in different culture conditions of oxygen concentration in nutrient solution for six weeks.	97
3.3.5.3	Nutrient content (mg plant <sup>-1</sup> ) of Chainat1 when grown in different culture conditions of oxygen concentration in nutrient solution for six weeks.	98

- 4.2.1 Treatments of oxygen (aerated; A and stagnant; S) and phosphorus (High P; 200  $\mu\text{M}$  and Low P; 1.6  $\mu\text{M}$ ) levels imposed on Australian rice (cv. Amaroo) in different periods. 109
- 4.3.1.1 The number of adventitious roots per plant and root/shoot ratio of Amaroo when grown in aerated and stagnant nutrient solution culture at high P supply for two weeks. 116
- 4.3.1.2 Root porosity (%) of a whole root system and aerenchyma formation (%) at 50 mm from the root tip and 20 mm from the root base of Amaroo when grown in aerated and stagnant nutrient solution culture at high P supply for two weeks. 117
- 4.3.1.3 The maximum root length (cm), root/shoot ratio and photosynthesis rate ( $\mu\text{mol m}^2 \text{s}^{-1}$ ) of Amaroo after transferred from aerated or stagnant nutrient solution culture at high P supply to aerated and stagnant nutrient solution culture at high or low P supply for one week. 118
- 4.3.1.4 Root porosity (%), aerenchyma formation (%) at 20 and 50 mm from root tip of Amaroo after transferred from aerated or stagnant nutrient solution culture at high P supply to aerated and stagnant nutrient solution culture at high or low P supply for one week. 119
- 4.3.1.5 The maximum root and shoot length (cm) and number of leaves per plant of Amaroo after transferred from aerated or stagnant nutrient solution culture at high P supply to aerated and stagnant nutrient solution culture at high or low P supply for two weeks. 120

- 4.3.1.6 Root and shoot dry weight ( $\text{g plant}^{-1}$ ) and root/shoot ratio of Amaroo after transferred from aerated or stagnant nutrient solution culture at high P supply to aerated and stagnant nutrient solution culture at high or low P supply for two weeks. 121
- 4.3.1.7 Photosynthesis rate ( $\mu\text{mol m}^2 \text{s}^{-1}$ ), root porosity (%) of a whole root system and aerenchyma formation (%) at 50 mm from the root tip of Amaroo after transferred from aerated or stagnant nutrient solution culture at high P supply to aerated and stagnant nutrient solution culture at high or low P supply for two weeks. 122
- 4.3.2.1 Length of longest roots of rice (cv. Amaroo) grown in aerated solution with high P ( $200 \mu\text{M}$ ) and then transferred to aerated or stagnant nutrient solution at low ( $1.6 \mu\text{M}$ ) or high ( $200 \mu\text{M}$ ) P supply for four and eight days. Values are means of three replicates  $\pm$  standard errors. 127
- 4.3.2.2 Adventitious root and tiller numbers of rice (cv. Amaroo) grown in aerated solution with high P ( $200 \mu\text{M}$ ) level (initial condition) and then transferred to aerated or stagnant nutrient solution at low ( $1.6 \mu\text{M}$ ) or high ( $200 \mu\text{M}$ ) P supply for eight days. Values are means of three replicates  $\pm$  standard errors. 128
- 4.3.2.3 Root and shoot growth of rice (cv. Amaroo) in aerated solution with high P ( $200 \mu\text{M}$ ) supply (initial) and then transferred to aerated or stagnant nutrient solution at low ( $1.6 \mu\text{M}$ ) or high ( $200 \mu\text{M}$ ) P supply for eight days. Values are means of three replicates 129

± standard errors.

4.3.2.4 Score for fluorescence of lignin and/or suberin in the layer of sclerenchymatous fibre cells in rice roots when grown in aerated solution with high P (200  $\mu\text{M}$ ) and then transferred to aerated or stagnant nutrient solution at low (1.6  $\mu\text{M}$ ) or high (200  $\mu\text{M}$ ) P supply for eight days. Values are means of three replicates ± standard errors.

131

4.3.2.5  $\text{O}_2$  consumption rate (n moles  $\text{O}_2 \text{ g}^{-1}$  Fresh Weight  $\text{s}^{-1}$ ) of rice (cv. Amaroo) roots grown in aerated solution with high P (200  $\mu\text{M}$ ) and then transferred to aerated or stagnant nutrient solution at low (1.6  $\mu\text{M}$ ) or high (200  $\mu\text{M}$ ) P supply for four days. Roots were cut into 5 mm segments for each zone and measured in the same solution as the growth medium. Values are means of four replicates ± standard errors.

134

4.3.2.6 Phosphorus content in whole rice plants ( $\text{mg plant}^{-1}$ ) grown in aerated solution with high P (200  $\mu\text{M}$ ) and then transferred to aerated or stagnant nutrient solution at low (1.6  $\mu\text{M}$ ) or high P supply for 1, 2, 4 and 8 days. Values are means of three replicates ± standard errors.

136

5.3.1.1 Maximum root lengths (cm) of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks.

150

- 5.3.1.2 Root numbers of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks. 151
- 5.3.1.3 Root dry weight ( $\text{g plant}^{-1}$ ) of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks. 152
- 5.3.1.4 Shoot dry weight ( $\text{g plant}^{-1}$ ) of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks. 153
- 5.3.1.5 Tiller numbers of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks 154
- 5.3.1.6 Leaf numbers of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks. 155
- 5.3.1.7 Maximum shoot lengths (cm) of 10 rice cultivars when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) for six weeks. 156
- 5.3.3.1 Maximum root length (cm) of three rice cultivars when grown in aerated or stagnant at 50, 100, 200 and 400  $\mu\text{M}$  P of nutrient solution cultures for two weeks. 168

- 5.3.3.2 Root numbers per plant of three rice cultivars when grown in aerated or stagnant at 50, 100, 200 and 400  $\mu\text{M}$  P of nutrient solution cultures for two weeks. 169
- 5.3.3.3 Root dry weight ( $\text{g plant}^{-1}$ ) of three rice cultivars when grown in aerated or stagnant at 50, 100, 200 and 400  $\mu\text{M}$  P of nutrient solution cultures for two weeks. 170
- 5.3.3.4 Maximum shoot length (cm) of three rice cultivars when grown in aerated or stagnant at 50, 100, 200 and 400  $\mu\text{M}$  P of nutrient solution cultures for two weeks. 171
- 5.3.3.5 Maximum root length (cm) of three rice cultivars when grown in aerated or stagnant at 1.6, 8, 40 and 200  $\mu\text{M}$  P of nutrient solution cultures for four weeks. 174
- 5.3.3.6 Root numbers per plant of three rice cultivars when grown in aerated or stagnant at 1.6, 8, 40 and 200  $\mu\text{M}$  P of nutrient solution cultures for four weeks. 175
- 5.3.3.7 Root dry weight ( $\text{g plant}^{-1}$ ) of three rice cultivars when grown in aerated or stagnant at 1.6, 8, 40 and 200  $\mu\text{M}$  P of nutrient solution cultures for four weeks. 176
- 5.3.3.8 Root porosity (%) of whole root system of three rice cultivars when grown in aerated or stagnant at 1.6, 8, 40 and 200  $\mu\text{M}$  P of nutrient solution cultures for four weeks. 177



- 5.3.3.9 Shoot dry weight ( $\text{g plant}^{-1}$ ) of three rice cultivars when grown in aerated or stagnant nutrient solution at 1.6, 8, 40 and 200  $\mu\text{M P}$  for four weeks. 178
- 5.3.3.10 Leaf numbers per plant of three rice cultivars when grown in aerated or stagnant nutrient solution at 1.6, 8, 40 and 200  $\mu\text{M P}$  for four weeks. 179
- 5.3.3.11 Tiller numbers per plant of three rice cultivars when grown in aerated or stagnant nutrient solution at 1.6, 8, 40 and 200  $\mu\text{M P}$  for four weeks. 180
- 5.3.3.12 Maximum shoot length (cm) of three rice cultivars when grown in aerated or stagnant at 1.6, 8, 40 and 200  $\mu\text{M P}$  of nutrient solution cultures for four weeks. 181

## List of Illustrations

Figure		Page
1.2.1	The rice lands classified by water regime and predominant rice type.	11
1.3.2.2.1	Typical seasonal total dry matter production and total nitrogen accumulation of rice plant (Guindo <i>et al.</i> , 1994a).	22
2.3.1.1	Aerenchyma appearance in adventitious roots at 5 cm from the root tip of KN when grown in well drained (A) and waterlogged (B) soil conditions compared with KDML105 when grown in well-drained (C) and waterlogged (D) soil conditions.	52
4.3.2.1	Root porosity (% gas volume per unit root volume) of rice (cv. Amaroo) grown in aerated solution with high (200 $\mu\text{M}$ ) P (HP) then transferred to aerated (open symbols) and stagnant (closed symbols) solutions at 1.6 $\mu\text{M}$ P ( $\odot$ ), and 200 $\mu\text{M}$ P ( $\square$ ). Root porosity (%) was measured for the whole root system of each plant. Bars represent standard errors of three replicates.	130
4.3.2.2	Typical transverse sections of rice roots showing autofluorescence of the walls in the outer cell layers. Plants were grown for eight days after transfer to high P level (200 $\mu\text{M}$ ) in aerated nutrient solution at 20 mm (A) and 70 mm (B) from the apex or in stagnant nutrient solution at 20 mm (C) and 70 mm (D). Increased autofluorescence in response to the stagnant treatment was evident	132

for the walls of the sclerenchymatous fibres, as well as those in the hypodermal cell layer and the cells to the immediate interior of the fibre cells. Average scoring of sections in triplicate is shown in Table 4.3.2.4.

4.3.2.3 Rate of radial O<sub>2</sub> loss (ROL) along adventitious roots of rice (cv. Amaroo) after transition to treatment solutions; aerated at 1.6 μM P (○), and 200 μM P (□) and stagnant at 1.6 μM P (●) and 200 μM P (■). Rates of ROL were measured from one 101-127 mm adventitious root of each of four 28 d-old plants in each treatment, with the treatment imposed for 1 (a), 2 (b), 4 (c) or 8 (d) days. Bars represent standard errors of four replicates.

133

4.3.2.4 Phosphorus uptake of rice (mg P g<sup>-1</sup> root dry weight) after transfer to treatment at 0-4 days and 4-8 days; aerated at 1.6 μM P (A LP), and 200 μM P (A HP) and stagnant at 1.6 μM P (S LP) and 200 μM P (S HP) nutrient solution culture. Plants were 28 d-old (initial) and the uptake was calculated as the final P content minus the initial P content. Bars represent means ± standard errors of three replicates. Different letters indicate significant differences at 5% level.

135


5.3.1.1 Correlation between P level and oxygen effects on root numbers of rice when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200 μM (high P) for six weeks

157

- 5.3.2.1 Maximum root length (cm) of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days. 159
- 5.3.2.2 Root numbers per plant of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days. 160
- 5.3.2.3 Root dry weight ( $\text{g plant}^{-1}$ ) of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days. 161
- 5.3.2.4 Root/shoo ratio of CNT1 (A) and RD7 (B) when grown in aerated at low or high P and stagnant at low or high P solution cultures at initial, 2, 4 and 8 days. 162
- 5.3.2.5 Root porosity (%) of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days. 163
- 5.3.2.6 Tiller numbers per plant of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days. 164
- 5.3.2.7 Phosphorus content ( $\text{mg plant}^{-1}$ ) of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days. 165

5.3.2.8 Phosphorus uptake efficiency ( $\text{mgP g}^{-1}$  root dry weight) of CNT1 (A) and RD7 (B) when grown in aerated and stagnant nutrient solution at 1.6 (low P) and 200  $\mu\text{M}$  (high P) at initial, 2, 4 and 8 days.

166



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright © by Chiang Mai University  
All rights reserved