

TABLE OF CONTENTS

	Page
Acknowledgements	iii
Abstract in Thai	iv
Abstract in English	vii
Table of Contents	x
List of Tables	xii
List of Figures	xvi
Chapter 1	
Introduction	1
- Objectives of the experiment	3
Chapter 2	
Literature Reviews	4
- Introduction	4
- Fruit growth and development	5
- Harvest maturity	5
- Harvesting	6
- Ripening processes	8
- Nutritional values	18
- Respiration and ethylene production	21
- 1- Methylcyclopropene (1-MCP)	22
Chapter 3	
Materials and Methods	25
- Changes of quality and enzyme activity during ripening of 'Keaw Morakot' mango fruit at ambient temperature	25
- Changes of quality and enzyme activity during ripening of 'Keaw Morakot' mango fruit at low temperature	38
- Effects of 1-MCP on the changes of quality and enzyme activity during ripening of 'Keaw Morakot' mango fruit at low temperature	39

TABLE OF CONTENTS

	Page
Chapter 4	
Results	40
- Changes of quality and enzyme activity during ripening of 'Keaw Morakot' mango fruit at ambient temperature	40
- Changes of quality and enzyme activity during ripening of 'Keaw Morakot' mango fruit at low temperature	67
- Effects of 1-MCP on the changes of quality and enzyme activity during ripening of 'Keaw Morakot' mango fruit at low temperature	109
Chapter 5	
Discussions	149
Chapter 6	
Conclusions	165
References	167
Appendix	182
Curriculum Vitae	226

LIST OF TABLES

Table	Page
1 Total soluble solids (TSS, %), (titratable acid TA, %) contents and TSS/TA ratio of 'Kaew' mango fruit at different maturity stored at 15°C after storage for 4, 8, 12 and 16 days	16
2 Citric acid and malic acid contents ($\mu\text{g}/\text{mg}$ pulp) in mango fruit cultivars during maturation and ripening	17
3 Proximate analysis of mango fruit (in 100 g edible portion as ripen)	19
4 Physical and chemical characteristics and nutritive value of nine Thai mango fruits (Chok Anan, Nam Dok Mai #4, Rad, Mon Duen Gao, Keaw Sawoei, Okrong Keaw, Okrong Thong, Kaew and Maha Chanok) evaluated at fully ripe stage	20
5 Carotenoids in the pulp of some Indian mango cultivars	21
6 Effects of maturity on firmness, L^* , $^{\circ}\text{H}$ and chroma values of pulp and peel of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 0 and 7 days	41
7 Effects of maturity on TSS, glucose, fructose and sucrose contents of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 0 and 7 days	47
8 Effects of maturity on pH value, TA, citric acid, malic acid and vitamin C contents of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 0 and 7 days	51
9 Effects of maturity on dry matter, starch, chlorophylls and β -carotene contents of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 0 and 7 days	56
10 Effects of maturity on respiration rate and ethylene production of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 0 and 7 days	61

LIST OF TABLES

Table	Page
11 Effects of maturity on WSP content, PG and PME activities of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 0 and 7 days	64
12 Effects of maturity on texture, consistency, feeling and taste of 'Keaw Morakot' mango fruit stored at ambient condition (26-31°C, 60-70% RH) for 7 days	64
13 Effects of maturity on firmness, L*, °H and chroma values of pulp and peel of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH for 12 days	68
14 Effects of maturity on firmness, L*, °H and chroma values of pulp and peel of 'Keaw Morakot' mango fruit stored at 25 °C, 70-75% RH for 0 and 7 days, after storage at 13°C, 85-90% RH for 21 days	68
15 Effects of maturity on TSS, glucose, fructose and sucrose contents of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH for 12 days	78
16 Effects of maturity on TSS, glucose, fructose and sucrose contents of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH for 0 and, 7 days after storage at 13°C, 85-90% RH for 21 days	78
17 Effects of maturity on pH value, TA, citric acid, malic acid and vitamin C contents of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH for 12 days	85
18 Effects of maturity on pH value, TA, citric acid, malic acid and vitamin C contents of 'Keaw Morakot' mango fruit placed in 25°C, 70-75% RH for 0 and 7 days, after storage at 13°C, 85-90% RH for 21 days	85
19 Effects of maturity on dry matter, starch, chlorophylls and β-carotene contents of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH for 12 days	92

LIST OF TABLES

Table		Page
20	Effects of maturity on dry matter, starch, chlorophylls and β -carotene contents of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH for 0 and 7 days, after storage at 13°C, 85-90% RH for 21 days	92
21	Effects of maturity on respiration rate and ethylene production of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH for 12 days	100
22	Effects of maturity on respiration rate and ethylene production of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH for 0 and 7 days, after storage at 13°C, 85-90% RH for 21 days	100
23	Effects of maturity on WSP content, PG and PME activities of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH for 12 days	104
24	Effects of maturity on WSP content, PG and PME activities of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH for 0 and 7 days, after storage at 13°C, 85-90% RH for 21 days	105
25	Effects of maturity on texture, consistency, feeling, taste and off-odor of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH for 7 days, after storage at 13°C, 85-90% RH for 21 days	105
26	Effects of 1-MCP on firmness, L*, a^* and chroma values of pulp and peel of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH, for 0 and 12 days	110
27	Effects of 1-MCP on TSS, glucose, fructose and sucrose contents of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH, for 0, 12 and 21 days	119

LIST OF TABLES

Table	Page
28 Effects of 1-MCP on pH value, titrable acid, citric acid, malic acid and vitamin C contents of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH, for 0 and 12 days	126
29 Effects of 1-MCP on dry matter, starch, chlorophylls and β -carotene contents of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH, for 0 and 12 days	133
30 Effects of 1-MCP on respiration rate and ethylene production of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH, for 0 and 12 days	141
31 Effects of 1-MCP on WSP content, PG and PME activities of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH, for 0 and 12 days	145

LIST OF FIGURES

Figure	Page
1. Changes in the firmness of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	42
2. Changes in the L* value of 'Keaw Morakot' mango pulp during storage at ambient condition (26-31°C, 60-70% RH)	42
3. Changes in the L* value of 'Keaw Morakot' mango peel during storage at ambient condition (26-31°C, 60-70% RH)	43
4. Changes in the °H value of 'Keaw Morakot' mango pulp during storage at ambient condition (26-31°C, 60-70% RH)	43
5. Changes in °H value of 'Keaw Morakot' mango peel during storage at ambient condition (26-31°C, 60-70% RH)	44
6. Changes in the chroma value of 'Keaw Morakot' mango pulp during storage at ambient condition (26-31°C, 60-70% RH)	44
7. Changes in the chroma value of 'Keaw Morakot' mango peel during storage at ambient condition (26-31°C, 60-70% RH)	45
8. 'Keaw Morakot' mango fruit at different maturity after storage for 0 and 7 days at ambient condition (26-31°C, 60-70% RH)	45
9. Changes in the TSS content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	48
10. Changes in the glucose content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	48
11. Changes in the fructose content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	49
12. Changes in the sucrose content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	49

LIST OF FIGURES

Figure	Page
13	52
Changes in the pH value of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
14	52
Changes in the TA content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
15	53
Changes in the citric acid content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
16	53
Changes in the malic acid content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
17	54
Changes in the vitamin C content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
18	57
Changes in the dry matter content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
19	57
Changes in the starch content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
20	58
Changes in the total chlorophyll content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
21	58
Changes in the chlorophyll a content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
22	59
Changes in the chlorophyll b content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
23	59
Changes in the β -carotene content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	
24	62
Changes in the respiration rate of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	

LIST OF FIGURES

Figure	Page
25 Changes in the ethylene production of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	62
26 Changes in the WSP content of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	65
27 Changes in the PG activity of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	65
28 Changes in the PME activity of 'Keaw Morakot' mango fruit during storage at ambient condition (26-31°C, 60-70% RH)	66
29 Changes in the firmness of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	69
30 Changes in the firmness of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	69
31 Changes in the L* value of 'Keaw Morakot' mango pulp stored at 13°C, 85-90% RH	70
32 Changes in the L* value of 'Keaw Morakot' mango pulp stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	70
33 Changes in the L* value of 'Keaw Morakot' mango peel stored at 13°C, 85-90% RH	71
34 Changes in the L* value of 'Keaw Morakot' mango peel stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	71
35 Changes in the °H value of 'Keaw Morakot' mango pulp stored at 13°C, 85-90% RH	72
36 Changes in the °H value of 'Keaw Morakot' mango pulp stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	72

LIST OF FIGURES

Figure	Page
37 Changes in the °H value of 'Keaw Morakot' mango peel stored at 13°C, 85-90% RH	73
38 Changes in the °H value of 'Keaw Morakot' mango peel stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	73
39 Changes in the chroma value of 'Keaw Morakot' mango pulp stored at 13°C, 85-90% RH	74
40 Changes in the chroma value of 'Keaw Morakot' mango pulp stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	74
41 Changes in the chroma value of 'Keaw Morakot' mango peel stored at 13°C, 85-90% RH	75
42 Changes in the chroma value of 'Keaw Morakot' mango peel stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	75
43 'Keaw Morakot' mango fruit after storage at 13°C, 85-90% RH for 12 days	76
44 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH for 0 and 7 days after storage at 13°C, 85-90% RH for 21 days	76
45 Changes in the TSS content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	79
46 Changes in the TSS content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	79
47 Changes in the glucose content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	80
48 Changes in the glucose content of 'Keaw Morakot' mango fruit stored	

at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days

80

LIST OF FIGURES

Figure		Page
49	Changes in the fructose content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	81
50	Changes in the fructose content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	81
51	Changes in the sucrose content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	82
52	Changes in the sucrose content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	82
53	Changes in the pH value of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	86
54	Changes in the pH value of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	86
55	Changes in the TA content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	87
56	Changes in the TA content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	87
57	Changes in the citric acid content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	88

LIST OF FIGURES

Figure	Page
58 Changes in the citric acid content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	88
59 Changes in the malic acid content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	89
60 Changes in the malic acid content of 'Keaw Morakot' mango fruit at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	89
61 Changes in the vitamin C content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	90
62 Changes in the vitamin C content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	90
63 Changes in the dry matter content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	93
64 Changes in the dry matter content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	93
65 Changes in the starch content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	94
66 Changes in the starch content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	94
67 Changes in the total chlorophyll content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	95
68 Changes in the total chlorophyll content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90%RH for 21 days	95
69 Changes in the chlorophyll a content of 'Keaw Morakot' mango fruit	

stored at 13°C, 85-90% RH

96

LIST OF FIGURES

Figure		Page
70	Changes in the chlorophyll a content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90%RH for 21 days	96
71	Changes in the chlorophyll b content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	97
72	Changes in the chlorophyll b content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	97
73	Changes in the β -carotene content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	98
74	Changes in the β -carotene content of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	98
73	Changes in the respiration rate of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	101
76	Changes in the respiration rate of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	101
77	Changes in the ethylene production of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	102
78	Changes in the ethylene production of 'Keaw Morakot' mango fruit stored at 25°C, 70-75%RH, after storage at 13°C, 85-90% RH for 21 days	102
79	Changes in the WSP content of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	106
80	Changes in the WSP of 'Keaw Morakot' mango fruit stored at 25°C,	

	70-75% RH, after storage at 13°C, 85-90% RH for 21 days	106
81	Changes in the PG activity of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	107

LIST OF FIGURES

Figure		Page
82	Changes in the PG activity of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	107
83	Changes in the PME activity of 'Keaw Morakot' mango fruit stored at 13°C, 85-90% RH	108
84	Changes in the PME activity of 'Keaw Morakot' mango fruit stored at 25°C, 70-75% RH, after storage at 13°C, 85-90% RH for 21 days	108
85	Changes in the firmness of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90%RH	111
86	Changes in the L* value of 'Keaw Morakot' mango pulp treated with 1-MCP during storage at 13°C, 85-90%RH	112
87	Changes in the L* value of 'Keaw Morakot' mango peel treated with 1-MCP during storage at 13°C, 85-90%RH	113
88	Changes in the °H value of 'Keaw Morakot' mango pulp treated with 1-MCP during storage at 13°C, 85-90%RH	114
89	Changes in the °H value of 'Keaw Morakot' mango peel treated with 1-MCP during storage at 13°C, 85-90%RH	115
90	Changes in the chroma value of 'Keaw Morakot' mango pulp treated with 1-MCP during storage at 13°C, 85-90%RH	116
91	Changes in the chroma value of 'Keaw Morakot' mango peel treated with 1-MCP during storage at 13°C, 85-90%RH	117
92	Changes in the TSS content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	120

93	Changes in the glucose content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	121
94	Changes in the fructose content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	122

LIST OF FIGURES

Figure		Page
95	Changes in the sucrose content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	123
97	Changes in the pH value of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	127
98	Changes in the TA content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	128
99	Changes in the citric acid content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	129
100	Changes in the malic acid content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	130
101	Changes in the vitamin C content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	131
102	Changes in the dry matter content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	134
103	Changes in the starch content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	135
104	Changes in the total chlorophyll content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	136
105	Changes in the chlorophyll a content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	137
106	Changes in the chlorophyll b content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	138

107	Changes in the β -carotene content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	139
108	Changes in the respiration rate of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	142

LIST OF FIGURES

Figure		Page
109	Changes in the ethylene production of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	143
110	Changes in the WSP content of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	146
111	Changes in the PS activity of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	147
112	Changes in the PME activity of 'Keaw Morakot' mango fruit treated with 1-MCP during storage at 13°C, 85-90% RH	148