

## **REFERENCES**

- Barbier, P.1989. Genetic variation and ecotypic differentiation in the wild rice species *Oryza rufipogon*. II. Influence of the mating system and life history traits on the genetic structure of populations. Jpn J. Genet. 64:273-285p.
- Brown, A. H. D. 2000. The genetic structure of crop landrace and the challenge to conserve them *in situ* on farm. In Gene in the field: On-Farm Conservation of Crop Diversity, Lewis Publishing Co., U. S. A. pp. 29-48.
- Cai, H.W., and Morishima, H. 2000. Genomic regions affecting seed shattering and seed dormancy in rice. Theor. Appl. Genet. 100:840–846.
- Chang, T.T. 1976. The origin, evolution, cultivation, dissemination, and diversification of Asian and African rices. Euphytica 25:435–441.
- Chang, T.T. 1984. Conservation of rice genetic resources: luxury or necessity? Science 224:251–256.
- Chen, X., Temnykh, S., Xu, Y.G., and McCouch. S.R. 1997. Development of a microsatellite framework map providing genome-wide coverage in rice (*Oryza sativa* L.). Theoretical and applied genetics. 95: 553-567.

- Chen, W.M. 2001. Wild rice biological characteristics and integrated control. Paper to be presented at the 18<sup>th</sup> Asian Pacific Weed Science Society Conference, 28 May-2 June 2001, Beijing, China.
- Chen, L.J., Lee, D.S., Song, Z.P., Shu, H. S., and Lu, B.R. 2004. Gene flow from cultivated rice (*Oryza sativa*) to its weedy and wild relatives. Annals of Botany 93: 67-73.
- Chitrakon, S. 1995. Characterization, evaluation and utilization of wild rice germplasm in Thailand. PhD Thesis, Hokkaido University.
- Doyle, J.J. and J.L. Doyle. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. Focus 12: 13-15.
- Ellstrand N.C., Prentice H.C. and J.F. Hancock. 1999. Gene flow and introgression from domesticated plants into their wild relatives. Annu. Rev. Ecol. Syst. 30: 539-563.
- Gao, L.Z., Ge, S., and Hong, D.Y. 2000. Allozymic diversity and genetic structure of common wild rice *Oryza rufipogon* Griff, China. Therotical and Applied Genetics 101: 494-502.
- Gao L.Z. 2002. The conservation of rice biodiversity in China: significance, genetic erosion, ethnobotany and prospect. *Genetic Resources and Crop Evolution*, 50 (1), 17-32.
- Goudet, J. 2001. FSTAT, a program to estimate and test gene diversities and fixation indices (version 2.9.3). <http://www.unil.ch/izea/softwares/fstat.html>

- Gu, XY, Kianian, S.F., and Foley, M.E. 2004. Multiple loci and epistases control genetic variation for seed dormancy in weedy rice (*Oryza sativa*) Genetics 166: 1503–1516.
- Hamrick J.L., and Godt M.J.W. 1989. Allozyme diversity in plant species. In plant population genetics, breeding and genetic resources (ed. Brown AHD, Clegg MT, Kahler AL, Weir BS), pp.43-63. Sinauer Association Inc., Sunderland.
- Harlan, J.R. 1992. Crops and man. Second Edition. Madison. Madison, Wisconsin, USA. pp. 90-93.
- IPGRI. 2003. International Plant Genetic Resources Institute: Using molecular marker technology in studies on plant genetic diversity. Cornell University, Ithaca, Ny 14583, 3p.
- IRRI-INGER, Genetic Resource Center. 4<sup>th</sup> edition 1996. Standard Evaluation System for Rice. P.O. Box 993, Manila, Philippines. 52 p.
- Jackson M.T. 1997. Conservation of rice genetic resources: the role of the International Rice Genebank at IRRI. *Plant Molecular Biology*, 35, 61-67.
- Khush, G. S. 1997. Origin, dispersal, cultivation and variation of rice. *Plant Molecular Biology*, 35: 25-34.
- Kuroda, Y. 2004. Conservation Biological Studies on Asian Wild Rice (*O. rufipogon* and *O. nivara*) in the Vientiane Plain of Laos. Study research in Asian and African Area, Kyoto University.

- Kuroda, Y., Sato, Y.I., Bounphanousay, C., Kono, Y., and Tanaka, K. 2006. Genetic structure of three *Oryza* AA genome species (*O. rufipogon*, *O. nivara* and *O. sativa*) as assessed by SSR analysis on the Vientiane Plain of Laos. Conserv Genet (2007) 8: 149-158.
- Langevin, S.A., Clay, K., and Grace, J.B. 1990. The incidence and effect of hybridization between cultivated rice and its related weed red rice (*Oryza sativa* L.). Evolution 44(4): 1000-1008.
- Londo, J.P. Yu-Chung Chiang, Kuo-Hsiang Hung, Tzen-Yuh Chiang, and Barbara A. Schaal, 2006. Phylogeography of Asian wild rice, *Oryza rufipogon*, reveals multiple independent domestications of cultivated rice, *Oryza sativa*. vol. 103 no.25.
- Lu, B.R. 1997. A Report on IRRI-CIAP Cooperative Collecting for Wild *Oryza* Species in Cambodia, November 18-24, 1997 Genetic Resources Center, IRRI.
- Majumder, N.D., T. Ram and A.C. Sharma. 1997. Cytological and morphological variation in hybrid swarms and introgressed population of interspecific hybrids (*Oryza rufipogon* Griff. x *Oryza sativa* L.) and its impact on evolution of intermediate types. Euphytica 94: 295-302.
- Maneechote, C., S. Jamjod and B. Rerkasem. 2004. Invasion of weedy rice in rice fields in Thailand: problems and management. IRRN 29(2): 20-22.

- McCouch S.R., Chen X., Panaud O., Temnykh S., Xu Y., Cho Y.G., Huang N., Ishii T., and Blair M. 1997. Microsatellite marker development, mapping and applications in rice genetics and breeding. *Plant Molecular Biology*, 35, 89–99.
- McCouch, S.R., M. Sweeney, J. Li, H. Jiang, M. Thomson, E. Septiningsih, J. Edwards, P. Moncada, J. Xiao, A. Garris, T. Tai, C. Martinez, J. Tohme, M. Sugiono, A. McClung, L.P. Yuan and S.N. Ahn. 2007. Through the genetic bottleneck: *O. rufipogon* as a source of trait-enhancing alleles for *O. sativa*. *Euphytica* 154: 317-339.
- Men, S., and Pith, K. H. 2007. Rice Book in Cambodia. Genetic and Hybridization, CARDI 47: 78-123p.
- Morishima, H., and Oka, H.I. 1960. The pattern of interspecific variation in the genus *Oryza*: Its quantitative representation by statistical methods. *Evolution* 14: 153–165.
- Morishima, H., Oka, H.I., and Chang W.T. 1961. Directions of differentiation in populations wild rice, *Oryza perennis* and *O. sativa f. spontanea*. *Evolution*, 15, 326-339.
- Murray, M.G., and Thompson, W.F. 1980. Rapid isolation of high molecular weight plant DNA. *Nucl. Acids Res.* 8: 4321–4325.
- Nei, M. 1978. Estimation of average heterozygosity and genetic distance from a small number of individuals. *Genetics*, 89, 583-590.

- Nei, M., Tajima, F., and Tateno, Y. 1983. Accuracy of estimated phylogenetic trees from molecular data. *Journal of Molecular Evolution* 19: 153-170.
- Nei, M., and Kumer, S. 2000. *Molecular Evolution and Phylogenetics*. Oxford University press. London.
- Niruntrayakul, S. 2008. Gene Flow between Cultivated and Wild Rice. Thesis of Doctor of Philosophy in Agronomy, Chiang Mai University, Thailand.
- Oka, H.I. 1988. *Origin of Cultivated Rice*. Japan Scientific Society Press and Elsevier.
- Ouk *et al* 2006. Agrodiversity for in-situ conservation of local rice germplasm in and near its center of diversity: Cambodia, Lao and Thailand. Annual Report: Cambodia. 12 p.
- Panaud, O., X. Chen and S.R. McCouch. 1996. Development of microsatellite markers and characterization of sample sequence length polymorphism (SSLP) in rice (*Oryza sativa L.*). *Mol Gen Genet*. 252: 597-607.
- Pritchard, J.K., Stephens, M., and Donnelly, P. 2000. Inference of population structure using multilocus genotype data. *Genetics* 155: 943-959.
- Prommin, A. 2007. Segregation in F<sub>2</sub> Generation of Crosses Between Common Wild Rice (*O. rufipogon Griff.*) and Cultivated Rice (*Oryza sativa L.*). MS. Thesis, Faculty of Agriculture, Chiang Mai University.
- Punyalue, A. 2005. Characterization of common wild rice populations from main rice growing regions of Thailand. MS. Thesis, Faculty of Agriculture, Chiang Mai University.

- Robert, E.H., Craufurd, R.Q., and Le Cochet, F. 1961. Estimation of percentage of natural cross-pollination: experiment on rice. *Nature* 190: 1084-1085.
- Sano Y, Morishima H, Oka HI (1980) Intermediate perennial-annual populations of *Oryza perennis* found in Thailand and their evolutionary significance. *The Botanical magazine, Tokyo*, 93, 291-305.
- Song, Z.P., Lu, B.R., Zhu, Y.G., and Chen, J.K. 2002. Pollen competition between cultivated and wild rice species (*Oryza sativa* and *O. rufipogon*). *New Phytologist*, 153, 289-296 [68].
- Song, Z.P., Lu, B.R., Zhu, Y.G., and Chen, J.K. 2003. Gene flow cultivated rice to the wild species *Oryza rufipogon* under experimental field conditions. *New Phytologist* 157: 657-665.
- Song, Z.P., Xu, X., Wang, B., Chen, J.K., and Lu, B.R. 2003. Genetic diversity in the northernmost *Oryza rufipogon* populations estimated by SSR markers. *Theor Appl Genet* 107: 1492-1499.
- Sun, C.Q., Wang, X.K., Cai, H.W., Yoshimura, A., Doi, K., and Iwata, B. 1997. Genetic differentiation of nuclear DNA in common wild rice (*Oryza rufipogon* Griff.) and cultivated rice (*O. sativa* L.). *Journal of China Agricultural University*, 2(5):65-71.
- Temnykh, S., Park, W.D., Ayres, N.M., Cartinhour, S., Hauck, N., Lipovich, L., Cho, Y.G., Ishii, T., and McCouch, S.R. 2000. Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theoretical and applied genetics*. 100: 697-712.

- Temnykh, DeClerck, G., Lukashova, A., Cartinhour, S., Lipovich, L., and McCouch, S.R. 2001. Computational and experimental analysis of microsatellite in rice (*Oryza sativa* L.): frequency length variation transpose associations and genetic marker potential. *Genome research.* 11: 1441-1452.
- Tian F, Li DJ, Fu Q, Zhu ZF, Fu YC, Wang XK, Sun CQ (2006) Construction of introgression lines carrying wild rice (*Oryza rufipogon* Griff.) segments in cultivated rice (*O. sativa* L.) background and characterization of introgressed segments associated with yield-related traits. *Theor Appl Genet* 112:570–580.
- Vaughan, D.A. 1994. The wild relatives of rice: a genetic resources guide book. The International Rice Research Institute, Los Baños, Philipines. 137p.
- Vaughan, D.A., and Chang T.T. 1992. In situ conservation of rice genetic resources. *Economic Botany*, 46, 368-383.
- Vaughan, D.A., and Morishima, H. 2003. Biosystematics of the genus *Oryza*. In: W. Smith W. ed. Rice: Origin, History, Technology, and Production. New York, N.Y. (USA): John Wiley & Sons, Inc. p 27-65.
- Wongtamee, A. 2007. Evaluation of Exotic Genes Contamination from Crop Rice (*Oryza sativa* L.) in Common Wild Rice (*Oryza rufipogon* Griff.) Populations. MS. Thesis, Faculty of Agriculture, Chiang Mai University.
- Wongtamee, A. 2007. Genetic structure of common wild rice (*Oryza rufipogon* Griff.), populations from northeastern Thailand. *Journal of Agriculture*, 40<sup>th</sup> Anniversary, Faculty of Agriculture (CMU), ISSN 0857-0841, 83p.

- Wu, K.S., Tanksley. S.D. 1993. Abundance polymorphism and genetic mapping of microsatellite in rice. Mol Gen Genet. 241: 225-235.
- Xiao J, Li J, Grandillo S, Ahn SN, Yuan L, Tanksley SD, McCouch SR (1998) Identification of trait-improving quantitative trait loci alleles from a wild rice relative, *Oryza rufipogon*. Genetics 150:899–909.
- Xiong, L.Z., Liu, K.D., Dai, X.K., Xu, C.G., and Qifa Zhang. 1999. Identification of genetic factors controlling domestication-related traits of rice using an F<sub>2</sub> population of a cross between *Oryza sativa* and *O. rufipogon*. Theor Appl Genet, 98: 243-251.
- Yeh, F.C., Yang, R.C., and Boyle, T. 1999. POPGENE, the Microsoft Window-based User-Friendly Software for population Genetic Analysis of Co-dominant and Dominant Markers and Quantitative Traits. University of Alberta, Center for International Forestry Research, Alberta, Canada.
- Yuan, L.P. 1993. Advantages and constraints to the use of hybrid rice varieties. In: Wilson KJ, ed. Proceedings of the International Workshop on Apomixis in Rice. The Rockefeller Foundation and China National Center for Biotechnology Development, New York and Beijing: 1-4.