

## TABLE OF CONTENTS

|                                  | Page |
|----------------------------------|------|
| ACKNOWLEDGEMENT                  | iii  |
| ENGLISH ABSTRACT                 | iv   |
| THAI ABSTRACT                    | vi   |
| TABLE OF CONTENTS                | viii |
| LIST OF TABLES                   | ix   |
| LIST OF FIGURES                  | x    |
| CHAPTER 1 INTRODUCTION           | 1    |
| CHAPTER 2 LITERATURE REVIEWS     | 3    |
| CHAPTER 3 MATERIALS AND METHODS  | 14   |
| CHAPTER 4 RESULTS AND DISCUSSION | 19   |
| CHAPTER 5 CONCLUSION             | 40   |
| REFERENCES                       | 41   |
| APPENDIX                         | 48   |
| CURRICULUM VITAE                 | 62   |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
 Copyright© by Chiang Mai University  
 All rights reserved

## LIST OF TABLES


| Table  | Page |
|--|------|
| 2.1 Dielectric constant and Dielectric loss factor of selected foods   | 11   |
| 4.1 Effect of RF heat treatment on fungal seed-borne pathogens in barley seed evaluated by PDA method.         | 22   |
| 4.2 Effect of RF heat treatment on fungal seed-borne pathogens in barley seed evaluated by blotter method.     | 23   |
| 4.3 The total protein content of barley seeds treated with RF heat treatment at various temperatures and times | 37   |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
 Copyright© by Chiang Mai University  
 All rights reserved

## LIST OF FIGURES

| Figure |   | Page |
|--------|---|------|
| 2.1    | Structure of barley kernel  | 3    |
| 2.2    | The electromagnetic spectrum  | 7    |
| 2.3    | Schematic diagram of radio frequency application unit   | 8    |
| 2.4    | Space charge and dipolar polarization in an alternating electric field at radio frequencies   | 9    |
| 4.1    | The infection of seed-borne fungi in barley seed: <i>Alternaria</i> sp., <i>A. flavus</i> , <i>A. niger</i> , <i>Fusarium</i> sp., <i>Penicillium</i> sp. and <i>Rhizopus</i> sp. | 20   |
| 4.2    | Growth inhibition of seed-borne fungi in barley seeds, detected by PDA method, after RF heat treatment at different temperature and exposure time                                 | 21   |
| 4.3    | Growth inhibition of seed-borne fungi in barley seeds, detected by blotter method, after RF heat treatment at different temperature and exposure time                             | 24   |
| 4.4    | The moisture content of barley seeds treated with RF heat treatment at various temperatures and times   | 27   |
| 4.5    | The germination of barley seeds treated with RF heat treatment at various temperatures and times  | 29   |
| 4.6    | Different categories of barley seed viability characteristic determined by a tetrazolium test.  | 31   |
| 4.7    | The viability of barley seeds treated with RF heat treatment at various temperatures and times  | 32   |
| 4.8    | The germination after accelerated aging of barley seeds treated with RF heat treatment at various temperatures and times  | 33   |

| Figure |  | Page |
|--------|--|------|
| 4.9    | The germination index of barley seeds treated with RF heat treatment at various temperatures and times | 34   |
| 4.10   | The dehydrogenase activity of barley seeds treated RF heat treatment at various temperatures and times | 38   |



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่  
Copyright© by Chiang Mai University  
All rights reserved