

## REFERENCES

Adel, M.M., Sehnal, F. 2000. Azadirachtin potentiates the action of ecdysteroid agonist RH-2485 in *Spodoptera littoralis*. J. Insect Physiol. 46: 267-274.

Ascher KRS (1993). Non conventional insecticidal effects of pesticides available from the Neem tree, *Azadirachta indica*. Arch. Insect Biochem. Physiol. 22: 433-449.

Avtzis, D.N. 1989. Die wichtigsten Schadinsekten der griechischen Wälder. AFZ für Waldwirtschaft und Umweltsorge (4): 95.

Avtzis, N., 1986. Development of *Thaumetopoea pityocampa* Schiff. (Lepidoptera: Thaumetopoeidae) in relation to food consumption. For. Ecol. Manage. 15, 65–68.

Breuer, M., Devkota, B. Control of *Thaumetopoea pityocampa* (Den. & Schiff.) by extracts of *Melia azedarach* L. (Meliaceae), J. Appl. Entmol. 110 (1990) 128–135.

Breuer, M., Schmidt, G. H. Influence of a short period treatment with *Melia azedarach* extract on food intake and growth of the larvae of *Spodoptera frugiperda* (J.E. Smith) (Lep., Noctuidae), Z. PflKrankh. PflSchutz 102 (1995) 633– 654.

Breuer, M., De Loof, A., Meliaceous plant preparations as potential insecticides for control of the oak processionary moth, *Thaumetopoea processionea* (L.) (Lepidoptera: Thaumetopoeidae), Med. Fac. Landbouww. Univ. Gent. 63/2b (1998) 529–536.

Breuer M, De Loof A (1998). Laboratory trials with NeemAzal-T/S on the allergenic forest pest *Thaumetopoea processionea* (L.). In: Kleeberg H, Zebitz CPW (eds) Practice oriented results on use and production of Neem-ingredients and pheromones: Proceedings of the 8<sup>th</sup> workshop, Hohensolms, Germany, pp. 23-30.

Breuer, M., Kontzog, H. G., Guerrero, A., Camps, F. and de Loof, A., 2003. Field trials with the synthetic sex pheromone of the Oak processionary moth *Thaumetopoea processionea*. *Journal of Chemical Ecology*, Vol. 29, 2003: 2461 – 2468.

Breuer, M., De Loof, A. Efficacy of an enriched *Melia azedarach* L. fruit extract for insect control, in: H. Kleeberg, C.P.W. Zebitz (Eds.), Practice Oriented Results on Use and Production of Neem Ingredients and Pheromones, vol. VI, 2000, pp. 173–183.

Breuer, M., Hoste, B., de Loof, A. and Naqvi, S. N. H., 2003. Effect of *Melia azedarach* extract on the activity of NADPH-cytochrome c reductase and cholinesterase in insects. *Pesticide Biochemistry and Physiology* 76 (2003) 99–103.

Champagne, D. E., Koul, O., Isman, M. B., Scudder, M.G. E., Towers, G. H. N., 1992. Biological activity of limonoids from the Rutales. *Phytochemistry* 31, 377 – 394

Devakumar, C. and S. Dev, 1993. Chemistry. In: *Neem Research and Development*, Randhawa, N.S. and B.S. Parmar (Eds.). Society of Pesticide Science, New Delhi, India, pp: 63-96.

Dureja, P. and Johnson, S. 2000. Photodegradation of azadirachtin-A: A neem-based pesticide. Division of Agricultural Chemicals, Indian Agricultural Research Institute, New Delhi 110 012, India.

Dwyer, J. F., Schroeder, H. W., Gobster, P. H., 1991. The significance of urban trees and forests: Towards a deeper understanding of values.

Ermel, K.; Pahlich, E.; Schmutterer, H., 1984: Comparison of the azadirachtin content of neem seeds for ecotypes of Asian and African origin. In: *Natural Pesticides from the Neem Tree and Other Tropical Plants*. Ed. by Schmutterer, H.; Ascher, K. R. S. Eschborn: GTZ, 91±93.

Ermel, K.; Pahlich, E.; Schmutterer, H., 1987: Azadirachtin content of neem kernels from different geographical locations, and its dependence on temperature, relative humidity and light. In: *Natural Pesticides from the Neem Tree and Other Tropical Plants*. Ed. By Schmutterer, H.; Ascher, K. R. S. Eschborn: GTZ, 171±184.

Fagoonee, I. 1984. Effects of Azadirachtin and of a neem extract on food utilization by *Crocidolomia binotalis*. Proc. 2<sup>nd</sup> Int. Neem Conf. (Rauischholzhausen, 1983), 211 – 224.

Ferenczy, A., Eppich, B., Varga, R. D., Biro, I., Kovacs, A., Petranyi, G., Hirka, A., Szaboki, CS., Isepy, I., Priszter, SZ., Türei, D., Gimesi, L., Garamvölgyi, A., Homorodi, R., Hufnagel, L., 2010. Comparative analysis of the relationship between phenological phenomena and meteorological indicators based on insect and plant monitoring. *Applied Ecology and Environmental Research* 8(4): 367-376.

Grahn, P. and Stigsdotter, U.A. (2003) 'Landscape planning and stress', in *Urban Forestry and Urban Greening*, 2, 1-18

Gregorová, B., Černý, K., Holub, V., Strnadová, V., 2010. Effects of climatic factors and air pollution on damage of London plane (*Platanus hispanica* Mill.) Hort. Sci. (Prague) Vol. 37, 2010, No. 3: 109–117.

Hartig, T., Böök, A., Garvill, J., Olsson, T., & Garling, T. (1996). Environmental influences on psychological restoration. *Scandinavian Journal of Psychology*, 37, 378–393.

Hough-Goldstein, J. and C.B. Keil. 1991. Prospects for integrated control of the Colorado potato beetle (Coleoptera:Chrysomelidae) using *Perillus bioculatus* (Hemiptera:Pentatomidae) and various pesticides. *J. Econ. Entomol.* 84: 1645-1651.

Isman, M.B., 1999. Neem and Related Natural Products. In: *Biopesticides: Use and Delivery*. Hall, F.R. and J.J. Menn (Eds.). Humana Totowa, New Jersey, pp: 139-153.

Jacobson, M. 1989. Botanical pesticides. Past, present and future. In Arnason, J. T., Philogéne, B. J. R., Morand, P. (Eds.) *Insecticides of Plant Origin*. American Chemical Society, Symp. Ser. No. 387, pp. 1 – 10, Washington, DC.

Jactel, H., Menassieu, P., Ve´tillard, F., Barthe´le´my, B., Piou, D., Fre´rot, B., Rousselet, J., Goussard, F., Branco, M., Battisti, A., 2006. Population monitoring of the pine processionary moth (*Lepidoptera: Thaumetopoeidae*) with pheromone-baited traps. *Forest Ecology and Management* 235 (2006) 96–106.

Kaethner, M. (1991) Untersuchung über die Eignung von Niemsamenprodukten zur Bekämpfung des Waldmaikäfers *Melolontha hippocastani* F. und des Feldmaikäfers *M. melolontha* L. (Col., Scarabaeidae).

Ketkar, C. M.; Ketkar, M. S. (1993): Different Uses of Neem S. 1-12; Proceedings of 2<sup>nd</sup> Workshop: Practice Oriented Results on Use and Production of Neem-Ingredients and Pheromones (ed. h. Kleeberg), Druck & Graphic, Giessen.

Kleeberg, H. and Zebitz, C.P.W. (Eds.) (1997): Practice oriented results on use and production of neem ingredients and pheromones: Proceedings of the fifth workshop. Wetzlar, 22 nd - 25 th January 1996, Giessen. 81-92.

Koul, O. and Isman, M. B. 1991. Effects of azadirachtin on the dietary utilization and development of the variegated cutworm, *Peridroma saucia*. *J. Insect Physiol.*, 37: 591-598.

Koul, O. and Wahab, S.: “Neem: Today and in the New Millennium,” Springer, Netherlands, 2004.

Kweon, B.S., Sullivan, W.C. & Wiley, A.R. (1998). Green common spaces and the social integration of inner-city older adults. *Environment and Behavior* 30(6), 832-858.

Ladanyi, M. and Hufnagel, L., 2006. The effect of climate change on the population of sycamore lace bug *Corythucha ciliata* (SAY) (Heteroptera: Tingidae) based on a simulation model with phenological response. *Applied Ecology and Environmental Research* 4(2): 85-112.

Maceljski, M. (1986): New development in the status of *Corythucha ciliata* in Europe. II. Meeting "Integr. control of *C. ciliata*", Padova, 1985, Bull. IOBC/WPRS IX/1. Bruxelles.

Maceljski, M. (1986): Current status of *Corythucha ciliata* in Europe. - Europ. Plant Prot. Organ. Conference, Nancy, 1986. Bull. OEPP, 16. : 621-624

McPherson, E.G., and J.R. Simpson. 2002. A comparison of municipal forest benefits and costs in Modesto and Santa Monica, California, USA. *Urban For. Urban Green.* 1:61-74.

Mordue, A.J., Simmonds, M.S.J., Ley, S.V., Blaney, W.M., Mordue, W., Nasiruddin *et al.* (1998) Action of azadirachtin, a plant allelochemical, against insects. *Pestic. Sci.* 54:277-284.

Mordue (Luntz) A. J., Evans K. A. and Charlet M. (1986) Azadirachtin, ecdysteroids and ecdysis in *Locusta migratoria*. *Comp. Biochem. Physiol.* 85c, 297-301

Mortimer, M.J., and Kane, B. C. P. 2004. Hazard tree law in the United States. *Urban Forestry and Urban Greening* 2:208–215.

Musolin, D. L., 2005. Insects in a warmer world: ecological, physiological and life-history responses of true bugs (*Heteroptera*) to climate change. *Global Change Biology* (2007) 13, 1565–1585.

Netherer, S., Schopf, A., 2009. Potential effects of climate change on insect herbivores in European forests—General aspects and the pine processionary moth as specific example. *Forest Ecology and Management* 259 (2010) 831–838.

Öszi, B., Ladanyi, M., Hufnagel, L., 2005. Population dynamics of the sycamore lace bug *Corythucha ciliata* (SAY) (Heteroptera: Tingidae) in Hungary. *Applied Ecology and Environmental Research* 4(1): 135-150.

Parmal, B.S. 1987: An overview of neem research and use in India during the years 1983-1986. In : *Natural Pesticides from Neem Tree and other Tropical Plants*. Proc. of the 3rd Int. Neem Conf. Nairobi, Kenya, 1986, 55-80.

Pavela, R. and Kalinkin, V. M. 2010. Systemic applications of Neem in the control of *Corythucha ciliate* Say., a Pest of *Platanus hispanica* Mill. Crop Research Institute, Drnovská 507, 161 06 Prague 6, Czech Republic

Quero, C., Bau, J., Guerrero, A., Breuer, M., de Loof, A., Kontzog, H. G. and Camps, F., 2003. Sex pheromone of the Oak processionary moth *Thaumetopoea processionea*. Identification and biological activity. *J. Agric. Food Chem.* 2003, 51: 2987-2991.

Randhawa, N. S. & Parmar B. S. (eds.) (1993): Neem – Research and Development, Soc. Of Pesticide Sci., India, New Delhi

Rembold, H. and Sieber, K.P. (1981): Inhibition of ovogenesis and ovarian ecdysteroid synthesis by azadirachtin in *Locusta migratoria migratorioides*. Z. Naturf., 36:466 - 469.

Rembold, H., Sharma, G.K. and Czoppelt, C., (Eds. Schmutterer, H., Ascher, K.R.S., Rembold, H.), *Natural Pesticides from the Neem Tree (Azadirachta indica)*, GTZ. Eschborn, 1982, Germany, 121–128.

Rembold, H.; Forster, H.; Czoppelt, C.H.; Rao, P.J. and Sieber, K.P. (1984): The azadirachtin, a group of insect growth regulator from the neem tree. In “Natural Pesticides from the neem tree (*Azadirachta indica*. A. Juss) and other Tropical Plants”

Rembold, H., Uhl, M., Müller, Th. 1987: Effect of Azadirachtin A on hormone titers during the gonadotropic cycle of *Locusta migratoria*. In : Natural Pesticides from Neem Tree and other Tropical Plants. Proc. of 3rd Int. Neem Conf. (Nairobi, Kenya, 1986), 289-298.

Rojht, H., Mesko, A., Vidrih, M., Trdan, S., 2009. Insecticidal activity of four different substances against larvae and adults of sycamore lace bug (*Corythucha ciliata* [Say], Heteroptera, Tingidae) Acta agriculturae Slovenica, 93 – 2009.

Rosell, G., Quero, C., Coll, J. and Guerrero, A., 2008. Biorational insecticides in pest management. J. Pestic. Sci., 33(2), 103–121 (2008).



Ruch, B., Kleeberg, H. 2000: Abschätzung des Rückstandverhaltens von NeemAzal-T/S aus Analysen der Leitsubstanz Azadirachtin A. In: Pflanzen-schutz im ökologischen Landbau, 4. Fachgespräch, Darmstadt, Juni 2000, (Kühne, ed), Saphir Verlag, S. 84-88.

Schenke, D., Knutzen, F., Jäckel, B., Doobe, G., Hilfert, G., 2009. Aufnahme von Dimethoat in Blätter von Spitzahorn, Linde und Kastanie nach Stammbehandlung mit Baumpflaster. 57. Deutsche Pflanzenschutztagung am 06.-09. September 2010, Berlin. PO-208

Schlüter, U. und Schulz, W. D. 1984. Structural damages caused by neem in *Epilachna varivestis*; A summary of histological and ultrastructural data. I. Tissues affected in larvae. Proc. 2<sup>nd</sup> Int. Neem Conf. (Rauischholzhausen, 1983), 227 – 236.

Schmutterer, H., and Ascher, K. R. S., (eds.): “Natural Pesticides from the Neem Tree (*Azadirachta indica* A. Juss.) and Other Tropical Plants,” Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ), Eschborn, Germany, 1984.

Schmutterer, H.; Zebitz, C.P.W. 1984: Effect of in ethanolic extracts from seeds of single neem trees of African and Asian origin, on *Epilachna variivestis* and *Aedes aegypti*. In: n: Natural Pesticides from the Neem Tree and other tropical Plants. Proc. of the 2nd Int. Neem Conf. (Rauischholzhausen,1983), 83-90.

Schmutterer, H. 1987: Fecundity reducing and sterilising effects of neem seed kernel extracts in the Colorado potato beetle, *Leptinotarsa decemlineata*. In: Natural Pesticides from the Neem Tree and other tropical Plants. Proc. of the 3rd Int. Neem Conf (Nairobi, Kenya, 1986), 351-260.

Schmutterer, H., and Ascher, K. R. S., (eds.): "Natural Pesticides from the Neem Tree (*Azadirachta indica* A. Juss.) and Other Tropical Plants," Deutsche Gesellschaft für Technische Zusammenarbeit (GTZ), Eschborn, Germany, 1987.

Schweizer, C., 2005. Bekämpfung der Rosskastanienminiermotten (*Cameraria ohridella*) mit NeemAzal. Andermatt BIOCONTROL AG, ag-journal 2005.

Schmutterer, H., Ascher, K.R.S.; Rembold, H. 1981: Natural pesticides from the neem tree (*Azadirachta indica* A. Juss). In: Natural Pesticides from the Neem Tree and other Tropical Plants. Proc. 1st Int. Neem Conf. (Rottach-Egern, 1980), 297 pp.

Schmutterer, H. 1990. Properties and potential of natural pesticides from the neem tree, *Azadirachta indica*. Annu. Rev. Entomol. 35: 271-297.

Schmutterer, H., 1995. The neem tree, *Azadirachta indica* A. Juss and other meliaceous plants: Sources of unique natural products for integrated pest management, medicine, industry and other purposes. VCH Publishers Inc., New York, NY (USA).

Schweizer, C., 2006. Rosskastanienminiermotten können mit NeemAzal-T/S bekämpft werden. Andermatt BIOCONTROL AG, ag-journal 2006.

Sieber, K. P. and Rembold H., 1983. The effects of Azadirachtin on the endocrine control of moulting in *Locusta migratoria*. J. Insect Physiol., 29: 523 – 527

- Szőcs, G., Balázs, K., Nagy, Z., Tóth,Cs., Nemestóthy, K., Demeter, T., Ujváry,I., Hummel, E., 2007. Alien Arthropods in South East Europe – crossroad of three continents *Cameraria Ohridella*: Do we know all about it? University of Forestry 2007, Sofia, Bulgaria.
- Tanzubil, P. B. and McCaffery, A. R. 1990. Effects of Azadirachtin and aqueous neem seed extracts on survival, growth and development of the African armyworm, *Spodoptera exempta*. – Crop Protection 9, 383 – 386.
- Unal, S. and Akkuzu E., 2009. Larvaecidal effects of azadirachtin on the pine processionary moth. Faculty of Forestry, Kastamonu University, 37100, Kastamonu-Turkey. African Journal of Biotechnology Vol. 8 (19), pp. 5128-5131, 2009.
- Wilps, H. 1987. Growth and adult molting of larvae and pupae of the blowfly *Phorria terrae-novae* in relationship to Azadirachtin concentrations. Proc. 3<sup>rd</sup> Int. Neem Conf. (Nairobi, 1986), 299-314.