

7. REFERENCES

- Asami T, Nakano T, Sakai K. Study on the relation between renal tubular disorders and glomerular dysfunction in the early phase of insulin-dependent diabetes mellitus in children. *Nippon Jinzo Gakkai Shi* 1992; 34: 57-63.
- Baker JR, Metcalf PA, Johnson RN, *et al.* Use of protein-based standards in automated colorimetric determinations of fructosamine in serum. *Clin Chem* 1985; 31: 1550-4.
- Bishop ML, *et al.* Basic Physiology, Analytical Procedures and Clinical Correlations. *Clinical Chemistry*, 2nd edition 1992; 460-3.
- Chanarat N, Suksriwong S. Microalbuminuria in diabetes mellitus, Presented at the 21st World Congress of Medical Technology, Hong kong, 25-29 July, 1994.
- Chesham J, Andertons SW, Kingdon CFM. Rapid, Competitive enzyme immunoassay for albumin in urine. *Clin Chem* 1986; 32: 669-71.
- Fielding BA, Price DA, Honlton CA. Enzyme immunoassay for urinary albumin. *Clin Chem* 1983; 29: 355-7.
- Furuhata N, Shiba K, Nara N. N-acetyl-beta-D-glucosaminidase . *Nippon Rinsho*. 1995; 63: 1267-76.
- Gyre WL. Comparison of several methods for semiquantitative determination of urinary protein. *Clin Chem* 1977; 23: 876-9.
- Harmoinen A, Vworimen P, Jokela H. Turbidimetric measurement of microalbuminuria. *Clin Chim Acta* 1987; 166 : 85-9.
- Hirai M, Nakano H, Oba K, *et al.* Microalbumin and N-acetyl-beta-D-glucosaminidase in random spot urine samples as predictors of diabetic

- nephropathy in the elderly non-insulin dependent diabetic patients. *Nippon Ika Daigaku Zasshi* 1997; 64: 518-25.
- Horak E, Hopfer SM, Sunderman FW. Spectrophotometric assay for urinary N-acetyl-beta-D-glucosaminidase activity. *Clin Chem* 1981; 27: 1180-5.
- Hsiao PH, Tsai WS, Tsai WY, *et al.* Urinary N-acetyl-beta-D-glucosaminidase activity in children with insulin-dependent diabetes mellitus. *Am J Nephrol* 1996; 16: 300-3.
- Hultberg B, Ockerman PA. Artificial substrates in the assay of acid glycosidases. *Clin Chim Acta* 1972; 39: 49-58.
- Ikenaga H, Suzuki H, Ishii N, *et al.* Enzymuria in non-insulin dependent diabetic patients : signs of tubular cell dysfunction. *Clin Sci Colch* 1993; 84: 469-75.
- Itoh Y, Numata Y, Morita A, *et al.* Varies value of urinary N-acetyl-beta-D-glucosaminidase isoenzyme B in male of reproductive age. *Kidney Intern* 1994; 46: 38-42.
- Jones AP, Lock S, Griffiths KD. Urinary N-acetyl-beta-glucosaminidase activity in type I diabetes mellitus. *Ann Clin Biochem* 1995: 58- 62.
- Jung K, Diego J, Scholz D, *et al.* Urinary enzyme excretion by renal transplant recipients in relation to interval after transplantation. *Clin Chem* 1982; 8: 1762-4
- Kanauchi M, Ishihara K, *et al.* Glomerular lesions in patients with non-insulin-dependent diabetes mellitus and microalbuminuria. *Intern Med* 1993; 32: 753-7.
- Kaplan A, Szabo LL. *Clinical Chemistry : Interpretation and Laboratory* 2nd edition, 1983, Philadelphia : Lea & Febiger, 1023-74.

- Killingsworth LM. Clinical application of protein determinations in biological fluids other than blood. *Clin Chem* 1982; 28: 1093-107.
- Klenk DC, Hermanson GJ, Krohn RI, *et al.* Determination of glycosylated hemoglobin by affinity chromatography : comparison with colorimetric and ion exchange method and effects of common interferences. *Clin Chem* 1982; 28: 2088-94.
- Knoll E, Wisser H, Rautenstrauch H. Comparative study of the diagnostic value of disc electrophoresis of urine proteins and measurement of the excretion of N-acetyl- β -D-glucosaminidase for the detection of renal tubule damage in chronic polyarthritis. *J Clin Chem Clin Bioc* 1980; 18: 53-8.
- Kumar P, Clark M. Diabetes Mellitus and Other Disorders of Metabolism. *Clinical Medicine* : 3rd edition 1994.
- Kunin CM, Chesney RW, Craig WA, *et al.* Enzymuria as a marker of renal injury and disease : Study of N-acetyl- β -D-glucosaminidase in the general population and in patients with renal disease. *Pediatrics* 1978; 62: 751-60.
- Lockwood TD, Bosmann HB. The use of urinary N-acetyl- β -D-glucosaminidase in human toxicology. II. Partial biochemical characterization and excretion in humans and release from the isolated perfused rat kidney. *Toxicol Appl Pharm* 1979; 49: 323-36.
- Lorini R, Scaramuzza A, *et al.* Increased urinary N-acetyl-beta-glucosaminidase (NAG) excretion in young insulin-dependent diabetic patients. *Diabetes Res Clin Pract* 1995; 29: 99-105.

- Makise J, Salto E, Obuchi M, *et al.* Kinetic rate assay of urinary N-acetyl-beta-D-glucosaminidase in urine with 2-chloro-4-nitrophenyl-N-acetyl- β -D-glucosaminidase as substrate. *Clin Chem* 1988; 34: 2140-3.
- Mancini G, Carbonara AO, Heremans IF. Immunochemical quantitation of antigen by single radial immunodiffusion. *Imm Chem* 1965; 2: 235-54.
- Martin D, Walton C, Chapman C, Bodansky HJ, Stickland MH. Increased urinary excretion of transferrin in children with type I diabetes mellitus. *Diabet Med* 1990; 7: 35-40.
- Mocan Z, Erem C, *et al.* Urinary beta-2-microglobulin levels and urinary N-acetyl-beta-D-glucosaminidase enzymes activities in early diagnosis of non-insulin-dependent diabetes mellitus nephropathy. *Diabetes Res* 1994; 26: 101-7.
- Morganson CE, Christensen CK. Predicting diabetic nephropathy in insulin-dependent patients. *N Engl J Med* 1984; 311: 89-93.
- Morganson CE, Chachati A, Christensen CK, *et al.* Microalbuminuria : an early marker of renal involvement in diabetes. *Uremia Invest* 1986; 9: 85-95.
- Nakamura S. Clinical evaluation of urinary alanine aminopeptidase in the patients with diabetes mellitus comparison among AAP, microalbumin and N-acetyl-beta-D-glucosaminidase. *Hokkaido Igaku Zasshi* 1991; 66: 522-33.
- National Working group for Control of Diabetes. Diabetes mellitus in Thailand 1987; Review and prospective.1988, Ministry of Health, Bangkok 14-27.

- Noto A, Nagawa Y, Mori S, *et al.* Simple rapid spectrophotometry of urinary N-acetyl-beta-D-glucosaminidase, with use of new chromogenic substrate. *Clin Chem* 1983; 29: 1713-6.
- Noto A, Kunihiro N, Kazuyuki S, Tsuomu S. Method for determination of NAGase and reagent therefor. *Eur Pat Application* 1988:88109201 .g.
- Paoli C, Bardelli F, Tarli P, *et al.* A simple latex agglutination test for urinary albumin screening. *Clin Chim Acta* 1987; 166: 67-71.
- Pocsi I, Csathy L, Olah VA, Price RG. Assay of urinary N-acetyl-beta-D-glucosaminidase in urine from neonates : comparison of two new colorimetric methods using MNP-GlcNAc and VRA-GlcNAc as substrates. *Ann Clin Biochem* 1992; 29: 292-5.
- Polnamin D, Determination of Urinary N-acetyl-beta-D-glucosaminidase (NAG) in diabetics. Term paper, Faculty of Associated Medical Sciences, 1995.
- Pongsomboon S, Chanarat N. Colorimetric determination of microalbumin in urine. Term Paper, Faculty of Associated Medical Sciences, 1992.
- Powell SC, Scaro J, Wilson E and Shihabi ZK. Assay of urinary N-acetyl-beta-D-glucosaminidase in a centrifugal analyzer. *Clin Chem* 1983; 29: 1717-9.
- Price RG. Urinary enzymes, nephrotoxicity and renal disease. *Toxicology* 1982; 23: 99-134.
- Price RG. Urinary N-acetyl- β -D-glucosaminidase (NAG) as an indicator of renal disease. *Curr Probl Clin Biochem* 1979; 9: 150-63.

- Rowe DJF, Dawney A, Watts GF. Microalbuminuria in diabetes mellitus : review and recommendations for the measurement of albumin in urine. *Ann Clin Biochem* 1990; 27: 297-312.
- Ruston R, Costigan M, *et al.* Proteinuria and renal tubular damage : urinary N-acetyl-beta-D-glucosaminidase and isozymes in dissimilar renal disease. *Am J Nephrol* 1998; 18: 179-85.
- Saitoh A, Tomino Y, *et al.* Determination of glomerular NAG fraction in urine samples of patients with diabetic nephropathy using the NAG test kit. *Nephron* 1992; 61: 114.
- Sanchez HMC, Mateo CJ, *et al.* Influence of glycemic blood glucose control and incipient diabetic nephropathy on the urinary excretion of NAG in diabetes mellitus. *Ann Med Intern* 1995; 12: 216-20
- Schneider RE, Rockstroh JK, Munch HG, *et al.* Elevated serum activity of N- acetyl- beta-glucosaminidase in essential hypertension : : diagnostic value and reversal to normal values after antihypertensive therapy. *Am J Kidney Dis* 1991; 18: 38-48.
- Schosinsky KH, Vargas M, Esquivel AL, Chavarria MA. Simple spectrophotometric determination of urinary albumin by dye binding with use of bromphenol blue. *Clin Chem* 1987; 3: 223-6.
- Sheth KJ, Good TA. Urinary acidic hydrolase in renal diseases in children. *Clin Nephrol* 1978; 10: 151-7.
- Shrivastava KS, Awasthi YC, Yoshida A. Studies on human- β -D-N-acetylhexosaminidase. I. Purification and properties. *J Biol Chem* 1974; 249: 2043-8.
- Silver AC, Dawney A, Landon J, Cattell WR. Immunoassays for low concentrations of albumin in urine. *Clin Chem* 1986; 32: 1303-6.

- Silver AC, Dawney A, Landon J. Specimen preparation for assay of albumin in urine. *Clin Chem* 1987; 33: 199-200.
- Skrha J, Sperl M, *et al.* Albuminuria and N-acetyl-beta-D-glucosaminidase activity in a prospective study of type I diabetics. *Vnitr Lek* 1993; 39: 699-705.
- Stolarek I, Howey JEA, Fraser CG. Biological variation of urinary N-acetyl- β -D-glucosaminidase : practical and clinical implications. *Clin Chem* 1989; 35: 560-3.
- Tanami J. Diabetic control and complications in diabetic clinic, Lerdsin Hospital. *Bull Dept Med Serv* 1995; 20: 147-54.
- The Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. Report on the Expert Committee on the Diagnosis and Classification of diabetes mellitus. *Diabetes Care* 1997; 20: 1183-97.
- Townsend JC. Effect of storage temperature on the precipitation of albumin from urine. *Clin Chem* 1986; 32: 1986-7.
- Viberti GC, Jarrett RJ, *et al.* Microalbuminuria as a predictor of clinical nephropathy in insulin-dependent diabetes mellitus. *Lancet* 1982; 2:430-2.
- Wellwood JM, Ellis BG, Price RG, *et al.* Urinary N-acetyl- β -D-glucosaminidase activities in patients with renal disease. *Br Med J* 1975; 3: 408-11.
- Whiting PH, Ross IS, Borthwick L. Serum and urine N-acetyl-beta-D-glucosaminidase in diabetes on diagnosis and subsequent treatment and stable insulin dependent diabetics. *Clin Chim Acta* 1979a; 92: 459-64.

- Whiting PH, Ross IS, Borthwick LJ. N-acetyl-beta-D-glucosaminidase levels and diabetic microangiopathy. *Clin Chim Acta* 1979b; 97: 191-5.
- Widstam AU, Berg U. Urinary protein excretion and renal function in young people with diabetes mellitus. *Nephrol Dial Transplant* 1992; 7: 487-92.
- Woo J, Floyd M, Cannon DC, Kahan B. Radioimmunoassay for urinary albumin. *Clin Chem* 1978; 1464-7.
- Wu JT. Review of Diabetes : Identification of markers for early detection, glycaemic control and monitoring clinical complications. *J Clin Lab Anal* 1993; 7: 293-300.
- Wu JT. Advanced Glycosylation End Products : A new disease marker for diabetes and aging. *J Clin Lab Anal* 1993; 7: 252-5.
- Yagame M, Suzuki D, *et al.* Urinary albumin fragments as a new clinical parameter for the early detection of diabetic nephropathy. *Intern Med* 1995; 15: 463-8.
- Yagi T, Hisada R, Shibata H. 3,4-Dinitrophenyl N-acetyl- β -D-glucosaminidase, a synthetic substrate for direct spectrophotometric assay of N-acetyl- β -D-glucosaminidase. *Anal Biochem* 1989; 183: 245-9.
- Yaqoob M, McClelland P, Patrick AW, *et al.* Tubular damage in microalbumic patients with primary glomerulonephritis and diabetic nephropathy. *Ren Fail* 1995; 17: 43-9.
- Yuen CT, Corbett CRP, Kind PRN, Thompson AE, Price RG. Isoenzymes of urinary N-acetyl- β -D-glucosaminidase (NAG) in patients with renal transplants. *Clin Chim Acta* 1987; 164: 339-50.