

TABLE OF CONTENTS

	Page
APPROVAL PAGE	ii
ACKNOWLEDGEMENTS	iii
ABSTRACT	iv
LIST OF TABLES	x
LIST OF ILLUSTRATIONS	xi
ABBREVIATIONS	xiv
CHAPTER I: INTRODUCTION	1
CHAPTER II: LITERATURE REVIEWS	5
1. Virology	5
2. Viral growth cycle	6
3. Host immune response	8
4. Immune evasion of HCMV	11
5. Persistence and latency	12
6. Pathology	12
7. Epidemiology	13
8. Laboratory diagnosis	15
9. Trends in laboratory diagnosis of Cytomegalovirus infection	21
10. Development of PCR technique for HCMV detection	22
11. Prevention of contamination in PCR process	28
CHAPTER III: RESEARCH DESIGN, MATERIALS AND METHODS	31
1. Research design	31
2. Materials and methods	34
2.1 Subjects	34
2.2 Oligonucleotide primers and probe	35

2.3 Specimen collection and preparation	36
2.4 Cloning of the HCMV MIE gene fragment	38
2.5 Preparation of standard DNA control	42
2.6 Quantitation of the DNA controls	44
2.7 Dot blot hybridization technique	46
2.8 Optimization of the nested PCR and duplex PCR condition	48
2.9 The HCMV DNA detection in the clinical specimens by the optimized duplex PCR	52
2.10 Data analysis	52
CHAPTER IV: RESULTS	54
1. Preparation of standard controls	54
2. Determination of conventional PCR assay sensitivity	60
3. Optimization of the PCR conditions	62
4. Detection of HCMV DNA in patients	78
4.1 Retinitis patients	78
4.2 Renal transplant patients	85
CHAPTER V: DISCUSSION AND CONCLUSION	90
REFERENCES	98
APPENDIX	110
CIRRICULUM VITAE	117

LIST OF TABLES

Table	Page
1 Vertical transmission of human CMV infection.	2
2 Rates of HCMV infection and disease in kidney and liver transplant patients, according to type of infection.	3
3 Immunogenic proteins of HCMV.	10
4 Interpretation of laboratory tests for HCMV infection and immunity.	20
5 HCMV MIE PCR Primers and oligonucleotide probe information.	35
6 The conditions of conventional and optimized nested PCR for amplification of the HCMV MIE gene fragment and beta-globin gene	76
7 The results of HCMV DNA detection by conventional and optimized duplex nested PCR in HCMV suspected retinitis patients	83
8 Comparison of positive results in conjunctival scraping, aqueous and vitreous humor samples after HCMV DNA detection by conventional and optimized duplex nested PCR.	84

LIST OF ILLUSTRATIONS

Figure	Page
1 HCMV genome structure	6
2 Human cytomegalovirus genome structure. The HCMV MIE gene is located between 165-178 kbp of EcoRI fragment. It is composed of 4 exon region (Exon1, Exon2, Exon3, and Exon4).	8
3 The schematic diagram of the research design in this study.	33
4 Structure of the human eye	34
5 The locations of primers (FP, RP, NFP, NRP) and probe on HCMV MIE gene.	36
6 The amplification of the HCMV MIE gene fragment by the conventional first round PCR.	54
7 The amplification of the HCMV MIE gene fragment by the conventional second round PCR.	55
8 The amplification of HCMV MIE gene fragment from colonies of transformed <i>E. coli</i> by the conventional first round PCR.	56
9 The circular plasmid with the HCMV MIE gene fragment insert was purified by miniprep technique.	57
10 The amplification of the HCMV MIE gene fragment from miniprep plasmid DNA by the conventional first round PCR.	58
11 The amplification of a beta-globin gene from 1 to 500 ng of human genomic DNA controls by the conventional PCR assay.	60
12 The sensitivity of the conventional first round PCR assay.	61
13 The sensitivity of the conventional second round PCR assay.	62
14 The amplification of 10^6 copies of the HCMV MIE gene fragment (351 bp) in the presence of various $MgCl_2$ concentrations by the first round PCR.	64

15	The amplification of the HCMV MIE gene fragment (170 bp) in the presence of various $MgCl_2$ concentrations by the second round PCR.	64
16	Comparison in the first round HCMV MIE gene (10^6 copies) amplification efficiency after adding various dNTP concentrations.	65
17	Comparison in the second round HCMV MIE gene (10^6 copies) amplification efficiency after adding various dNTP concentrations.	66
18	The amplification of a ten-fold concentration of 0.1 to 10^7 copies of the HCMV MIE gene fragment by the optimized first round PCR.	67
19	The amplification of a ten-fold concentration of 0.1 to 10^7 copies of the HCMV MIE gene fragment by the optimized second round PCR.	68
20	The detection of optimized second round PCR product of 1 to 10^7 HCMV DNA by dot blot hybridization.	69
21	The amplification of 50 ng human genomic DNA combined to 10^4 , 10^5 and 10^6 copies of HCMV MIE gene fragment in the optimized first round duplex PCR.	71
22	The amplification of 100 ng human genomic DNA combined with ten-fold concentration of 1 to 10^7 copies of HCMV MIE gene fragment in the optimized first round duplex PCR.	72
23	The amplification of 500 ng of human genomic DNA combined with a ten-fold concentration of 1 to 10^8 copies of the HCMV MIE gene fragment in the optimized first round duplex PCR.	73
24	The optimized first round PCR amplification of 1, 10, 20, 30, 40 and 50 ng beta-globin gene in the presence of 10^5 HCMV MIE gene fragment.	74
25	PCR products achieved when different primer concentration of beta-globin gene and HCMV MIE gene primers were used in the duplex PCR assay.	75
26	The amplification of a ten-fold concentration of 1 to 10^7 copies of the HCMV MIE gene fragment by the optimized second round duplex nested PCR.	77
27	The amplification of HCMV DNA in ocular samples by the first round of conventional PCR.	79

28	The amplification of HCMV DNA in ocular samples by the second round of conventional PCR.	79
29	The amplification of HCMV DNA in ocular samples by the optimized first round duplex nested PCR.	81
30	The amplification of HCMV DNA in ocular samples by the optimized second round duplex nested PCR.	82
31	Comparison of the positive results of HCMV DNA amplification in ocular samples by the conventional and optimized duplex nested PCR.	84
32	Percentage of HCMV DNA positive specimens in plasma and PBMCs at 2 week intervals after transplantation.	88
33	The amplification of HCMV DNA from PBMCs and plasma samples in transplantation patients by the optimized first round duplex nested PCR.	89
34	The amplification of HCMV DNA from PBMCs (lane 1-6) and plasma samples (lane 7-9) in transplantation patients by the optimized second round duplex nested PCR.	89

ABBREVIATIONS

α	alpha
β	beta
γ	gamma
$^{\circ}\text{C}$	degree celsius
μl	microliter
A	adenine
C	cytosine
G	guanine
T	thymine
dATP	deoxyadenosine triphosphate
dCTP	deoxycytosine triphosphate
dGTP	deoxyguanine triphosphate
DNA	deoxyribonucleic acid
dNTP	deoxyribonucleotide triphosphate
dTTP	deoxythymine triphosphate
ELISA	enzyme linked immunosorbent assay
IgA	immunoglobulin A
IgG	immunoglobulin G
IgM	immunoglobulin M
Kb	kilobasepair
LB medium	Luria-Bertani medium
M	molarity
mg	milligram
MgCl_2	magnesium chloride
min	minute
ml	milliliter
mM	millimolar

mRNA	messenger ribonucleic acid
MW	molecular weight
N	normality
ng	nanogram
nm	nanometer
O.D.	Optical Density
PBMC	peripheral blood mononuclear cells
PCR	Polymerase Chain Reaction
pmol	picomole
RNA	ribonucleic acid
Tag	<i>Thermus aquaticus</i>

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved