

TABLE OF CONTENTS

CONTENT	PAGE
ACKNOWLEDGEMENT	iii
ABSTRACT ENGLISH	iv
ABSTRACT THAI	vi
TABLE OF CONTENTS	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
ABBREVIATIONS	xvi
CHAPTER I: INTRODUCTION	1
1.1 Statement and significance of the problem	1
1.2 Literature reviews	3
1.2.1 Introduction to polyclonal and monoclonal antibody	3
1.2.2 Monoclonal antibody	5
1.2.2.1 Hybridoma technology	6
1.2.2.1.1 Immunization protocol	17
1.2.2.1.2 Developing the screening procedure	18
1.2.2.1.3 Producing hybridomas	19
1.2.2.1.3.1 Single cell cloning	19
1.2.3 Feeder cells	22
1.2.4 Conditioned medium	23
1.3 Objectives	25
CHAPTER II: MATERIALS AND METHODS	26

2.1 Chemicals, antibodies, cell lines and instruments used in this study	26
are shown in Appendix A-C.	
2.2 Mycoplasma Detection	26
2.2.1 DNA preparation	26
2.2.2 Detection of mycoplasma contamination by PCR	27
2.3 Preparation of conditioned media	29
2.4 Study the utilization of conditioned media for hybridoma single cell cloning	29
2.4.1 Single cell cloning using stable hybridoma line	29
2.4.2 Single cell cloning using unstable hybridoma line	30
2.4.3 Generation of hybridomas by immunoprecipitated bead immunization	30
2.4.3.1 Preparation of immunoprecipitated beads	30
2.4.3.2 Mouse immunization with OKT3 immunoprecipitated beads	31
2.4.3.3 Determination of antibody response in the immunized mouse by indirect immunofluorescence staining	31
2.4.3.4 Hybridoma technique	32
2.4.3.4.1 Preparation of spleen cells	32
2.4.3.4.2 Preparation of myeloma cells	33
2.4.3.4.3 Cell fusion	33
2.4.3.4.4 Screening of hybridomas by indirect immunofluorescence staining	34
2.4.3.4.5 Single cell cloning	34
2.5 Study the utilization of conditioned media for generation of hybridomas by hybridoma technique	34
2.5.1 Preparation of Hb Portland and Hb A ₂	34


2.5.1.1 Preparation of hemolysates	34
2.5.1.2 Preparation of purified Hb Portland	35
2.5.1.3 Preparation of purified Hb A ₂	35
2.5.2 Mouse immunization with Hb Portland and A ₂	36
2.5.3 Determination of antibody response in the immunized mice by indirect ELISA	36
2.5.4 Hybridoma technique	37
2.5.4.1 Preparation of spleen cells	37
2.5.4.2 Preparation of myeloma cells	37
2.5.4.3 Preparation of HAT and HT medium	37
2.5.4.4 Cell fusion	37
2.5.5 Screening of hybridomas by ELISA	38
2.5.6 Single cell cloning	38
2.6 Analysis of protein in the produced conditioned medium by sodium dodesyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE)	38
CHAPTER III: RESULTS	40
3.1 Preparation of conditioned media	40
3.2 Study the utilization of the generated conditioned media for hybridoma single cell cloning	43
3.2.1 Single cell cloning using stable hybridoma line	43
3.2.1.1 Determination of the appropriate conditioned medium	43
3.2.1.2 Determination of the optimal concentration of the produced conditioned medium	48
3.2.2 Single cell cloning using unstable hybridoma line	50

3.3 Utilization of the produced conditioned medium for generation of hybridomas by hybridoma technique	55
3.3.1 Mice immunization and antibody responses	55
3.3.2 Cell fusion	57
3.3.3 Single cell cloning	60
3.4 Analysis of protein contained in conditioned medium by sodium dodesyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE)	68
3.5 Analysis and comparison of cost effectiveness between the produced conditioned medium and commercial reagent	72
CHAPTER IV: DISCUSSION AND CONCLUSION	74
REFERENCES	85
APPENDIX	91
CURRICULUM VITAE	119

LIST OF TABLES

TABLE	PAGE
2.1 The reaction mixture of mycoplasma DNA detection	28
3.1 Various culture conditions and culture supernatants (or conditioned media) used in this study	42
3.2 Growth supporting effect of using BW5147 mouse thymoma Conditioned media on stable hybridoma Thal N/B single cell cloning	45
3.3 Growth supporting effect of using myeloma conditioned media on Stable hybridoma Thal N/B single cell cloning	47
3.4 Growth supporting effect of using various concentrations of BW5147 mouse thymoma conditioned media on stable hybridoma single cell cloning	49
3.5 Growth supporting effect of using various concentrations of BW5147 mouse thymomas conditioned media on un-stable hybridoma single cell cloning	53
3.6 Generation of hybridomas and antibody reactivity screening by using BW5147 conditioned medium and BM condimed H1	59
3.7 Single cell cloning of hybridoma produce anti-Hb Portland using BW5147 conditioned medium and BM condimed H1	63
3.8 Comparison of using BW5147 conditioned medium and BM condimed H1 in generation of hybridomas produce anti-HB E/A ₂ antibody	65
3.9 Comparison of using BW5147 conditioned medium and BM condimed H1 in generation of hybridoma produce anti- activated platelet	66

antibody	
3.10 Comparison of using BW5147 conditioned medium and BM condimed H1 in generation of hybridoma produce anti-CD99 associated molecules	67
3.11 Analysis and comparison of cost effectiveness between BW5147 conditioned medium and commercial reagent BM condimed H1	73



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved


LIST OF FIGURES

FIGURES	PAGE
1.1 The fusion of a myeloma and an antibody-producing cell results in an immortal antibody-producing hybridoma	7
1.2 Metabolic pathways relevant to hybrid selection in medium containing hypoxanthine, aminopterin and thymidine (HAT medium)	11
1.3 Pathway of nucleotide synthesis in the presence of antifolate drug	12
1.4 Drug selections for viable hybridomas	13
1.5 Outlines the procedure of monoclonal antibody production by hybridoma technique	16
1.6 Cloning techniques to obtain cloned hybridomas	21
3.1 Agarose gel electrophoresis of PCR product for determination of mycoplasma infection	41
3.2 Antibody responses of mouse immunized with OKT3 immunoprecipitated beads	52
3.3 Antibody reactivity of culture supernatants of hybridoma clones using the produced BW5147 mouse thymoma condition medium and BM conditioned H1 supplementation	54
3.4 Antibody responses in BALB/c mice after immunizations with Hb Portland and Hb A ₂	56
3.5 Proteins analysis of BW5147 conditioned medium, BM conditioned H1 and 10%FCS-IMDM using 12.5% SDS-PAGE	69
3.6 Proteins analysis of BW5147 conditioned medium, BM conditioned H1	70

and 10%FCS-IMDM using 10% SDS-PAGE

3.7 Proteins analysis of BW5147 conditioned medium, BM condimed H1 71

and 10%FCS-IMDM using 7.5% SDS-PAGE



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved

ABBREVIATIONS

Ab	antibody
APS	ammonium persulphate
BSA	bovine serum albumin
°C	degree Celcius
CD	cluster of differentiation
DEAE-Dextran	diethylaminoethyl-Dextran
DMSO	dimethyl sulfoxide
DNA	deoxyribonucleic acid
EDTA	ethylene diamine tetraacetic acid
ELISA	enzyme-linked immunosorbent assay
EtBr	ethidium bromide
FCS	fetal calf serum
FITC	fluorescein isothiocyanate
g	gram or gravity
HAT	hypoxanthine aminopterin and thymidine
Hb	hemoglobin
HGPRT	hypoxanthine guanine phosphoribosyltransferase
HMI	humoral-mediated immunity
Ig	immunoglobulin
Igs	immunoglobulins
IL-6	interleukin-6
IMDM	iscove's modified Dulbecco's medium

kDa	kilodalton
M	molar
mg	milligram
ml	milliliter
mM	millimolar
mAb	monoclonal antibody
mA	milli ampere
NaN ₃	sodium azide
O.D.	optical density
PBS	phosphate buffered saline
pH	power of hydrogen
PMA	phorbol myristate acetate
rpm	revolution per minute
SDS-PAGE	sodium dodecyl sulfate-polyacrylamide gel electrophoresis
TBE	tris-borate-EDTA
V	volt
μg	microgram
μl	microliter
μm	micrometer