# TABLE OF CONTENTS

9181860 Pa	σo
ACKNOWLEDGMENT	iii
ABSTRACT (in English)	iv
ABSTRACT (in Thai)	vi
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xvi
CHAPTER	
I GENERAL INTRODUCTION	1
1. Flavonoids	2
1.1 Description	2
1.2 Physicochemical properties	4
1.3 Bioactivity	7
1.3.1 Antioxidant activity	7
1.3.2 Anticancer activity	7
1.3.3 Intracellular target	9
1.3.4 Interaction of flavonoids with membrane	10
2. Distribution and localization	11
2.1 Anatomy and physico-chemical properties of lipid	<b>1</b> J
Copyright <sup>©</sup> 2.2 Enthalpy/Entropy of Mai Univers	14
2.2.1 Phase of lipid	14
A 1 2.2.2 Mono lipid system S C 1 V C	16
2.2.3 Lipid mixture system	16
2.3 Transition phase	16
OBJECTIVES	18
THE ORGANIZATION OF THE THESIS DISSERTATION	19

### REFERENCES

<b>II</b> SPECTROPHOTOMETRIC CHARACTERIZATION	27
OF BEHAVIOR AND THE PREDOMINANT SPECIES	
<b>OF FLAVONOIDS IN PHYSIOLOGICAL BUFFER:</b>	
DETERMINATION OF SOLUBILITY, LIPOPHILICITY	
AND ANTICANCER EFFICACY	
ABSTRACT	28
1. INTRODUCTION	29
2. MATERIALS AND METHODS	30
2.1 Chemical and solutions	30
2.2 Spectrophotometric measurements	32
2.3 Cell culture and cytotoxicity assay	32
2.4 Determination of lipophilicity	33
2.5 HPLC analysis	33
3. RESULTS	33
3.1 Spectrophotometric determination of the	33
tautomerization of the flavonoids	
3.2 Determination of log P values	37
3.3 Cytotoxicity of flavonoids	40
4. DISCUSSION	41
5. REFERENCES	43
ົລາອີກຄົ້ນນັດຄົກແດດ້ແມ່ນີ້ແດງໃ	
<b>C O C III INTERACTION OF FLAVONOIDS WITH MYRISTYL</b>	47
MYRISTATE AND TWEEN20 BILAYER MEMBRANE :	
RELATION BETWEEN THE CHEMICAL STRUCTURES	ILY
AND MEAN RATE INFLUX COEFFICIENT ABSTRACT	<b>2</b> 48
1. INTRODUCTION	49
2. MATERIALS AND METHODS	50
2.1 Construction of the multilamellar mysistyl-myristate -Tween20 <sup>®</sup> vesicles	50

20

	2.2 Micrograph of liposomes	50
	2.3 Measurements of doxorubicin entrapped in liposomes	50
	2.4 Measurements of the mean rate influx coefficient	51
	$(k_{+})$ of flavonoids	
	2.5 Measurement the critical temperature of lipid bilayer	51
	2.6 Theoretical approach	51
	3. RESULTS	52
	3.1 Measurements of doxorubicin accumulation and size of	f 52
9	liposome	
0	3.2 Measurement of the mean rate influx coefficient	53
	$(k_{+})$ of flavonoids	
	3.3 Thermal properties of the myristyl myristate and	55
R	Tween20 bialyer	
	3.4 Flavonoids affected the thermal properties of the	55
	myristyl myristate-Tween20 bialyer	
H	4. DISCUSSION	57
5	5. REFERENCES	62
IV	GENERAL DISCUSSION AND CONCLUSION	65

# APPENDIX

IN VITRO AND IN VIVO ASSESSMENT OF CANCER	70
CELL RESPONSE TO FLAVONOIDS BY USING	1411
MOLECULAR IMAGING TECHNIQUES	
Copyrigh ABSTRACT Chiang Mai Univer	SI <sup>71</sup> 72
2. MATERIALS AND METHODS <b>C S C V</b>	<b>e</b> 72
2.1 Cell culture and apoptotic induction assay	72
2.2 Animal experiments	72
2.3 99 <sup>m</sup> Tc-hynic-Annexin V Scintigraphy	73
2.4 Statistical analyses	73

I UNIVER

3. RESULTS	73
3.1 Cytotoxicity of Siamois® polyphenols against	73
myoblast and cancer cells	
3.2 Decrease in the ROSi and alteration of $\Delta \Psi m$	73
of cancer cells	
3.3 Determination of apoptosis against cancer cells	74
3.4 Determination of apoptosis in living xenografted mice	76
4. CONCLUSION	77
5. REFERENCES	77
CURRICURUM VITAE	79
3	
F.	
MAL BERST	
UNIVE UNIVE	

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright<sup>©</sup> by Chiang Mai University All rights reserved

### LIST OF TABLES

# Table Page CHAPTER II 1 Log P values, Ionization rate constants (pKa) and 38 38 Neutral form (%) of flavonoids 2 Cytotoxicity of flavonoids against GLC4, 41 GLC4/adr, K562 and K562/adr 41 1 The effects of flavonoids studied on the gel-to-fluid 60 phase transition of the lipids

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright<sup>©</sup> by Chiang Mai University All rights reserved

# LIST OF FIGURES

	Figur	. กมยนต์ .	Page
CHAP	TER I		8
	1	Albert Szent-GyÖrgyi	2
	2	Nucleus structure of flavonoids	2
	3	Classified of flavonoids	3
	4	An amphiphilic lipid	12
	5	The monomolecular layer of lipid	13
50	6	Scheme of vesicle formation	13
20	7	Phase transition	15
	8	Diagram represent of energy transfer in the system related	15
	1	to temperature	
CHAP'	TER I	u A A A A A A A A A A A A A A A A A A A	
	1	Chemical structure of catechin, eriodictyol, apigenin,	31
	<u> </u>	kaempferol, quercetin, WP279, WP280 and WP283	
	2	Absorption spectra (a) catechin and (b) eriodictyol	34
	3	Absorption spectra apigenin	35
	4	Absorption spectra kaempferol, quercetin	36
	5	Absorption spectra of WP 280, WP 279, and WP283	37
ลิขสา	6	Chemical structure of flavonoid nucleus indicating	39
CIUCII		different functional group substitutions and its log P	
Copyr	igh	difference V Chiang Mai Univ	ersity
	7	Relationship between the experimental and calculated	40
	ľ	log P values I S I E S E I	veo
CHAPTER III			
	1	Micrograph of multilamellar vesicles and histogram of	53
		vesicular size distribution	
	2	Emission spectra of doxorubicin and variation of	54

	fluoreseenee intensity at 500 nm as a function of lineseme	
	nuorescence intensity at 590 mill as a function of hposome	
	contents	
3	Variation of temperature difference of myristyl myristate-	56
	Tween20 bilayer membrane and first derivative spectra	
	of the data	
4	A typical results of apigenin caused the thermal pattern	58
	of myristyl myristate-Tween20 bilayer changes	
5 9	A typical results of WP 283 caused the thermal pattern	58
8	of myristyl myristate-Tween20 bilayer changes.	
Scheme 1	Hydration of keto group of myristyl myristate-	61
	Tween20 bilayer membrane catalysis by acid or base	
Scheme 2	Ketalisation of flavonoids with keto group of	61
275	myristyl myristate-Tween20 bilayer membrane	25
CHAPTER I	IV IV	
Diagram 1	The behavior of flavonoids in aqueous physiological	69
E I	condition	
APPENDIX		
1	Effects of Siamois, Siamois 1 and Siamois 2, doxorubicin	74
	on myocyte cell growth and Siamois 1 and Siamois 2	
	on MDA-MB 435 cells.	
2	Chemical structure of quercetin, kaempferol, apigenin	75
	and eriodictyol	
$\mathbf{S}$	Kinetics of DHCF-DA oxidation	75
adan	Variation of Absolute values of $\Delta \Psi m$ and percentage	76
	of early apoptotic cell as a function of ROSi in K562	
Copyrign	and K562/adr cell.	ersity
<b>A</b> 5	Gamma scintigraphic images showing the enhanced	77 e d
	accumulation of 99mTc-Annexin V in the MDA-MB 435	
	breast tumor xenografted in athymic nude mice and	l ex vivo
	histological tumor tissue.	

	ABBREVIATIONS
K562	Adriamycin-sensitive erythroleukemia cells
K562/adr	Adriamycin-resistant erythroleukemia cells
GLC4	Adriamycin-sensitive small cell lung carcinoma
GLC4/adr	Adriamycin-resistant small cell lung carcinoma
P-gp	P-glycoprotein
MRP1	Multidrug resistance associated protein 1
MTT	Tetrazolium salt,3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl
Q	-2 <i>H</i> -tetrazolium bromide
HEPES	N-2-Hydroxyethylpiperazine-N'-2-ethane-sulfonic acid
IC <sub>50</sub>	The concentration of molecule required for 50% inhibition of the
	proliferation of cell
рКа	Ionization rate constant
Log P	log partition coefficient
D°H	Neutral Form
	Positive charge
Cocyrig	Degree Celsius Chiang Mai University
<b>DPH</b>	1,6-diphenyl-1,3,5-hexatrine <b>PESEIVEO</b>
$T_t$	Critical temperature
$\Delta S$	Entropy
$\Delta H$	Enthalpy



 $k_+$  Mean rate influx coefficient



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่ Copyright<sup>©</sup> by Chiang Mai University All rights reserved