

REFERENCES

- Aasheim B, Ogaard B (1993). Hypodontia in 9-year-old Norwegians related to need of orthodontic treatment. *Scand J Dent Res* 101(5):257-60.
- Albashaireh ZS, Khader YS (2006). The prevalence and pattern of hypodontia of the permanent teeth and crown size and shape deformity affecting upper lateral incisors in a sample of Jordanian dental patients. *Community Dent Health* 23(4):239-43.
- Altug-Atac AT, Erdem D (2007). Prevalence and distribution of dental anomalies in orthodontic patients. *Am J Orthod Dentofacial Orthop* 131(4):510-4.
- Amiel J, Bougeard G, Francannet C, Raclin V, Munnich A, Lyonnet S, et al. (2001). TP63 gene mutation in ADULT syndrome. *Eur J Hum Genet* 9(8):642-5.
- Aulehla A, Wehrle C, Brand-Saberi B, Kemler R, Gossler A, Kanzler B, et al. (2003). Wnt3a plays a major role in the segmentation clock controlling somitogenesis. *Dev Cell* 4(3):395-406.
- Bamshad MJ (2008). p63 and the Ectodermal Dysplasia, Ectrodactyly, and Cleft Lip and/or Palate (EEC), Limb-Mammary (LMS), Ankyloblepharon, Ectrodactyly, and Cleft Lip/Palate (AEC, Hay-Wells), and Acro-Dermato-Ungual-Lacrima-Digit (ADULT) Syndromes and Ectrodactyly (Split Hand/Foot Malformation). In: *Inborn Errors of Development; The Molecular Basis of Clinical Disorders of Morphogenesis*, 2nd ed. New York: Oxford university press, pp.1017-26.
- Barbieri CE, Pietsenpol JA (2006). p63 and epithelial biology. *Exp Cell Res* 312(6):695-706.
- Barrow LL, van Bokhoven H, Daack-Hirsch S, Andersen T, van Beersum SE, Gorlin R, et al. (2002). Analysis of the p63 gene in classical EEC syndrome, related syndromes, and non-syndromic orofacial clefts. *J Med Genet* 39(8):559-66.
- Barski A, Frenkel B (2004). CHIP Display: novel method for identification of genomic targets of transcription factors. *Nucleic Acids Res* 32(12):e104.
- Baskin LS, Hayward SW, Young P, Cunha GR (1996). Role of mesenchymal-epithelial interactions in normal bladder development. *J Urol* 156(5):1820-7.
- Benard J, Douc-Rasy S, Ahomadegbe JC (2003). TP53 family members and human cancers. *Hum Mutat* 21(3):182-91.

- Boehm CD (1989). Use of polymerase chain reaction for diagnosis of inherited disorders. *Clin Chem* 35(9):1843-8.
- Brunner HG, Hamel BC, Van Bokhoven H (2002). The p63 gene in EEC and other syndromes. *J Med Genet* 39(6):377-81.
- Buss PW (1994). Cleft hand/foot: clinical and developmental aspects. *J Med Genet* 31(9):726-30.
- Buss PW, Hughes HE, Clarke A (1995). Twenty-four cases of the EEC syndrome: clinical presentation and management. *J Med Genet* 32(9):716-23.
- Candi E, Terrinoni A, Rufini A, Chikh A, Lena AM, Suzuki Y, et al. (2006). p63 is upstream of IKK alpha in epidermal development. *J Cell Sci* 119(Pt 22):4617-22.
- Carlson B (2004). Head and neck. In: Human embryology and development biology, 3rd ed. Philadelphia: Mosby, pp. 317-51.
- Celli J, Duijf P, Hamel BC, Bamshad M, Kramer B, Smits AP, et al. (1999). Heterozygous germline mutations in the p53 homolog p63 are the cause of EEC syndrome. *Cell* 99(2):143-53.
- Chan I, Harper JJ, Mellerio JE, McGrath JA (2004). ADULT ectodermal dysplasia syndrome resulting from the missense mutation R298Q in the p63 gene. *Clin Exp Dermatol* 29(6):669-72.
- Chan I, McGrath JA, Kivirikko S (2005). Rapp-Hodgkin syndrome and the tail of p63. *Clin Exp Dermatol* 30(2):183-6.
- Cheng W, Jacobs WB, Zhang JJ, Moro A, Park JH, Kushida M, et al. (2006). DeltaNp63 plays an anti-apoptotic role in ventral bladder development. *Development* 133(23):4783-92.
- Chung CJ, Han JH, Kim KH (2008). The pattern and prevalence of hypodontia in Koreans. *Oral Dis*. 14:620-25.
- Clements SE, Techanukul T, Coman D, Mellerio JE, McGrath JA (2009). Molecular basis of EEC (ectrodactyly, ectodermal dysplasia, clefting) syndrome: five new mutations in the DNA-binding domain of the TP63 gene and genotype-phenotype correlation. *Br J Dermatol* 162:201-7.
- Comperat E, Camparo P, Haus R, Chartier-Kastler E, Bart S, Delcourt A, et al. (2006). Immunohistochemical expression of p63, p53 and MIB-1 in urinary bladder carcinoma. A tissue microarray study of 158 cases. *Virchows Arch* 448(3):319-24.

- Daniels TE, Silverman S, Jr., Michalski JP, Greenspan JS, Sylvester RA, Talal N (1975). The oral component of Sjogren's syndrome. *Oral Surg Oral Med Oral Pathol* 39(6):875-85.
- Das P, Stockton DW, Bauer C, Shaffer LG, D'Souza RN, Wright T, et al. (2002). Haploinsufficiency of PAX9 is associated with autosomal dominant hypodontia. *Hum Genet* 110(4):371-6.
- Davis PJ (1987). Hypodontia and hyperdontia of permanent teeth in Hong Kong schoolchildren. *Community Dent Oral Epidemiol* 15(4):218-20.
- Di Como CJ, Urist MJ, Babayan I, Drobnjak M, Hedvat CV, Teruya-Feldstein J, et al. (2002). p63 expression profiles in human normal and tumor tissues. *Clin Cancer Res* 8(2):494-501.
- Dianzani I, Garelli E, Gustavsson P, Carando A, Gustafsson B, Dahl N, et al. (2003). Rapp-Hodgkin and AEC syndromes due to a new frameshift mutation in the TP63 gene. *J Med Genet* 40(12):e133.
- Dong J, Amor D, Aldred MJ, Gu T, Escamilla M, MacDougall M (2005). DLX3 mutation associated with autosomal dominant amelogenesis imperfecta with taurodontism. *Am J Med Genet A* 133(2):138-41.
- Duijf PH, van Bokhoven H, Brunner HG (2003). Pathogenesis of split-hand/split-foot malformation. *Hum Mol Genet* 12 Review Issue I:R51-60.
- Endo T, Ozoe R, Kubota M, Akiyama M, Shimooka S (2006). A survey of hypodontia in Japanese orthodontic patients. *Am J Orthod Dentofacial Orthop* 129(1):29-35.
- Fekonja A (2005). Hypodontia in orthodontically treated children. *Eur J Orthod* 27(5):457-60.
- Ferguson MW (1988). Palate development. *Development* 103 Suppl:41-60.
- Fomenkov A, Huang YP, Topaloglu O, Brechman A, Osada M, Fomenkova T, et al. (2003). P63 alpha mutations lead to aberrant splicing of keratinocyte growth factor receptor in the Hay-Wells syndrome. *J Biol Chem* 278(26):23906-14.
- Ghioni P, Bolognese F, Duijf PH, Van Bokhoven H, Mantovani R, Guerrini L (2002). Complex transcriptional effects of p63 isoforms: identification of novel activation and repression domains. *Mol Cell Biol* 22(24):8659-68.
- Gorlin RI, Cohen MMj, Hannekam RCM (2001). Teratogenic Agents. In: Syndromes of the head and neck. New York: Oxford University Press, pp. 14-34.

- Haldeman RJ, Cooper LF, Hart TC, Phillips C, Boyd C, Lester GE, et al. (2004). Increased bone density associated with DLX3 mutation in the tricho-dento-osseous syndrome. *Bone* 35(4):988-97.
- Hamada T, Chan I, Willoughby CE, Goudie DR, McGrath JA (2002). Common mutations in Arg304 of the p63 gene in ectrodactyly, ectodermal dysplasia, clefting syndrome: lack of genotype-phenotype correlation and implications for mutation detection strategies. *J Invest Dermatol* 119(5):1202-3.
- Han D, Gong Y, Wu H, Zhang X, Yan M, Wang X, et al. (2008). Novel EDA mutation resulting in X-linked non-syndromic hypodontia and the pattern of EDA-associated isolated tooth agenesis. *Eur J Med Genet* 51(6):536-46.
- Hassan MQ, Javed A, Morasso MI, Karlin J, Montecino M, van Wijnen AJ, et al. (2004). Dlx3 transcriptional regulation of osteoblast differentiation: temporal recruitment of Msx2, Dlx3, and Dlx5 homeodomain proteins to chromatin of the osteocalcin gene. *Mol Cell Biol* 24(20):9248-61.
- Holder-Espinasse M, Martin-Coignard D, Escande F, Manouvrier-Hanu S (2007). A new mutation in TP63 is associated with age-related pathology. *Eur J Hum Genet* 15(11):1115-20.
- Hu JC, Simmer JP (2007). Developmental biology and genetics of dental malformations. *Orthod Craniofac Res* 10(2):45-52.
- Hu MC, Wang Y, Qiu WR, Mikhail A, Meyer CF, Tan TH (1999). Hematopoietic progenitor kinase-1 (HPK1) stress response signaling pathway activates IkkappaB kinases (IKK-alpha/beta) and IKK-beta is a developmentally regulated protein kinase. *Oncogene* 18(40):5514-24.
- Ianakiev P, Kilpatrick MW, Toudjarska I, Basel D, Beighton P, Tsipouras P (2000). Split-hand/split-foot malformation is caused by mutations in the p63 gene on 3q27. *Am J Hum Genet* 67(1):59-66.
- Kantaputra PN, Pruksachatkunakorn C, Vanittanakom P (1998). Rapp-Hodgkin syndrome with palmoplantar keratoderma, glossy tongue, congenital absence of lingual frenum and of sublingual caruncles: newly recognized findings. *Am J Med Genet* 79(5):343-6.
- Kantaputra PN, Hamada T, Kumchai T, McGrath JA (2003). Heterozygous mutation in the SAM domain of p63 underlies Rapp-Hodgkin ectodermal dysplasia. *J Dent Res* 82(6):433-7.
- Kettunen P, Karavanova I, Thesleff I (1998). Responsiveness of developing dental tissues to fibroblast growth factors: expression of splicing alternatives of FGFR1, -2, -3, and of FGFR4; and stimulation of cell proliferation by FGF-2, -4, -8, and -9. *Dev Genet* 22(4):374-85.

- King KE, Ponnampereuma RM, Yamashita T, Tokino T, Lee LA, Young MF, et al. (2003). deltaNp63alpha functions as both a positive and a negative transcriptional regulator and blocks in vitro differentiation of murine keratinocytes. *Oncogene* 22(23):3635-44.
- Koillinen H, Lahermo P, Rautio J, Hukki J, Peyrard-Janvid M, Kere J (2005). A genome-wide scan of non-syndromic cleft palate only (CPO) in Finnish multiplex families. *J Med Genet* 42(2):177-84.
- Koster MI, Roop DR (2004a). The role of p63 in development and differentiation of the epidermis. *J Dermatol Sci* 34(1):3-9.
- Koster MI, Roop DR (2004b). p63 and epithelial appendage development. *Differentiation* 72(8):364-70.
- Koster MI, Dai D, Marinari B, Sano Y, Costanzo A, Karin M, et al. (2007). p63 induces key target genes required for epidermal morphogenesis. *Proc Natl Acad Sci U S A* 104(9):3255-60.
- Lammi L, Arte S, Somer M, Jarvinen H, Lahermo P, Thesleff I, et al. (2004). Mutations in AXIN2 cause familial tooth agenesis and predispose to colorectal cancer. *Am J Hum Genet* 74(5):1043-50.
- Laurikkala J, Mikkola ML, James M, Tummers M, Mills AA, Thesleff I (2006). p63 regulates multiple signalling pathways required for ectodermal organogenesis and differentiation. *Development* 133(8):1553-63.
- Lee SK, Lee ZH, Lee SJ, Ahn BD, Kim YJ, Lee SH, et al. (2008). DLX3 mutation in a new family and its phenotypic variations. *J Dent Res* 87(4):354-7.
- Leoyklang P, Siriwan P, Shotelersuk V (2006). A mutation of the p63 gene in non-syndromic cleft lip. *J Med Genet* 43(6):e28.
- Li H, Marijanovic I, Kronenberg MS, Erceg I, Stover ML, Velonis D, et al. (2008). Expression and function of Dlx genes in the osteoblast lineage. *Dev Biol*.
- Lo Iacono N, Mantero S, Chiarelli A, Garcia E, Mills AA, Morasso MI, et al. (2008). Regulation of Dlx5 and Dlx6 gene expression by p63 is involved in EEC and SHFM congenital limb defects. *Development* 135(7):1377-88.
- Maas SM, de Jong TP, Buss P, Hennekam RC (1996). EEC syndrome and genitourinary anomalies: an update. *Am J Med Genet* 63(3):472-8.
- Maclean K, Holme SA, Gilmour E, Taylor M, Scheffer H, Graf N, et al. (2007). EEC syndrome, Arg227Gln TP63 mutation and micturition difficulties: Is there a genotype-phenotype correlation? *Am J Med Genet A* 143A(10):1114-9.

- Mangiulli M, Valletti A, Caratozzolo MF, Tullo A, Sbisa E, Pesole G, et al. (2009). Identification and functional characterization of two new transcriptional variants of the human p63 gene. *Nucleic Acids Res* 37(18):6092-104.
- Mattheeuws N, Dermaut L, Martens G (2004). Has hypodontia increased in Caucasians during the 20th century? A meta-analysis. *Eur J Orthod* 26(1):99-103.
- McDaniell R, Warthen DM, Sanchez-Lara PA, Pai A, Krantz ID, Piccoli DA, et al. (2006). NOTCH2 mutations cause Alagille syndrome, a heterogeneous disorder of the notch signaling pathway. *Am J Hum Genet* 79(1):169-73.
- McGrath JA, Duijf PH, Doetsch V, Irvine AD, de Waal R, Vanmolkot KR, et al. (2001). Hay-Wells syndrome is caused by heterozygous missense mutations in the SAM domain of p63. *Hum Mol Genet* 10(3):221-9.
- McKeon F (2004). p63 and the epithelial stem cell: more than status quo? *Genes Dev* 18(5):465-9.
- Medeiros RB, Papenfuss KJ, Hoium B, Coley K, Jadrach J, Goh SK, et al. (2009). Novel sequential ChIP and simplified basic ChIP protocols for promoter co-occupancy and target gene identification in human embryonic stem cells. *BMC Biotechnol* 9:59.
- Melino G, Lu X, Gasco M, Crook T, Knight RA (2003). Functional regulation of p73 and p63: development and cancer. *Trends Biochem Sci* 28(12):663-70.
- Mikkola ML, Thesleff I (2003). Ectodysplasin signaling in development. *Cytokine Growth Factor Rev* 14(3-4):211-24.
- Mikkola ML (2007). p63 in skin appendage development. *Cell Cycle* 6(3):285-90.
- Mills AA, Zheng B, Wang XJ, Vogel H, Roop DR, Bradley A (1999). p63 is a p53 homologue required for limb and epidermal morphogenesis. *Nature* 398(6729):708-13.
- Moll UM, Slade N (2004). p63 and p73: roles in development and tumor formation. *Mol Cancer Res* 2(7):371-86.
- Morasso MI, Mahon KA, Sargent TD (1995). A *Xenopus* distal-less gene in transgenic mice: conserved regulation in distal limb epidermis and other sites of epithelial-mesenchymal interaction. *Proc Natl Acad Sci U S A* 92(9):3968-72.
- Morasso MI, Grinberg A, Robinson G, Sargent TD, Mahon KA (1999). Placental failure in mice lacking the homeobox gene *Dlx3*. *Proc Natl Acad Sci U S A* 96(1):162-7.

- Morasso MI, Radoja N (2005). Dlx genes, p63, and ectodermal dysplasias. *Birth Defects Res C Embryo Today* 75(3):163-71.
- Murray-Zmijewski F, Lane DP, Bourdon JC (2006). p53/p63/p73 isoforms: an orchestra of isoforms to harmonise cell differentiation and response to stress. *Cell Death Differ* 13(6):962-72.
- Neilson DE, Brunger JW, Heeger S, Bamshad M, Robin NH (2002). Mixed clefting type in Rapp-Hodgkin syndrome. *Am J Med Genet* 108(4):281-4.
- O'Quinn JR, Hennekam RC, Jorde LB, Bamshad M (1998). Syndromic ectrodactyly with severe limb, ectodermal, urogenital, and palatal defects maps to chromosome 19. *Am J Hum Genet* 62(1):130-5.
- Ohazama A, Hu Y, Schmidt-Ullrich R, Cao Y, Scheidereit C, Karin M, et al. (2004). A dual role for Ikk alpha in tooth development. *Dev Cell* 6(2):219-27.
- Osada M, Park HL, Nagakawa Y, Begum S, Yamashita K, Wu G, et al. (2006). A novel response element confers p63- and p73-specific activation of the WNT4 promoter. *Biochem Biophys Res Commun* 339(4):1120-8.
- Park BJ, Lee SJ, Kim JI, Lee CH, Chang SG, Park JH, et al. (2000). Frequent alteration of p63 expression in human primary bladder carcinomas. *Cancer Res* 60(13):3370-4.
- Parsa R, Yang A, McKeon F, Green H (1999). Association of p63 with proliferative potential in normal and neoplastic human keratinocytes. *J Invest Dermatol* 113(6):1099-105.
- Payne AS, Yan AC, Ilyas E, Li W, Seykora JT, Young TL, et al. (2005). Two novel TP63 mutations associated with the ankyloblepharon, ectodermal defects, and cleft lip and palate syndrome: a skin fragility phenotype. *Arch Dermatol* 141(12):1567-73.
- Perez CA, Pietenpol JA (2007). Transcriptional programs regulated by p63 in normal epithelium and tumors. *Cell Cycle* 6(3):246-54.
- Pispa J, Thesleff I (2003). Mechanisms of ectodermal organogenesis. *Dev Biol* 262(2):195-205.
- Pozo G, Canun S, Kofman-Alfaro S, Zenteno JC (2004). A de novo heterozygous point mutation in the p63 gene causing the syndrome of ectrodactyly, ectodermal dysplasia and facial clefting. *Br J Dermatol* 151(4):930-2.
- Price JA, Bowden DW, Wright JT, Pettenati MJ, Hart TC (1998a). Identification of a mutation in DLX3 associated with tricho-dento-osseous (TDO) syndrome. *Hum Mol Genet* 7(3):563-9.

- Price JA, Wright JT, Kula K, Bowden DW, Hart TC (1998b). A common DLX3 gene mutation is responsible for tricho-dento-osseous syndrome in Virginia and North Carolina families. *J Med Genet* 35(10):825-8.
- Price JA, Wright JT, Walker SJ, Crawford PJ, Aldred MJ, Hart TC (1999). Tricho-dento-osseous syndrome and amelogenesis imperfecta with taurodontism are genetically distinct conditions. *Clin Genet* 56(1):35-40.
- Propping P, Friedl W, Wienker TF, Uhlhaas S, Zerres K (2000). ADULT syndrome allelic to limb mammary syndrome (LMS)? *Am J Med Genet* 90(2):179-82.
- Pummila M, Fliniaux I, Jaatinen R, James MJ, Laurikkala J, Schneider P, et al. (2007). Ectodysplasin has a dual role in ectodermal organogenesis: inhibition of Bmp activity and induction of Shh expression. *Development* 134(1):117-25.
- Radoja N, Guerrini L, Lo Iacono N, Merlo GR, Costanzo A, Weinberg WC, et al. (2007). Homeobox gene Dlx3 is regulated by p63 during ectoderm development: relevance in the pathogenesis of ectodermal dysplasias. *Development* 134(1):13-8.
- Rapp RS, Hodgkin WE (1968). Anhidrotic ectodermal dysplasia: autosomal dominant inheritance with palate and lip anomalies. *J Med Genet* 5(4):269-72.
- Reisler TT, Patton MA, Meagher PP (2006). Further phenotypic and genetic variation in ADULT syndrome. *Am J Med Genet A* 140(22):2495-500.
- Rice DP (2005). Craniofacial anomalies: from development to molecular pathogenesis. *Curr Mol Med* 5(7):699-722.
- Rinne T, Hamel B, van Bokhoven H, Brunner HG (2006a). Pattern of p63 mutations and their phenotypes--update. *Am J Med Genet A* 140(13):1396-406.
- Rinne T, Spadoni E, Kjaer KW, Danesino C, Larizza D, Kock M, et al. (2006b). Delineation of the ADULT syndrome phenotype due to arginine 298 mutations of the p63 gene. *Eur J Hum Genet* 14(8):904-10.
- Rinne T, Brunner HG, van Bokhoven H (2007). p63-associated disorders. *Cell Cycle* 6(3):262-8.
- Rinne T, Clements SE, Lamme E, Duijf PH, Bolat E, Meijer R, et al. (2008). A novel translation re-initiation mechanism for the p63 gene revealed by amino-terminal truncating mutations in Rapp-Hodgkin/Hay-Wells-like syndromes. *Hum Mol Genet* 17(13):1968-77.
- Roelfsema NM, Cobben JM (1996). The EEC syndrome: a literature study. *Clin Dysmorphol* 5(2):115-27.

- Ropke A, Kujat A, Graber M, Giannakudis J, Hansmann I (2003). Identification of 36 novel Jagged1 (JAG1) mutations in patients with Alagille syndrome. *Hum Mutat* 21(1):100.
- Rufini A, Weil M, McKeon F, Barlattani A, Melino G, Candi E (2006). p63 protein is essential for the embryonic development of vibrissae and teeth. *Biochem Biophys Res Commun* 340(3):737-41.
- Sahin MT, Turel-Ermertcan A, Chan I, McGrath JA, Ozturkcan S (2004). Ectodermal dysplasia showing clinical overlap between AEC, Rapp-Hodgkin and CHAND syndromes. *Clin Exp Dermatol* 29(5):486-8.
- Sasaki Y, Ishida S, Morimoto I, Yamashita T, Kojima T, Kihara C, et al. (2002). The p53 family member genes are involved in the Notch signal pathway. *J Biol Chem* 277(1):719-24.
- Satokata I, Maas R (1994). Msx1 deficient mice exhibit cleft palate and abnormalities of craniofacial and tooth development. *Nat Genet* 6(4):348-56.
- Schultz J, Ponting CP, Hofmann K, Bork P (1997). SAM as a protein interaction domain involved in developmental regulation. *Protein Sci* 6(1):249-53.
- Scoumanne A, Harms KL, Chen X (2005). Structural basis for gene activation by p53 family members. *Cancer Biol Ther* 4(11):1178-85.
- Serber Z, Lai HC, Yang A, Ou HD, Sigal MS, Kelly AE, et al. (2002). A C-terminal inhibitory domain controls the activity of p63 by an intramolecular mechanism. *Mol Cell Biol* 22(24):8601-11.
- Shimada T, Kawai T, Takeda K, Matsumoto M, Inoue J, Tatsumi Y, et al. (1999). IKK-i, a novel lipopolysaccharide-inducible kinase that is related to IkappaB kinases. *Int Immunol* 11(8):1357-62.
- Sifakis S, Basel D, Ianakiev P, Kilpatrick M, Tsipouras P (2001). Distal limb malformations: underlying mechanisms and clinical associations. *Clin Genet* 60(3):165-72.
- Sisman Y, Uysal T, Gelgor IE (2007). Hypodontia. Does the prevalence and distribution pattern differ in orthodontic patients? *Eur J Dent* 1(3):167-73.
- Spak CJ, Johnson G, Ekstrand J (1994). Caries incidence, salivary flow rate and efficacy of fluoride gel treatment in irradiated patients. *Caries Res* 28(5):388-93.
- Strachan T, Read AP (2004). DNA structure and gene expression. In: Human molecular genetics, 3rd ed. London and New York: Garland Science, pp. 3-32.

- Temtamy SA, McKusick VA (1978). Absence deformities as isolated malformations. In: The genetics of hand malformations. *Birth Defects Orig Artic Ser* 14(3):iv, 36-71.
- Testoni B, Borrelli S, Tenedini E, Alotto D, Castagnoli C, Piccolo S, et al. (2006). Identification of new p63 targets in human keratinocytes. *Cell Cycle* 5(23):2805-11.
- Thanos CD, Bowie JU (1999). p53 Family members p63 and p73 are SAM domain-containing proteins. *Protein Sci* 8(8):1708-10.
- Thesleff I (2003). Epithelial-mesenchymal signalling regulating tooth morphogenesis. *J Cell Sci* 116(Pt 9):1647-8.
- Thilander B, Myrberg N (1973). The prevalence of malocclusion in Swedish schoolchildren. *Scand J Dent Res* 81(1):12-21.
- Thomason HA, Dixon MJ, Dixon J (2008). Facial clefting in Tp63 deficient mice results from altered Bmp4, Fgf8 and Shh signaling. *Dev Biol* 321(1):273-82.
- Tucker A, Sharpe P (2004). The cutting-edge of mammalian development; how the embryo makes teeth. *Nat Rev Genet* 5(7):499-508.
- Tucker AS, Al Khamis A, Sharpe PT (1998). Interactions between Bmp-4 and Msx-1 act to restrict gene expression to odontogenic mesenchyme. *Dev Dyn* 212(4):533-9.
- Urist MJ, Di Como CJ, Lu ML, Charytonowicz E, Verbel D, Crum CP, et al. (2002). Loss of p63 expression is associated with tumor progression in bladder cancer. *Am J Pathol* 161(4):1199-206.
- van Bokhoven H, Jung M, Smits AP, van Beersum S, Ruschendorf F, van Steensel M, et al. (1999). Limb mammary syndrome: a new genetic disorder with mammary hypoplasia, ectrodactyly, and other Hand/Foot anomalies maps to human chromosome 3q27. *Am J Hum Genet* 64(2):538-46.
- van Bokhoven H, Hamel BC, Bamshad M, Sangiorgi E, Gurrieri F, Duijf PH, et al. (2001). p63 Gene mutations in eec syndrome, limb-mammary syndrome, and isolated split hand-split foot malformation suggest a genotype-phenotype correlation. *Am J Hum Genet* 69(3):481-92.
- van Bokhoven H, Brunner HG (2002). Splitting p63. *Am J Hum Genet* 71(1):1-13.
- van Bokhoven H, McKeon F (2002). Mutations in the p53 homolog p63: allele-specific developmental syndromes in humans. *Trends Mol Med* 8(3):133-9.

- van der Reijden WA, van der Kwaak JS, Veerman EC, Nieuw Amerongen AV (1996). Analysis of the concentration and output of whole salivary constituents in patients with Sjogren's syndrome. *Eur J Oral Sci* 104(4 (Pt 1)):335-40.
- Vieira AR (2003). Oral clefts and syndromic forms of tooth agenesis as models for genetics of isolated tooth agenesis. *J Dent Res* 82(3):162-5.
- Vigano MA, Mantovani R (2007). Hitting the numbers: the emerging network of p63 targets. *Cell Cycle* 6(3):233-9.
- Wessagowit V, Mellerio JE, Pembroke AC, McGrath JA (2000). Heterozygous germline missense mutation in the p63 gene underlying EEC syndrome. *Clin Exp Dermatol* 25(5):441-3.
- Westfall MD, Mays DJ, Sniezek JC, Pietenpol JA (2003). The Delta Np63 alpha phosphoprotein binds the p21 and 14-3-3 sigma promoters in vivo and has transcriptional repressor activity that is reduced by Hay-Wells syndrome-derived mutations. *Mol Cell Biol* 23(7):2264-76.
- Westfall MD, Pietenpol JA (2004). p63: Molecular complexity in development and cancer. *Carcinogenesis* 25(6):857-64.
- Wright JT, Kula K, Hall K, Simmons JH, Hart TC (1997). Analysis of the trichodento-osseous syndrome genotype and phenotype. *Am J Med Genet* 72(2):197-204.
- Yang A, Kaghad M, Wang Y, Gillett E, Fleming MD, Dotsch V, et al. (1998). p63, a p53 homolog at 3q27-29, encodes multiple products with transactivating, death-inducing, and dominant-negative activities. *Mol Cell* 2(3):305-16.
- Yang A, Schweitzer R, Sun D, Kaghad M, Walker N, Bronson RT, et al. (1999). p63 is essential for regenerative proliferation in limb, craniofacial and epithelial development. *Nature* 398(6729):714-8.
- Yang A, Zhu Z, Kapranov P, McKeon F, Church GM, Gingeras TR, et al. (2006). Relationships between p63 binding, DNA sequence, transcription activity, and biological function in human cells. *Mol Cell* 24(4):593-602.
- Zhang H, Hu G, Wang H, Sciavolino P, Iler N, Shen MM, et al. (1997). Heterodimerization of Msx and Dlx homeoproteins results in functional antagonism. *Mol Cell Biol* 17(5):2920-32.