

BIBLIOGRAPHY

1. English JD. Early treatment of skeletal open bite malocclusions. *Am J Orthod Dentofacial Orthop* 2002; 121(6): 563-5.
2. Ohmae M, Saito S, Morohashi T, Seki K, Qu H, Kanomi R, Yamasaki K, Okano T, Yamada S, Shibasaki Y. A clinical and histological evaluation of titanium mini-implants as anchors for orthodontic intrusion in the beagle dog. *Am J Orthod Dentofacial Orthop* 2001; 119(5): 489-97.
3. Park HS, Kwon OW, Sung JH. Nonextraction treatment of an open bite with microscrew implant anchorage. *Am J Orthod Dentofacial Orthop* 2006; 130(3): 391-402.
4. Poggio PM, Incorvati C, Velo S, Carano A. "Safe Zone": A guide for miniscrew positioning in the maxillary and mandibular arch. *Angle Orthod* 2006;76(2):191-97.
5. Beane RA, Jr. Nonsurgical management of the anterior open bite: a review of the options. *Semin Orthod* 1999; 5(4): 275-83.
6. Sherwood KH, Burch JG, Thompson WJ. Closing anterior open bites by intruding molars with titanium miniplate anchorage. *Am J Orthod Dentofacial Orthop* 2002; 122(6): 593-600.
7. Yoa CCJ, Lee JJ, Chang ZCJ, Chang HF, Chen YJ. Maxillar Molar Intrusion with Fixed Appliances and Mini-implant anchorage Studied in Three Dimensions. *Angle Orthod* 2005; 75(5): 754-60.

8. Xun C, Zeng X, Wang X. Microscrew Anchorage in Skeletal Anterior Open-bite Treatment. *Angle Orthod* 2007; 77(1): 47-56.
9. Park HS, Jang BK, Kyung HM. Maxillary molar intrusion with micro-implant anchorage (MIA). *Aust Orthod J* 2005; 21(2): 129-35.
10. Park HS, Kwon TG, Kwon OW. Treatment of open bite with microscrew implant anchorage. *Am J Orthod Dentofacial Orthop* 2004; 126(5): 627-36.
11. Lee JS, Kim DH, Park YC, Kyung SH, Kim TK. The efficient Use of Midpalatal Miniscrew Implants. *Angle Orthod* 2004; 74(5): 711-14.
12. Dudic A, Kiliaridis S, Mombelli A, Giannopoulou C. Composition changes in gingival crevicular fluid during orthodontic tooth movement: comparisons between tension and compression sides. *Eur J Oral Sci* 2006; 114: 416-22.
13. Jaito N, Jotikasthira D, Krisanaprakornkit S, Ong-chai S, Kongtawelert P. Monitoring of chondroitin-6-sulfate levels in gingival crevicular fluid during orthodontic canine movement. In: Davidovitch Z MJ, Suthanarak S, editor. *Biological Mechanisms of Tooth Eruption, Resorption and Movement*. Boston: Harvard Society for the Advancement of Orthodontics, Boston, Massachusetts; 2006. p. 289-95.
14. Sari E, Ucar C. Interleukine 1B levels around Microscrew implants during Orthodontic Tooth Movement. *Angle Orthod* 2007; 77(6): 1073-78.
15. Intachai I, Kritsanaprakornkit S, Kongtawelert P, Ong-chai S, Buranastidporn B, Suzuki EY, Jotikasthira. Chondroitin sulphate epitope (WF6 epitope) levels in peri-miniscrew implant crevicular fluid during orthodontic loading. *Eur J Orthod* 2010; 32: 60-65.

16. Subtelny JD, Sakuda M. Open-bite: Diagnosis and treatment. *Am J Orthod* 1964; May: 337-58.
17. Ferguson JW. The assessment and treatment of anterior open bite. *Dent Update* 1995; 22(4): 163-8.
18. Sung JH, Kyung HM, Bae SM, Park HS, Kwon OW, McNamara, Jr JA. *Microimplants in orthodontics*. 1 ed. Daegu: Dentos, Inc; 2006.
19. Papadopoulos MA. The use of miniscrew implants for temporary skeletal anchorage in orthodontics: A comprehensive review. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2007; 103: e6-e15.
20. Anka G. Use of miniscrews as temporary anchorage devices in orthodontic practice. I-Introduction. *Aust Orthod J* 2006; 22: 131-39.
21. Umemori M, Sugawara J, Mitani H, Nagasaka H, Kawamura H. Skeletal anchorage system for open-bite correction. *Am J Orthod Dentofacial Orthop* 1999; 115(2): 166-74.
22. TanaKa E, Iwabe T, Kawai N, Nishi M, Dalla-Bona D, Hasegawa T, Tanne K. An adult case of skeletal open bite with a large lower anterior facial height. *Angle Orthod* 2005; 75(3): 465-71.
23. Gurton AU, Akin E, Karacay S. Initial intrusion of the molars in the treatment of anterior open bite malocclusions in growing patients. *Angle Orthod* 2004; 74(4): 454-64.
24. Liou E, Pai B, Lin J. Do miniscrews remain stationary under orthodontic force? *Am J Orthod Dentofacial Orthop* 2004; 126: 42-47.

25. Giannobile WV, Riviere GR, Gorski JP, Tira DE, Cobb CM. Glycosaminoglycans and periodontal disease: Analysis of GCF by Safranin O. *J Periodontol* 1993; 64: 186-90.
26. Crutis MA, Griffiths GS, Price SJ, Coulthurst SK, Johnson NW. The total protein concentration of gingival crevicular fluid. *J Clin Periodontol* 1988; 15: 628-32.
27. Waddington RJ, Embery G. Proteoglycans and orthodontic tooth movement. *J Orthod* 2001; 28(4): 281-90.
28. Kavadia-Tsatala S, Kaklamanos EG, Tsalikis L. Effects of orthodontic treatment on gingival crevicular fluid flow rate and composition: Clinical implications and applications. *Int J Adult Orthod Orthognath Surg* 2002; 17(3): 191-205.
29. Kagayama M, Samano y, Mizoguchi I, Kamo N, Takahashi I, Mitani H. Localization of glycosaminoglycans in periodontal ligament during physiological and experimental tooth movement. *J Periodont Res* 1996; 31: 229-34.
30. Last KS, Donkin C, Embery G. Glycosaminoglycans in human gingival crevicular fluid during orthodontic movement. *Arch Oral Biol* 1988; 33: 907-12.
31. Bischof M, Nedir R, Szmukler-Moncler S, Bernard J, Samson J. Implant stability measurement of delayed and immediately loaded implants during healing. A clinical resonance-frequency analysis study with sandblasted-and-etched ITI implants. *Clin Oral Impl res* 2004; 24: 529-39.
32. Kravitz ND, Kusnoto B. Risks and complications of orthodontic miniscrews. *Am J Orthod Dentofacial Orthop* 2007; 131(4): s43-s51.

33. Miyawaki S, Koyama I, Inoue M, Mishima K, Sugahara T, Takano-Yamamoto T. Factors associated with the stability of titanium screws placed in the posterior region for orthodontic anchorage. *Am J Orthod Dentofacial Orthop* 2003; 124(4): 373-78.
34. Cheng SJ, Tseng IY, Lee JJ, Kok SH. A prospective study of the risk factors associated with failure of mini-implants used for orthodontic anchorage. *Int J Oral Maxillofac Implants* 2004; 19: 100-06.
35. Sewerin IP, Gotfredsen K, Stoltze K. Accuracy of radiographic diagnosis of peri-implant radiolucencies - an *in vitro* experiment. *Clin Oral Impl res* 1997; 8: 299-304.
36. Jepson S, Ruhling A, Jepson K, Ohlenbusch B, Alber H-K. Progressiv peri-implantitis. Incidence and prediction of peri-implant attachment loss. *Clin Oral Impl res* 1996; 7: 133-42.
37. Alsaadi G, Quirynen M, Michiels K, Jacobs R, van Steenberghe D. A biomechanical assessment of the relation between the oral implant stability at insertion and subjective bone quality assessment. *J Clin Periodontol* 2007; 34: 359-66.
38. Kim JW, Ahn SJ, Chang YI. Histomorphometric and mechanical analyses of the drill-free screw as orthodontic anchorage. *Am J Orthod Dentofacial Orthop* 2005; 128: 190-94.
39. Geng JP, Tan KB, Liu GR. Application of finite element analysis in implant dentistry: a review of literature. *J Prosthet Dent* 2001; 85: 585-98.
40. Costa A, Raffaini M, Melsen B. Microscrews as orthodontic anchorage: A preliminary report. *Int J Adult Orthod Orthog Surg* 1998; 13: 201-09.

41. Motoyoshi M, Yano S, Tsuruika T, Shimizu N. Biomechanical effect of abutment on stability of orthodontic mini-implant. A finite element analysis. *Clin Oral Impl res* 2005; 16: 480-85.
42. Meridith N. Assessment of implant stability as a prognosis determinant. *Int J prosthodont* 1998; 11: 491-501.
43. Huja SS, Litsky AS, Beck FM, Johnson KA, Larsen PE. Pull-out strength of monocortical screws placed in the maxillae and mandibles of dogs. *Am J Orthod Dentofacial Orthop* 2005; 127: 307-13.
44. Okazaki J, Gonda Y, Kamada A, Sakaki T, Kitayama N, Kawamura T, Ueda M. Disaccharide analysis of chondroitin sulfate in peri-implant sulcus fluid from dental implants. *Eur J Oral Sci* 1996; 104: 141-43.
45. Johanson B, Smedberg J, Langley M, Embery G. Glycosaminoglycans in peri-implant sulcus fluid from implants placed in sinus-inlay bone grafts. *Clin Oral Implants Res* 2001; 12: 202-06.
46. Pander N, Samuels RHA, Last KS. The monitoring of orthodontic tooth movement over a 2-year period by analysis of gingival crevicular fluid. *Eur J Oral Sci* 1994; 16: 511-20.
47. Brex MC, Schlegel K, Gehr P, Lang NP. Comparison between histological and clinical parameters during human experimental gingivitis. *J Periodont Res* 1987; 22: 50-57.
48. Baldwin PD, Pander N, Last KS. Effects on tooth movement of force delivery from nickel-titanium archwires. *Eur J Orthod* 1999; 21: 481-89.

49. Kao RT, Curtis DA, Richard DW, Preble J. Increase in interleukine-1B in the crevicular of diseased implants. *Int J Oral Maxillofac Implants* 1995; 10: 969-701.
50. Samuels RHA, Pender N, Last KS. The effects of orthodontic tooth movement on the glycosaminoglycan components of gingival crevicular fluid. *J Clin Periodontol* 1993; 20: 371-77.
51. Shibutani Y, Nishino W, Shiraki M, Iwayama Y. ELISA detection of glycosaminoglycan (GAG)-linked proteoglycans in gingival crevicular fluid. *J Periodont Res* 1993; 28: 17-20.
52. Pothachareon P. The quantitative analysis of chondroitin sulphate epitopes and hyaluronan as diagnosis marker for degenerative joint disease by ELISA technique. Graduated school, Chiang Mai University, Thailand 2000.
53. Tiengburanatum N. Production and characterization of monoclonal antibody against chondroitin-6-sulphate. Graduated school, Chiang Mai University, Thailand 1996.
54. Kongtawelert P. WF6: a CS monoclonal antibody and its application in monitoring joint pathology in vitro and in vivo. Satellite Symposium of IUBMB 2006 Extracellular Glycomatrix in Health and Disease 2006; June;17:92-93.
55. Suzuki Y, Suzuki B. The Suzuki Reef Knot for Attaching Closed-Coil Springs to Miniscrew Heads. *J Clin Orthod* 2007; XLI(3): 148-51.
56. Park HS, Jeong SH, Kwon OW. Factors affecting the clinical success of screw implants used as orthodontic anchorage. *Am J Orthod Dentofacial Orthop* 2006; 130: 18-25.

57. Waddington RJ, Embrey G, Samuels RH. Characterization of proteoglycan metabolites in human gingival crevicular fluid during orthodontic tooth movement. *Arch Oral Biol* 1994; 39: 361-68.
58. Khongkhunthian S, Srimueang N, Krisanaprakornkit S, Pattanaporn K, Ong-Chia S, Kongtawelert P. Raise chondroitin sulphate WF6 epitope levels in gingival crevicular fluid in chronic periodontitis. *J Clin Periodontol* 2008; 35: 871-76.
59. Last KS, Stanbury JB, Embrey G. Glycosaminoglycans in human gingival crevicular fluid as indicators of active periodontal disease. *Arch Oral Biol* 1985; 30: 275-81.
60. Beck CB, Embrey G, Langley MS, Waddington RJ. Levels of glycosaminoglycans in pre-implant sulcus fluid as a means of monitoring bone response to endosseous dental implants. *Clin Oral Implants Res* 1991; 2: 179-85.
61. Smedberg JI, Beck CB, Embrey G. Glycosaminoglycans in peri-implant sulcus fluid from implants supporting fixed or removable prosthesis. *Clin Oral Implants Res* 1993; 4: 137-43.
62. Last KS, Smith S, Pender N. Monitoring of IMZ titanium endosseous dental implants by glycosaminoglycan analysis of peri-implant sulcus fluid. *Int J Oral Maxillofac Implants* 1995; 10: 58-66.
63. Faltin RM, Faltin K, Sander FG, Arana-Chavez VE. Ultrastructure of cementum and periodontal ligament after continuous intrusion in humans: a transmission electron microscopy study. *Eur J Orthod* 2001; 23: 35-49.

64. Embery G, Oliver W, Stanbury JB, Purvis J A. The electrophoretic detection of acid glycosaminoglycans in human gingival sulcus fluid. *Arch Oral Biol* 1982; 27: 177-79.



ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่
Copyright© by Chiang Mai University
All rights reserved