

## TABLE OF CONTENTS

| CONTENTS   | PAGE |
|--|------|
| ACKNOWLEDGEMENTS   | iii  |
| ABSTRACT   | iv   |
| LIST OF TABLES   | x    |
| LIST OF ILLUSTRATIONS  | xi   |
| CHAPTER I INTRODUCTION   | 1    |
| Principle, theory, rationale, and hypothesis   | 1    |
| Anticipated benefits   | 2    |
| CHAPTER II LITERATURE REVIEW   | 3    |
| I) Effect of orthodontic treatment on periodontium   | 3    |
| II) The periodontal tissue (periodontium)  | 5    |
| III) Extracellular matrix (ECM) of the periodontal tissue  | 7    |
| IV) Glycosaminoglycans and chondroitin sulfate in<br>periodontal tissue                                  | 8    |
| V) Gingival crevicular fluid (GCF)   | 10   |
| VI) Glycosaminoglycans (GAGs) in gingival crevicular fluid<br>associated with orthodontic tooth movement | 12   |
| VII) Methods for detecting GAGs in GCF   | 13   |
| Electrophoresis  | 13   |
| Enzyme-linked immunosorbent assay (ELISA)  | 15   |
| VIII) Monoclonal antibodies (mAb) and mAb WF6  | 17   |
| CHAPTER III MATERIALS AND METHODS  | 19   |
| Materials  | 19   |

| CONTENTS  | PAGE      |
|---|-----------|
| Methods   | 20        |
| <b>CHAPTER IV RESULTS</b>   | <b>24</b> |
| Part I: Longitudinal changes of WF6 epitope levels during orthodontic tooth movement  | 24        |
| Part II: Comparison of WF6 epitope levels during the first four weeks of canine movement phase and after complete canine movement phase | 28        |
| <b>CHAPTER V DISCUSSION</b>   | <b>33</b> |
| C-6-S levels in human GCF as a biomarker for alveolar bone resorption during orthodontic tooth movement                                 | 33        |
| A cyclical pattern of C-6-S levels in human GCF collected from teeth undergoing orthodontic tooth movement                              | 35        |
| Limitation of the study   | 37        |
| Suggestions for further study   | 37        |
| <b>CHAPTER VI CONCLUSION</b>  | <b>39</b> |
| <b>BIBLIOGRAPHY</b>   | <b>40</b> |
| <b>APPENDIX</b>   | <b>45</b> |
| Estimation of the volume and weight of GCF samples  | 45        |
| <b>CIRRICULUM VITAE</b>   | <b>54</b> |

## LIST OF TABLES

| TABLES  | PAGE |
|---|------|
| 4.1 All subjects and their teeth in this study  | 24   |
| 4.2 A summary of the medians of WF6 epitope levels (ng/ $\mu$ g total protein) of canines (n=7) and incisors (n=4) for the first four weeks of canine movement phase or M phase             | 29   |
| 4.3 A summary of the medians of WF6 epitope levels (ng/ $\mu$ g total protein) of canines (n=7) and incisors (n=4) for the first four weeks after complete canine movement phase or S phase | 31   |
| A.1 Protein levels, WF6 epitope levels, WF6 epitope levels per unit protein in GCF samples from tooth C11 and C13   | 46   |
| A.2 Protein levels, WF6 epitope levels, WF6 epitope levels per unit protein in GCF samples from tooth D11, D13 ,and D23   | 48   |
| A.3 Protein levels, WF6 epitope levels, WF6 epitope levels per unit protein in GCF samples from tooth E11, E13, and E23   | 50   |
| A.4 Protein levels, WF6 epitope levels, WF6 epitope levels per unit protein in GCF samples from tooth H11, H13, and H23   | 52   |

ลิขสิทธิ์มหาวิทยาลัยเชียงใหม่

Copyright© by Chiang Mai University

All rights reserved

## LIST OF ILLUSTRATIONS

| FIGURES  | PAGE |
|--|------|
| 2.1 Structure of periodontium  | 6    |
| 2.2 Models of typical proteoglycans. NH <sub>2</sub> and COOH termini of core proteins are indicated by N and C, respectively. Glycosaminoglycans are depicted by red lines (i.e., solid line, chondroitin sulfate / dermatan sulfate CS / DS; broken line, heparan sulfate HS; wavy line, keratan sulfate KS) | 8    |
| 2.3 Classification and repeating disaccharide units of glycosaminoglycans  | 9    |
| 2.4 Compositions of gingival crevicular fluid  | 11   |
| 2.5 Separation of molecules after moving through an electric field   | 14   |
| 2.6 ELISA and monoclonal antibody in detecting antigen   | 16   |
| 3.1 Paper strips used to collect GCF   | 21   |
| 4.1 Longitudinal changes in WF6 epitope levels of a canine (ng/ $\mu$ g total protein). The red arrows indicated the time when a new closed coil spring was changed and the force was therefore applied to the canine  | 26   |
| 4.2 Longitudinal changes in WF6 epitope levels of an incisor (ng/ $\mu$ g total protein). The red arrows indicated the time when a new closed coil spring was changed and the force was therefore applied to the canine  | 27   |
| 4.3 A linear graph showed the medians of WF6 epitope levels (ng/ $\mu$ g total protein) of canines (n=7) and incisors (n=4) for the first four weeks of canine movement phase or M phase   | 29   |
| 4.4 A linear graph showed the medians of WF6 epitope levels (ng/ $\mu$ g total protein) of canines (n=7) and incisors (n=4) for the first four weeks after complete canine movement phase or S phase   | 31   |