

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	iii
ABSTRACT (THAI)	iv
ABSTRACT (ENGLISH)	vi
TABLE OF CONTENTS	viii
LIST OF TABLES	xiii
LIST OF FIGURES	xv
CHAPTER 1 INTRODUCTION	1
1.1 Principles, rationale and hypothesis	1
1.2 Research objectives	3
1.3 Research scope	3
1.4 Usefulness of the research	3
1.5 Research locations	3
CHAPTER 2 LITERATURE REVIEW	4
2.1 Introduction	4
2.2 Chemical composition, nutritive and medicinal value	5
2.3 Anatomy of citrus fruit	6
2.4 Postharvest technology	9
2.4.1 Harvest maturity indices	9
2.4.2 Harvesting methods	10
2.4.3 Packinghouse procedures	10
2.4.3.1 Delivery and cleaning	10
2.4.3.2 Washing, waxing, antifungal treatment and sorting	10
2.4.3.3 Sizing and packing	10
2.5 Postharvest physiology	11
2.6 Refrigerated storage of tangerine fruit	13

2.7 Green mold rot of tangerine	15
2.7.1 Disease cycle and epidemiology	15
2.7.2 Symptoms	16
2.7.3 Causal agent	16
2.8 Chilling injury	19
2.8.1 Chilling temperature of tangerine fruit	19
2.8.2 Symptoms of chilling injury	20
2.8.3 Mechanism of chilling injury	21
2.8.3.1 Lipid composition and chilling injury	22
2.8.3.2 Membrane permeability and leakage	23
2.9 Heat treatment	24
2.10 Hot water treatment (HWT)	26
2.11 Heat treatment and host pathogen interactions	27
2.11.1 Pathogen response to heat treatment	27
2.11.2 Fruit responses affecting pathogen defence	29
2.12 Heat treatment and chilling injury	33
CHAPTER 3 MATERIALS AND METHODS	36
3.1 Effect of hot water treatment (HWT) on green mold infection in tangerine fruit cv. Sai Num Pung	36
3.1.1 Effect of HWT on spore germination of <i>Penicillium digitatum</i> <i>in vitro</i>	36
3.1.1.1 Preparation of spore suspension	36
3.1.1.2 Hot water treatments	36
3.1.2 Effect of HWT on infection of <i>Penicillium digitatum</i> in tangerine fruit cv. Sai Num Pung	37
3.1.2.1 Fruit preparation	37
3.1.2.2 Postharvest HWT and storage conditions	37
3.1.2.3 Inoculum preparation	37
3.1.2.4 Hot water treatments	38

3.2 Effect of HWT on anatomy and chemical component changes in tangerine fruit during infection of <i>Penicillium digitatum</i> and chilling injury under low-temperature storage	41
3.2.1 Effect of HWT on infection of <i>Penicillium digitatum</i> in tangerine fruit under low-temperature storage	41
3.2.1.1 Fruit preparation	41
3.2.1.2 Postharvest HWT and storage conditions	41
3.2.1.3 Inoculum preparation	41
3.2.1.4 Hot water treatments	41
3.2.2 Effect of HWT on chemical component changes and chilling injury in tangerine fruit under low-temperature storage	42
3.2.2.1 Fruit preparation	42
3.2.2.2 Chilling injury evaluation	42
3.2.2.3 Determination of electrolyte leakage	43
3.2.2.4 Determination of soluble solid content (SSC)	43
3.2.3 Effect of HWT on anatomy changes of tangerine fruit peel during infection of <i>Penicillium digitatum</i> under low-temperature storage	44
3.2.3.1 Preparation of tangerine fruit peel for SEM	44
3.3 Effect of HWT on biochemical changes in tangerine fruit peel during infection of <i>Penicillium digitatum</i> and chilling injury under low-temperature storage	45
3.3.1 Effect of HWT on activities of the defensive enzymes in tangerine fruit peel during infection of <i>Penicillium digitatum</i> under low-temperature storage	45
3.3.1.1 Extraction of the defensive enzymes	45
3.3.1.2 Enzyme assay and protein determination	45
3.3.2 Effect of HWT on protein patterns in tangerine fruit peel during infection of <i>Penicillium digitatum</i> under low-temperature storage	47
SDS-PAGE (SDS-Polyacrylamide gel electrophoresis) of protein	47
3.3.2.1 Preparation of separating gel	47
3.3.2.2 Preparation of stacking gel	48

3.3.2.3 Running the gel	48
3.3.2.4 Sample preparation	48
3.3.2.5 Staining of gels with Coomassie Brilliant Blue (CBB) R-250	48
3.3.2.6 Rf Value of protein and marker dye	49
3.3.3 Effect of HWT on lipid peroxidation in tangerine fruit peel under low-temperature storage	49
3.3.3.1 Determination of malondialdehyde (MDA) concentration	49
3.3.3.2 Extraction of lipoxygenase	50
3.3.3.3 Enzyme assay and protein determination	50
3.4 Statistical data analysis	50
CHAPTER 4 RESULTS AND DISCUSSION	51
4.1 Effect of hot water treatment (HWT) on green mold infection in tangerine fruit cv. Sai Num Pung	51
4.1.1 Effect of HWT on spore germination of <i>Penicillium digitatum</i> <i>in vitro</i>	51
4.1.2 Effect of HWT on infection of <i>Penicillium digitatum</i> in tangerine fruit cv. Sai Num Pung	60
4.2 Effect of HWT on anatomy and biochemical changes in tangerine fruit during infection of <i>Penicillium digitatum</i> under low-temperature storage	81
4.2.1 Effect of HWT on infection of <i>Penicillium digitatum</i> in tangerine fruit under low-temperature storage	81
4.2.2 Effect of HWT on anatomy changes of tangerine fruit peel during infection of <i>Penicillium digitatum</i> under low-temperature storage	92
4.2.3 Effect of HWT on activities of the defensive enzymes and protein patterns in tangerine fruit peel during infection of <i>Penicillium digitatum</i> under low-temperature storage	104
4.3 Effect of HWT on chemical component changes and chilling injury in tangerine fruit under low-temperature storage	113

CHAPTER 5 CONCLUSIONS	123
5.1 Effect of hot water treatment (HWT) on green mold infection in tangerine fruit cv. Sai Num Pung	123
5.2 Effect of HWT on anatomy and biochemical changes in tangerine fruit during infection of <i>Penicillium digitatum</i> under low-temperature storage	123
5.3 Effect of HWT on chemical component changes and chilling injury in tangerine fruit under low-temperature storage	124
BIBLIOGRAPHY	125
PUBLICATIONS	141
CURRICULUM VITAE	142

LIST OF TABLES

Table	Page
2.1 The chemical compositions and nutritional values of fresh tangerine fruit per 100 gram of the edible part	7
2.2 Storage temperature, relative humidity and storage life of fresh mandarin and tangerine fruit	14
2.3 Approximate lowest safe temperatures, freezing points and characteristic low-temperature injury in citrus fruit	21
2.4 The thermodynamic and operational characteristics of heat treatments	25
2.5 Hot water treatments for citrus fruit, optimal temperature and aim of heat treatments	28
4.1 Effect of hot water treatments (HWT) on green mold rot disease index of artificially-inoculated tangerine fruit during 5 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	65
4.2 Effect of hot water treatments (HWT) on green mold rot disease severity (lesion diameter) of artificially-inoculated tangerine fruit during 5 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	66
4.3 Effect of hot water treatments (HWT) on sporulation of green mold rot disease of artificially-inoculated tangerine fruit during 5 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	67
4.4 Effect of hot water treatments (HWT) on green mold rot disease index of artificially-inoculated tangerine fruit during 35 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	83
4.5 Effect of hot water treatments (HWT) on green mold rot disease severity (lesion diameter) of artificially-inoculated tangerine fruit during 35 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	84

4.6	Effect of hot water treatments (HWT) on chitinase activity in flavedo tissues of artificially-inoculated tangerine fruit during 30 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	106
4.7	Effect of hot water treatments (HWT) on β -1,3-glucanase activity in flavedo tissues of artificially-inoculated tangerine fruit during 30 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	107
4.8	Effect of hot water treatments (HWT) on peroxidase activity in flavedo tissues of artificially-inoculated tangerine fruit during 30 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	109
4.9	Effect of hot water treatments (HWT) at various temperature and time on chilling injury index of tangerine fruit storage at $2\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH for 20, 25 and 30 days	114
4.10	Effect of hot water treatments (HWT) on electrolyte leakage (EL) in flavedo tissues of tangerine fruit during 30 days storage at $2\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	117
4.11	Effect of hot water treatments (HWT) on malondialdehyde (MDA) concentration in flavedo tissues of tangerine fruit during 30 days storage at $2\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	120
4.12	Effect of hot water treatments (HWT) on soluble solids content (SSC) of tangerine fruit during 30 days storage at $2\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	122

LIST OF FIGURES

Figure	Page
2.1 Cross-section of a citrus fruit	8
2.2 Three stages of green mold rot on sweet oranges	17
2.3 <i>Penicillium digitatum</i> colonies on CYA (A) and MEA (B), 7 days at 25°C	17
2.4 <i>Penicillium digitatum</i> (A) penicilli x750 (B) conidia x1875	18
2.5 Postharvest chilling injury of citrus fruit	22
3.1 Scores of the fruit surface infected with <i>Penicillium digitatum</i>	39
3.2 Sporulation index of the fruit surface covered with green mold spores	40
4.1 Effect of hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 24 and 48 hours at 45±2, 50±2 and 55±2°C for 0.5, 1, 2 and 3 minutes	52
4.2 Hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 24 hours, at 45±2°C for 0.5, 1, 2 and 3 minutes (A-D) compared with control (E) showed no effect (100% germination)	53
4.3 Hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 48 hours, at 45±2°C for 0.5, 1, 2 and 3 minutes (A-D) compared with control (E) showed no effect (100% germination)	54
4.4 Hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 24 hours, at 50±2°C for 0.5, 1, 2 and 3 minutes	55
4.5 Hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 48 hours, at 50±2°C	

	for 0.5, 1, 2 and 3 minutes (A-D) compared with control (E) showed no effect (100% germination)	56
4.6	Hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 24 hours, at 55±2°C for 0.5, 1, 2 and 3 minutes	57
4.7	Hot water treatments on spore germination of <i>Penicillium digitatum</i> incubated at 25±2°C in darkness for 48 hours, at 55±2°C for 0.5, 1, 2 and 3 minutes	58
4.8	Disease index of green mold rot of artificially-inoculated tangerine fruit treated with hot water (HWT) before/after inoculation at 45±2, 50±2 and 55±2°C for 0.5, 1, 2 and 3 minutes during 5 days incubation at 24±2°C and 90±5% RH	61
4.9	Disease severity of green mold rot of artificially-inoculated tangerine fruit treated with hot water (HWT) before/after inoculation at 45±2, 50±2 and 55±2°C for 0.5, 1, 2 and 3 minutes during 5 days incubation at 24±2°C and 90±5% RH	62
4.10	Sporulation index of green mold rot of artificially-inoculated tangerine fruit treated with hot water (HWT) before/after inoculation at 45±2, 50±2 and 55±2°C for 0.5, 1, 2 and 3 minutes during 5 days incubation at 24±2°C and 90±5% RH	63
4.11A	Tangerine fruit after inoculated 0 day with <i>Penicillium digitatum</i> and treated with hot water (HWT) at 45±2°C for 0.5, 1, 2 and 3 minutes	68
4.11B	Tangerine fruit after inoculated 0 day with <i>Penicillium digitatum</i> and treated with hot water (HWT) at 50±2°C for 0.5, 1, 2 and 3 minutes	69
4.11C	Tangerine fruit after inoculated 0 day with <i>Penicillium digitatum</i> and treated with hot water (HWT) at 55±2°C for 0.5, 1, 2 and 3 minutes	70
4.11D	Two sets of control uninoculated and untreated fruit 0 day	71
4.12A	Tangerine fruit 3 days incubation at 24±2°C and 90±5% RH after	

- inoculated with *Penicillium digitatum* and treated with hot water (HWT) at $45\pm 2^{\circ}\text{C}$ for 0.5, 1, 2 and 3 minutes; arrows indicate where the symptoms begin to show 72
- 4.12B Tangerine fruit 3 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH after inoculated with *Penicillium digitatum* and treated with hot water (HWT) at $50\pm 2^{\circ}\text{C}$ for 0.5, 1, 2 and 3 minutes; arrows indicate where the symptoms begin to show 73
- 4.12C Tangerine fruit 3 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH after inoculated with *Penicillium digitatum* and treated with hot water (HWT) at $55\pm 2^{\circ}\text{C}$ for 0.5, 1, 2 and 3 minutes; arrows indicate where the symptoms begin to show 74
- 4.12D Two sets of control uninoculated fruit showed no symptom while inoculated and untreated fruit showed symptoms (arrow) 3 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH 75
- 4.13A Tangerine fruit 5 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH after inoculated with *Penicillium digitatum* and treated with hot water (HWT) at $45\pm 2^{\circ}\text{C}$ for 0.5, 1, 2 and 3 minutes; all treatments showed symptoms 76
- 4.13B Tangerine fruit 5 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH after inoculated with *Penicillium digitatum* and treated with hot water (HWT) at $50\pm 2^{\circ}\text{C}$ for 0.5, 1, 2 and 3 minutes; inoculation before HWT at $50\pm 2^{\circ}\text{C}$ for 3 minutes showed best result 77
- 4.13C Tangerine fruit 5 days incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH after inoculated with *Penicillium digitatum* and treated with hot water (HWT) at $55\pm 2^{\circ}\text{C}$ for 0.5, 1, 2 and 3 minutes; inoculation before HWT at $55\pm 2^{\circ}\text{C}$ for 2 and 3 minutes showed best result 78
- 4.13D Two sets of control uninoculated fruit showed no symptom while inoculated and untreated fruit showed serious symptoms 5 days

	incubation at $24\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	79
4.14	Effect of hot water treatments (HWT) on green mold rot disease index and severity (lesion diameter) of artificially-inoculated tangerine fruit during 35 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	82
4.15	Effect of hot water treatments (HWT) at various temperature and time on green mold rot disease development of artificially-inoculated tangerine fruit 10 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	85
4.16	Effect of hot water treatments (HWT) at various temperature and time on green mold rot disease development (percentage of disease index: DI) of artificially-inoculated tangerine fruit 15 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	86
4.17	Effect of hot water treatments (HWT) at various temperature and time on green mold rot disease development (percentage of disease index: DI) of artificially-inoculated tangerine fruit 20 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	87
4.18	Effect of hot water treatments (HWT) at various temperature and time on green mold rot disease development (percentage of disease index: DI) of artificially-inoculated tangerine fruit 25 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	88
4.19	Effect of hot water treatments (HWT) at various temperature and time on green mold rot disease development (percentage of disease index: DI) of artificially-inoculated tangerine fruit 30 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH	89
4.20	SEM showed effect of hot water treatments (HWT) on 'Sai Num Pung' tangerine fruit surfaces after treated with HWT at $50\pm 2^{\circ}\text{C}$ for 3 minutes (A) HWT at $55\pm 2^{\circ}\text{C}$ for 2 minutes (B) and 3 minutes (C) (0 day) the peel was smooth homogeneous but control (D) was rough	93
4.21	SEM showed effect of hot water treatments (HWT) (Day 0) on 'Sai Num Pung' tangerine fruit surfaces after treated with HWT at	

- 55±2°C for 3 minutes having narrow stoma (C) while HWT at 50±2°C for 3 minutes and HWT at 55±2°C for 2 minutes stomata were plugged with melted cuticle (A, B) compared with control stoma was wide opened and unplugged (D) 94
- 4.22 SEM showed effect of hot water treatments (HWT) on crack of the peel light after treated with HWT at 50±2°C for 3 minutes (A) HWT at 55±2°C for 2 minutes (B) and 3 minutes (C) (0 day) showed the cracks were shallow filled with melted cuticle whereas control (D) the crack was wider with a germinating spore (arrow) 95
- 4.23 SEM showed effect of hot water treatments (0 day) on *Penicillium digitatum* spore (Sp) number exist on the peel after treatment at various temperature and time, all treatments (A, B, C) showed small number of spore compared to high number in control (D) 97
- 4.24 SEM showed higher magnification of spore (Sp) on the fruit peel affected by hot water treatments (0 day) at various temperature and time, all treatments (A, B, C) showed denature of spore (ASp = abnormal spore) while the spore in control (D) was normal 98
- 4.25 SEM showed numerous spores (Sp) and mycelium (H = hyphae) of *Penicillium digitatum* on the peel of tangerine fruit (A, B, C) 15 days after inoculation stored at 4±2°C and 90±5% RH without hot water treatment (control) 99
- 4.26 SEM showed few hyphae (H), small number and denature of spore (Sp) of *Penicillium digitatum* on the peel of tangerine fruit affected by hot water treatment at 50±2°C for 3 minutes (A, B, C) 15 days after inoculation stored at 4±2°C and 90±5% RH; ASp = abnormal spore 100
- 4.27 SEM showed small number and denature of hyphae (H) of *Penicillium digitatum* on the peel of tangerine fruit affected by hot

- water treatment at $55\pm 2^{\circ}\text{C}$ for 2 minutes (A, B, C) 15 days after inoculation stored at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH; AH = abnormal hyphae 101
- 4.28 SEM showed few hyphae (H) and denature of spore (Sp) of *Penicillium digitatum* on the peel of tangerine fruit affected by hot water treatment at $55\pm 2^{\circ}\text{C}$ for 3 minutes (A, B, C) 15 days after inoculation stored at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH; ASp = abnormal spore 102
- 4.29 Effect of hot water treatments (HWT) on changes of chitinase, β -1,3-glucanase (beta-1,3-Glucanase) and peroxidase activities in flavedo tissue of tangerine fruit inoculated with *Penicillium digitatum* during 30 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH 105
- 4.30 Protein bands from tangerine fruit peel tissue in response to hot water treatments (HWT) 5 days storage at $4\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH by 10% SDS-PAGE 112
- 4.31 Effect of hot water treatments (HWT) at various temperature and time on chilling injury index of tangerine fruit storage at $2\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH for 20, 25 and 30 days 115
- 4.32 Effect of hot water treatments (HWT) at various temperature and time on chilling injury of tangerine fruit 30 days storage at $2\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH, no chilling injury were found 116
- 4.33 Effect of hot water treatments (HWT) on electrolyte leakage, malondialdehyde (MDA) and soluble solids content of tangerine fruit during 30 days storage at $0\pm 2^{\circ}\text{C}$ and $90\pm 5\%$ RH 118