

TABLE OF CONTENTS

	Page
Acknowledgement	iii
Thai abstract	iv
English abstract	vi
Table of Contents	viii
List of Tables	xii
List of Illustrations	xvi
Chapter 1 Introduction	1
1.1 Principles and Rationale	1
1.2 Study objectives	2
1.3 Usefulness of the research (theoretical and/or applied)	3
1.4 Research plan	3
Chapter 2 Literature review	4
2.1 Lime fruit (<i>Citrus aurantifolia</i> (Christm.) Swingle)	4
2.1.1 Scientific classification	4
2.1.2 General description	5
2.1.3 Constituents and composition	5
2.1.4 Botanical	6
2.1.5 Benefits	6
2.2 Citrus limonoids	8
2.2.1 Limonin and other limonoids	8

	2.2.2 Structure of limonin	9
	2.2.3 Properties of limonin	9
	2.2.4 Biosynthesis of limonoids	10
	2.2.5 Delayed bitterness in citrus juice	13
	2.3 Hydrocolloids	15
	2.3.1 Gum acasia	15
	2.3.2 Pectin	18
	2.3.3 Carboxymethylcellulose (CMC)	21
	2.3.4 Carrageenan	22
	2.4 High Pressure Processing (HPP)	24
	2.4.1 Principles of High Pressure Processing	26
	2.4.2 Engineering Aspects	27
	2.4.3 The advantage of High Pressure Processing	29
Chapter 3	Materials and Methods	30
	3.1 Materials and reagents	30
	3.1.1 Raw materials and hydrocolloids	30
	3.1.2 Standard reagents	30
	3.2 Lime juice sample preparation	30
	3.3 Methodology	31
	3.3.1 Characterization of fresh lime juice	32
	3.3.2 Limonin content of the lime fruit component	34
	3.3.3 Different quality parameters of lime juice during one month storage at 4-6°C and at ambient temperature	34

	3.3.4 The effect of different types and levels of hydrocolloids on the d-limonin content and qualities of lime juice during storage at ambient temperature	36
	3.3.5 The effect of High Pressure Processing (HPP) on the d-limonin content and the quality of lime juices during storage at 4-6°C and at ambient temperature	38
	3.4 Sensory evaluation	40
Chapter 4	Results and Discussion	41
	4.1 Proximate analysis and chemical characteristics of fresh lime juice	41
	4.2 Physical, chemical, nutritional and microbial qualities of fresh lime juice (non-processed lime juice) during storage at 4-6°C and at ambient temperature for 1 month period	43
	4.3 Distribution of d-limonin in lime fruit	53
	4.4 The level of d-limonin in fresh lime juice during one month storage at different storage temperatures	55
	4.5 The effect of hydrocolloids on the physical, chemical, nutritional and d-limonin content of lime juice during one month storage	56
	4.6 The effects of High Pressure Processing (HPP) in the presence and absence of CMC on the physical, chemical, nutritional, microbial, qualities and d-limonin of lime juice during one month storage	74

4.7 The effect of High Pressure Processing in the presence and absence of CMC on the sensory quality of lime juice during one month storage	96
Chapter 5 Conclusion	109
Reference	112
Appendices	
Appendix A Picture of lime juice	127
Appendix B Reagents and Instruments	131
Appendix C Estimation of d-limonin	135
Appendix D Physical, chemical, microbiological and nutritional analyses	139
Appendix E Statistical analysis	148
Curriculum vitae	156

LIST OF TABLES

Table		Page
2.1	Source of commercially important hydrocolloids	16
4.1	The proximate analysis and chemical characteristics of fresh lime juice	41
4.2	Chemical characteristics of fresh lime juice during one month storage at 4-6°C	44
4.3	Chemical characteristics of fresh lime juice during one month storage at ambient temperature	44
4.4	L*-value of lime juices added with different hydrocolloid types at a concentration of 0.5 g/l (w/v) during one month storage at ambient temperature	67
4.5	L*-value of lime juices added with different hydrocolloid types at a concentration of 1.0 g/l (w/v) during one month storage at ambient temperature	68
4.6	L*-value of lime juices added with different hydrocolloid types at a concentration of 1.5 g/l (w/v) during one month storage at ambient temperature	68
4.7	a*-value of lime juices added with different hydrocolloid types at a concentration of 0.5 g/l (w/v) during one month storage at ambient temperature	70

4.8	a*-value of lime juices added with different hydrocolloid types at a concentration of 1.0 g/l (w/v) during one month storage at ambient temperature	71
4.9	a*-value of lime juices added with different hydrocolloid types at a concentration of 1.5 g/l (w/v) during one month storage at ambient temperature	71
4.10	b*-value of lime juices added with different hydrocolloid types at a concentration of 0.5 g/l (w/v) during one month storage at ambient temperature	72
4.11	b*-value of lime juices added with different hydrocolloid types at a concentration of 1.0 g/l (w/v) during one month storage at ambient temperature	72
4.12	b*-value of lime juices added with different hydrocolloid types at a concentration of 1.5 g/l (w/v) during one month storage at ambient temperature	73
4.13	Total Plate Count (CFU/ml) of High Pressure Processing treated-lime juices during one month storage at 4-6°C	79
4.14	Total Plate Count (CFU/ml) of High Pressure Processing treated-lime juices during one month storage at ambient temperature	79
4.15	Yeast and Mold (CFU/ml) of High Pressure Processing treated-lime juices during one month storage at 4-6°C	80
4.16	Yeast and Mold (CFU/ml) of High Pressure Processing treated-lime juices during one month storage at ambient temperature	80

4.17	The effect of HPP on color (L*-value) of CMC-treated and non-treated lime juice during storage at 4-6°C	90
4.18	The effect of HPP on color (L*-value) of CMC-treated and non-treated lime juice during storage at ambient temperature	91
4.19	The effect of HPP on color (a*-value) of CMC-treated and non-treated lime juice during storage at 4-6°C	92
4.20	The effect of HPP on color (a*-value) of CMC-treated and non-treated lime juice during storage at ambient temperature	93
4.21	The effect of HPP on color (b*-value) of CMC-treated and non-treated lime juice during storage at 4-6°C	94
4.22	The effect of HPP on color (b*-value) of CMC-treated and non-treated lime juice during storage at ambient temperature	95
4.23	Effect of HPP on bitterness characteristic of CMC-treated and non-treated lime juice during the storage at 4-6°C	99
4.24	Effect of HPP on bitterness characteristic of CMC-treated and non-treated lime juice during storage at ambient temperature	100
4.25	Effect of HPP on color appearance of CMC-treated and non-treated lime juice during storage at 4-6°C	101
4.26	Effect of HPP on color appearance of CMC-treated and non-treated lime juice during storage at ambient temperature	102
4.27	Effect of HPP on aroma characteristic of CMC-treated and non-treated lime juice during storage at 4-6°C	103
4.28	Effect of HPP on aroma characteristic of CMC-treated and non-treated lime juice during storage at ambient temperature	104

4.29	Effect of HPP on sour taste of CMC-treated and non-treated lime juice during storage at 4-6 ⁰ C	105
4.30	Effect of HPP on sour taste of CMC-treated and non-treated lime juice during storage at ambient temperature	106
4.31	Effect of HPP on overall acceptance appearance of CMC-treated and non-treated lime juice during storage at 4-6 ⁰ C	107
4.32	Effect of HPP on overall acceptance appearance of CMC-treated and non-treated lime juice during storage at ambient temperature	108

LIST OF ILLUSTRATIONS

Figure		Page
2.1	Lime fruits (a) and the half fruits of lime juice (b)	4
2.2	Cross section diagram of lime fruit	7
2.3	Structure of limonin	9
2.4	The proposed biosynthetic pathways of limonin in citrus	11
2.5	Citrus limonoids	12
2.7	Distribution of limonin in a fruit and vegetable part (as $\mu\text{g}/100\text{ mg}$)	13
2.8	Mechanism of delayed bitterness	14
2.9	The nodules exudates of Gum acasia(a) and Gum acasia powder(b)	17
2.10	Possible structure of the carbohydrate component of gum from <i>A. senegal</i>	17
2.11	Pectin from citrus (a), from apple (b) and pectin powder (c)	19
2.12	Structure of pectins	20
2.13	Structure of Carboxymethylcellulose (CMC)	22
2.14	Carrageenan structures	24
2.15	High pressure machines with wire-wound vessels and two pressure transmission systems	28
3.1	A diagram for the production process of lime juice, storage of the juice and analyses of the juice samples (section 3.2, 3.3.1 and 3.3.3)	35

3.2	A diagram for d-limonin content of each lime fruit component and hydrocolloid treated - lime juice (section 3.2 and 3.3.4)	37
3.3	A diagram for hydrocolloid and HPP treated-lime juice during storage (section 3.3.5)	39
4.1	The content of ascorbic acid in fresh lime juices during one month storage at different storage temperatures	45
4.2	The L*-value of fresh lime juices during 1 month at different storage temperatures	47
4.3	The a*-value of fresh lime juices during 1 month at different storage temperatures	48
4.4	The b*-value of fresh lime juices during 1 month at different storage temperatures	50
4.5	Total plate count and the count of yeast and mold of fresh lime juice during one month storage at different storage temperatures	52
4.6	The distribution of d-limonin in lime fruits	53
4.7	Cross-section of lime fruit, showing the d-limonin distribution in each lime fruit part	54
4.8	The d-limonin content of fresh lime juices during one month storage at different storage temperatures	55
4.9	The d-limonin content of lime juices added with 0.5 g/l (w/v) hydrocolloids during one month storage at ambient temperature	60
4.10	The d-limonin content of lime juices added with 1.0 g/l (w/v) hydrocolloids during one month storage at ambient temperature	60

4.11	The d-limonin content of lime juices added with 1.5 g/l (w/v) hydrocolloids during one month storage at ambient temperature	61
4.12	Total acidity (% citric acid) of lime juices added with different types and levels of hydrocolloids during one month storage at ambient temperature	63
4.13	Total soluble solid (⁰ Brix) of lime juices added with different types and levels of hydrocolloids during one month storage at ambient temperature	64
4.14	pH value of lime juices added with different types and levels of hydrocolloids during one month storage at ambient temperature	64
4.15	Ascorbic acid of lime juices added with different types and levels of hydrocolloids during one month storage at ambient temperature	66
4.16	The d-limonin content (ppm) of HPP treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at 4-6°C	76
4.17	The d-limonin content (ppm) of HPP treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at ambient temperature	76
4.18	The ascorbic acid (mg/100 ml) of High Pressure Processing treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at 4-6°C	82
4.19	The ascorbic acid (mg/100 ml) of High Pressure Processing treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at ambient temperature	83

4.20	pH of High Pressure Processing treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at 4-6°C	84
4.21	pH of High Pressure Processing treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at ambient temperature	85
4.22	Total acidity (% citric acid) of High Pressure Processing treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at 4-6°C	87
4.23	Total acidity (% citric acid) of High Pressure Processing treated-lime juices in the presence and absence of 1.0 g/l (w/v) CMC during one month storage at ambient temperature	87